

# PLANKTONIC COELENTERATES COLLECTED IN THE NORTH ATLANTIC OCEAN

by

R. W. M. VAN SOEST

*Institute of Taxonomic Zoology, University of Amsterdam, The Netherlands*

## ABSTRACT

This short faunistic note contains a report on the species of Scyphozoa, Hydromedusae, Siphonophora and Ctenophora that were collected by OWS "Cirrus" and "Cumulus", R.V. "Tridens" and HMS "Snellius" in the North Atlantic Ocean between 22° N and 66° N. In all, 61 species have been captured. New records for *Calyropsis tya* Fewkes, 1882, *Cunina duplicata* Maas, 1893, *Sibogita geometrica* Maas, 1905 ssp. *occidentalis* Kramp, 1959, *Apolemia uvaria* (Lesueur, 1811), *Thalassophyes crystallina* Moser, 1925 and *Crystallophyes amygdalina* Moser, 1925, are given.

## INTRODUCTION

Although the north-eastern Atlantic Ocean is one of the most thoroughly investigated oceanic areas, the knowledge of the distribution of the oceanic plankton species, particularly the meso- and bathypelagic ones, is by no means complete. New species are still to be described and new records are frequently reported. It seems thus justified to publish a faunistic note on the coelenterates that were collected by Dutch government vessels during the years 1963-1972. Most of the material presented here was collected during the various cruises of the Ocean Weather Ships "Cirrus" and "Cumulus" at five more or less fixed stations: M (66° N, 01° E), A (62° N, 33° W), I (59° N, 19° W), J (52° N, 20° W) and K (45° N, 16° W). At these stations the ships were moving over an area the extent of which was at most one degree latitude and longitude. These "moving" stations were frequently sampled during the period of 1963-1972. Additional material was collected by HMS "Snellius" during 1964/1965 in the area between the west coast of Africa and Bermuda. Finally an important part of the material was collected by the Fisheries Investigation Vessel RV "Tridens" during a cruise to and from the Azores;

this cruise was attended by the author. All stations from which samples were studied are mentioned in table I and fig. 1.

I am much indebted to Mr. C. L. Bekkering, Mr. S. D. Koning and Mr. H. A. J. Donkers of the "Cirrus" and "Cumulus" and to the commanding officer and the crew of HMS "Snellius" for supplying the plankton samples, and to captain A. Krijgsman and his crew of the "Tridens" for their skill and cooperation in obtaining the Azores-plankton. Dr. C. L. Deelder of the Rijksinstituut voor Visserijonderzoek (RIVO) at IJmuiden was very kind in providing me with the opportunity to join the "Tridens" cruise. Dr. P. Pugh (National Institute of Oceanography, Great Britain) kindly identified some species of physonect siphonophores.

For the identification of the species the studies of Kramp (1959), Totton (1954, 1965), Bigelow (1911), Sears (1953), Moser (1925), and Russell (1953, 1970) were used.

## RESULTS

Tables II, III and IV contain a list of all species that were found in the plankton samples mentioned in table I. The stations (latitude only) at which they occurred are indicated by a cross. In this way tables II, III and IV give a latitudinal distribution pattern of the species as it appeared from the samples. This does not necessarily mean that the species do not occur outside the area indicated in the tables, as absence of a species from a sample does not prove that the species is absent in the area the sample is supposed to represent. In fact, many species of oceanic coelenterates seem to be quite rare judging from the number of records.

Table I. List of oceanic stations from which samples were studied.

Station	Position	Date	Time	Depth
CIRRUS M	66° N, 01° E *)	IX-1964	—	Several deep hauls
"	"	VII-1964	—	" " "
CIRRUS A	62° N, 33° W *)	VIII-IX-1963	—	" " "
"	"	II/III-1964	—	" " "
"	"	III-1965	—	" " "
"	"	X-1965	—	" " "
"	"	IV/V-1966	—	" " "
"	"	VI/VII-1968	—	" " "
"	"	V-1972	—	" " "
CIRRUS I	59° N, 19° W *)	XI/XII-1963	—	" " "
"	"	VI-1964	—	" " "
"	"	VIII-1965	—	" " "
"	"	I-1966	—	" " "
CIRRUS J	52° N, 20° W *)	I-1964	—	" " "
"	"	II-1966	—	" " "
CIRRUS K	45° N, 16° W *)	X-1963	—	" " "
"	"	IV/V-1964	—	" " "
"	"	XI-1965	—	" " "
"	"	V/VI-1966	—	" " "
"	"	III/IV-1970	—	" " "
"	"	I-1972	—	" " "
"	"	VI-1972	—	" " "
TRIDENS 4	43°30' N, 09°44' W	30-V-1972	08.30-11.00	350-0 m
TRIDENS 5	42°23' N, 12°38' W	30/31-V-1972	22.15-00.20	150-0 m
TRIDENS 7	39°15' N, 21°25' W	1-VI-1972	13.30-16.30	700-0 m
TRIDENS 8	38°58' N, 22°37' W	1-VI-1972	22.00-24.00	150-0 m
TRIDENS 9	39°26' N, 23°22' W	6-VI-1972	21.00-24.00	150-0 m
TRIDENS 10	42°30' N, 19°30' W	7-VI-1972	21.00-24.00	150-0 m
TRIDENS 11	46°25' N, 15°00' W	8-VI-1972	21.00-24.00	150-0 m
TRIDENS 12	49°00' N, 11°27' W	9-VI-1972	19.30-23.00	300-0 m
SNELLIUS J 3	36°44' N, 55°00' W	17-V-1965	21.00	—
SNELLIUS J 4	37°26' N, 38°46' W	20-V-1965	—	—
SNELLIUS J 5	37°15' N, 27°40' W	22-V-1965	—	2400 m wire out
SNELLIUS J 5a	37°15' N, 27°40' W	23-V-1965	18.00	—
SNELLIUS I 1	33°53' N, 12°55' W	9-IV-1965	22.00	—
SNELLIUS I 1a	33°53' N, 12°55' W	10-IV-1965	10.45	—
SNELLIUS I 2	33°52' N, 19°20' W	11-IV-1965	16.30	—
SNELLIUS I 2a	33°52' N, 19°20' W	12-IV-1965	07.00	50-0 m
SNELLIUS I 3	34°06' N, 30°08' W	14-IV-1965	11.45	—
SNELLIUS I 4	34°09' N, 39°03' W	16-IV-1965	19.30	—
SNELLIUS Navado F-1	25°12' N, 70°23' W	16-I-1965	21.45	10 m
SNELLIUS Navado F-7	25°11' N, 19°35' W	31-I-1965	20.00	110 m
SNELLIUS Echo 5	22°01' N, 49°55' W	13-XII-1964	18.50	100 m

\*) Note: The "Cirrus" and "Cumulus" stations (both indicated as CIRRUS in this table) are "moving" stations. The above listed positions are approximated. Deviations of these listed positions are small, not exceeding one degree latitude or longitude.

Common or fairly common species in the area west of the British Isles are *Aeginura grimaldi* Maas, 1904, *Aglantha digitale* (O. F. Müller, 1776), *Botrynema brucei* Browne, 1908, *Halicreas minimum* Fewkes, 1882, *Pantachogon haeckeli* Maas, 1893, *Atolla wyvillei* Haeckel, 1880, *Periphylla periphylla* (Péron & Lesueur, 1809), *Lenisia conoidea* (Kefferstein & Ehlers, 1860), *Chu-*

*niphyes multidentata* Lens & van Riemsdijk, 1908, and *Beroe cucumis* Fabricius, 1780. In the samples from north of 60° N, *Dimophyes arctica* Chun, 1897 and *Crossota rufobrunnea* (Kramp, 1913), were common. South of 50° N, *Hippopodius hippopus* (Forskål, 1776), *Rosacea plicata* Quoy & Gaimard, 1827, *Vogtia spinosa* Kefferstein & Ehlers, 1861, *Chelophyes appendiculata* (Eschscholtz,




Fig. 1. Map of stations from which samples were studied. Circumlined dots are "moving" stations (see text).

1829), *Eudoxoides spiralis* (Bigelow, 1911) and *Pelagia noctiluca* (Forskål, 1775), were frequently met. South of 35° N, *Liriope tetraphylla* (Chamisso & Eyssenhardt, 1821), *Aglaura hemistoma* Péron & Lesueur, 1809, *Bassia bassensis* (Quoy & Gaimard, 1833) and *Diphyes dispar* Chamisso & Eyssenhardt, 1821 were the dominating species.

*New or interesting records:*

*Calycopsis tya* Fewkes, 1882 (fig. 2A), has originally been described from waters over the

continental shelf off the east coast of North America. Bigelow (e.g. 1922) collected specimens in the same area. Kramp (1959) found several specimens near the Cape Verde Islands. These are the only certain records of this species so far, although possibly some West Indian specimens found by Kramp (l.c.) could in his opinion also belong to this species. RV "Tridens" collected one specimen north-east of São Miguel (one of the Azores) (station 7: 39° 15' N, 21° 25' W). This new record links the records of Fewkes (1882), Bigelow (l.c.) and Kramp (l.c.).

*Cunina duplicata* Maas, 1893. The northernmost

Table II. List of species of siphonophores collected in the North Atlantic Ocean. The stations at which the species occurred (latitudinal position only) are indicated by crosses. For the exact positions of the stations see table I.

	66°	62°	59°	52°	49°	46°	45°	43°	42°	39°	34°	33°	32°	25°	22°	N
<i>SIPHONOPHORA</i>																
<i>Thalassophyes crystallina</i>			x													
<i>Muggiaea kochi</i>			x													
<i>Dimophyes arctica</i>		x	x													
<i>Crystallophyes amygdalina</i>			x				x									
<i>Chuniphyes multidentata</i>			x	x	x		x				x					
<i>Apolemia uvaria</i>			x			x	x									
<i>Nanomia cara</i>			x	x												
<i>Physophora hydrostatica</i>			x	x										x		
<i>Lensia conoidea</i>			x	x				x	x	x	x					
<i>Lensia multicristata</i>				x				x								
<i>Lensia fowleri</i>					x			x								
<i>Rosacea plicata</i>					x	x	x	x		x	x					
<i>Vogtia pentacantha</i>				x			x	x						x		
<i>Vogtia spinosa</i>					x	x	x	x	x	x	x					
<i>Hippopodius hippopus</i>					x		x	x	x	x	x					x
<i>Velella velella</i>					x		x									
<i>Eudoxoides spiralis</i>					x	x	x	x	x	x	x					
<i>Chelophyes appendiculata</i>							x				x	x				x
<i>Halistemma rubrum</i>						x			x	x	x					
<i>Nectopyramis thetis</i>							x				x					
<i>Nectopyramis diomedae</i>							x									
<i>Vogtia serrata</i>											x					
<i>Rosacea cymbiformis</i>											x	x				
<i>Lensia subtilis</i>												x	x			
<i>Diphyes dispar</i>											x	x	x		x	x
<i>Diphyes bojani</i>												x	x			
<i>Ceratocymba sagittata</i>											x	x				
<i>Bassia bassensis</i>												x	x			
<i>Abylopsis tetragona</i>												x				x
<i>Abylopsis eschscholtzi</i>												x	x			
<i>Eudoxoides mitra</i>													x			
<i>Sulculeolaria monoica</i>																x

record of Kramp (1959) in the Atlantic Ocean is 37° 25' N, 41° 00' W. The one rather damaged specimen collected by RV "Tridens" north east of São Miguel (39° 15' N, 21° 25' W) represents a slight northward extension of the known range of this species.

*Pandea conica* (Quoy & Gaimard, 1827) (fig. 2B). One specimen was collected by RV "Tridens" at 38° 58' N, 22° 37' W (st. 8). It is an interesting specimen because it lacks the characteristic apical process, just as the specimen described by Bigelow (1918) from the Bermuda area. It has 32 marginal tentacles, which is in accordance with the number found in specimens off Bermuda (Bigelow, 1938). Its height is 20 mm, its width 14 mm, which makes it a fairly large specimen because of the absence of the apical projection. *P. conica* is common in the Mediterranean, but has been found in the Atlantic near Bermuda, along the west coast of Africa

near Patagonia and in the Azores area (Kramp, 1959). It also occurs in the Pacific.

*Sibogita geometrica* Maas, 1905 ssp. *occidentalis* Kramp, 1959 (fig. 2C). This rare form has been found at three locations in the Atlantic Ocean by Kramp (1959): 36° 41' N, 26° 21' W (one specimen); 39° 26' N, 21° 51' W (one specimen); and 47° 04' N, 07° 08' W (one specimen). RV "Tridens" collected one specimen at station 9 (39° 26' N, 23° 22' W), quite near one of the locations of Kramp (l.c.). Another specimen must be mentioned. It was not collected by one of the research vessels of which the material is at present reported upon, but taken by MS "Walther Herwig" near the equator (St. WH 471-II-71, 02° 29' S, 18° 58' W, 10-IV-1971, 19.50-20.50 h, 300-304 m). This brings the total number of specimens known of this subspecies to five.

*Neoturris pileata* (Forskål, 1775) (fig. 2D). Two specimens were collected by RV "Tridens" at

Table III. List of species of Hydromedusae collected in the North Atlantic Ocean. The stations at which the species occurred (latitudinal position only) are indicated by crosses. For the exact positions of the stations see table I.

	66°	62°	59°	52°	49°	46°	45°	43°	42°	39°	34°	33°	32°	25°	22° N
<b>HYDROMEDUSAE</b>															
<i>Crossota rufobrunnea</i>	x	x													
<i>Botrynema brucei</i>		x	x	x			x								
<i>Aglantha digitale</i>		x	x	x	x		x								
<i>Aeginura grimaldi</i>		x	x				x						x		
<i>Aeginura grimaldi</i>			x												
<i>Tiaranna rotunda</i>		x	x	x			x						x		
<i>Haliscera bigelowi</i>		x	x	x			x								
<i>Pantachogon haeckeli</i>		x	x	x			x								
<i>Neoturris pileata</i>									x						
<i>Rhopalonema funerarium</i>							x								
<i>Solmissus incisa</i> (fragm.)							x		x						
<i>Colobonema sericeum</i>							x								x
<i>Aegina citrea</i>						x	x		x						x
<i>Calyropsis typa</i>															x
<i>Cunina duplicata</i>															x
<i>Pandea conica</i>															x
<i>Sibogita geometrica</i> ssp. <i>occidentalis</i>															x
<i>Rhopalonema velatum</i>															x
<i>Liriope tetraphylla</i>														x	x
<i>Cunina frugifera</i>															x
<i>Aglaura hemistoma</i>															x

Table IV. List of species of Scyphomedusae and Ctenophora collected in the North Atlantic Ocean. The stations at which the species occurred (latitudinal positions only) are indicated by crosses. For the exact positions of the stations, see table I.


	66°	62°	59°	52°	49°	46°	45°	43°	42°	39°	34°	33°	32°	25°	22° N
<b>SCYPHOMEDUSAE</b>															
<i>Atolla wyvillei</i>	x	x	x	x			x								x
<i>Periphylla periphylla</i>		x	x	x	x	x									x
<i>Nausithoe globifera</i>		x					x								
<i>Atolla vanhoeffeni</i>					x		x								
<i>Atolla parva</i>							x								x
<i>Pelagia noctiluca</i>				x			x								x
<b>CTENOPHORA</b>															
<i>Beroe cucumis</i>				x		x	x	x	x	x					
<i>Hormiphora plumosa</i>														x	x

43° 30' N, 09° 44' W ( st. 4). The number of marginal tentacles is 53 and 62 respectively for both specimens. The apical projection is weakly developed.

*Apolemia uvaria* (Lesueur, 1811) (Fig. 2G). Although considered as a Mediterranean siphonophore this species was found by Fraser (1961) in the northern North Sea and the Atlantic waters west of Scotland. RV "Tridens" collected several fine nectophores at 49° 00' N, 11° 27' W (st. 12). OWS "Cirrus" collected 2 nectophores at

45° N, 16° W, and 1 nectophore at 62° N, 33° W. The known range of this species is thus considerably extended. The fact that a specimen of this species was found at 62° N, 33° W in the month of March (1965) supports Fraser's (1967) assumption that the species may survive the boreal winter and is not a seasonal immigrant.

*Thalassophyes crystallina* Moser, 1925. One gonophore and one anterior nectophore of this species were collected by OWS "Cirrus" at 62° N, 33° W, thus confirming Fraser's (1961) records



of occurrence of this species in the north-eastern Atlantic.

*Crystallophyes amygdalina* Moser, 1925 (fig. 2E). Several anterior and posterior nectophores belonging to this species were collected by OWS "Cirrus" at 62° N, 33° W. One anterior nectophore was collected at 45° N, 16° W. The species was hitherto only known from Antarctic waters and from the Subantarctic Atlantic and Indian Ocean and from one location in the North Atlantic (10°N) (Moser, 1925; Leloup, 1934; Totton, 1954). It is not the only species of siphonophore with a disrupt distribution in the North Atlantic as well as in the Southern Oceans (e.g. *Muggiaea bargmannae* Totton, 1954).

*Diphyes dispar* Chamisso & Eyssenhardt, 1821 (fig. 2F). One gigantic specimen of this species

collected by RV "Tridens" at st. 8 (38° 58' N, 22° 37' W) must be mentioned. Anterior and posterior nectophore of the polygastric phase each measure 40 mm in length, which makes it one of the biggest, if not the biggest specimen ever recorded of this well known and common species.

*Dimophyes arctica* (Chun, 1897). Quite a few specimens, both of the polygastric and the eudoxid phase, have been found in various samples taken by OWS "Cirrus" at 62° N, 33° W and 66° N, 01° E. Although Fraser (1961, 1967) considers it to be no longer an exclusively cold water form, since it has been found frequently in the central North Atlantic and even in tropical waters (e.g. Leloup, 1934; Alvarino, 1971), it was absent from all the samples studied at present except for the ones mentioned above.

#### BIBLIOGRAPHY

- ALVARINO, A., 1971. Siphonophores of the Pacific, with a review of the world distribution. *Bull. Scripps Inst. Oceanogr.*, **16**: 1—432.
- BIGELOW, H. B., 1911. Report on the scientific results of the expedition to the Eastern Tropical Pacific, U.S. Fish Commission Steamer "Albatross", 23. The Siphonophorae. *Mem. Mus. comp. Zool. Harv.*, **38** (2): 171—401.
- , 1918. Some medusae and siphonophorae from the Western Atlantic. *Bull. Mus. comp. Zool. Harv.*, **52** (8): 365—442.
- , 1922. Exploration of the coastal waters off the north-eastern United States in 1916 by the US Fisheries schooner "Grampus". *Bull. Mus. comp. Zool. Harv.*, **65**: 85—188.
- , 1938. Plankton of the Bermuda Oceanographic Expeditions, 8. Medusae taken during the years 1929 and 1930. *Zoologica*, New York, **23**: 99—189.
- FEWKES, J. W., 1882. On the Acalephæ of the east coast of New England. *Bull. Mus. comp. Zool. Harv.*, **9** (8): 291—310.
- FRASER, J. H., 1961. The oceanic and bathypelagic plankton of the north-east Atlantic. *Mar. Res.*, **1961** (4): 1—48.
- , 1967. Siphonophora in the plankton to the North and West of the British Isles. *Proc. roy. Soc. Edinburgh*, (B) **70** (1): 1—30.
- KRAMP, P. L., 1959. The Hydromedusae of the Atlantic Ocean and adjacent waters. *Dana Rep.*, **46**: 1—283.
- LELOUP, E., 1934. Siphonophores calycophorides de l'Océan Atlantique tropical et austral. *Bull. Mus. roy. Hist. nat. Belgique*, **10** (6): 1—64.
- MOSER, F., 1925. Die Siphonophoren der Deutschen Südpolar Expedition 1901-1903. *Deutsch. Südpolarexp.*, **17** (Zool. 9): 1—541.
- RUSSELL, F. S., 1953. The Medusae of the British Isles, **1**: 1—530, 35 pls. (University Press, Cambridge).
- , 1970. The Medusae of the British Isles, **2**: 1—284, 15 pls. (University Press, Cambridge).
- SEARS, M., 1953. Notes on Siphonophores 2. A revision of the Abylinae. *Bull. Mus. comp. Zool. Harv.*, **109** (1): 1—119.
- TOTTON, A. K., 1954. Siphonophora of the Indian Ocean. *Discovery Rep.*, **27**: 1—162.
- , 1965. A synopsis of the Siphonophora: 1—230, 40 pls. (Trustees British Museum, London).

Received: 11 January 1973

Fig. 2. A. *Calycopsis typa* Fewkes, 1882, from 39° 15' N, 21° 25' W (×2.6).

B. *Pandea conica* (Quoy & Gaimard, 1827) from 38° 28' N, 22° 37' W (×3.0).

C. *Sibogita geometrica* Maas, 1905 ssp. *occidentalis* Kramp, 1959, from 39° 26' N, 21° 51' W (×2.4).

D. *Neoturris pileata* (Forskål, 1775) from 43° 30' N, 09° 44' W (×2.4).

E. *Crystallophyes amygdalina* Moser, 1925, anterior nectophore from 45° 00' N, 16° 00' W (×16).

F. A very large specimen of *Diphyes dispar* Chamisso & Eyssenhardt, 1821 (polygastric phase) from 38° 58' N, 22° 37' W (×1.3).

G. *Apolemia uvaria* (Lesueur, 1811), three nectophores from 49° 00' N, 11° 27' W (×1.5).

(Photographs made by Mr. L. A. van der Laan).

### Errata

Bijdragen tot de Dierkunde vol. 43 no. 1 (1973)

R. W. M. van Soest, Planktonic coelenterates collected in the North Atlantic Ocean, page 123, table III, HYDROMEDUSAE

5th line - for *Aeginura grimaldi* read: *Tiaranna rotunda*

6th line - for *Tiaranna rotunda* read: *Halicreas minimum*.