

THE ASCOMYCETE GENUS SORDARIA

J. GUARRO* & J. A. VON ARX**

Sordaria is restricted to coprophilous, soil-, or seed-borne Pyrenomycetes with aseptate, elongate ascospores with a gelatinous, amorphous sheath. The genus is redescribed and a key to fourteen accepted species is given. A checklist of all taxa described as *Sordaria* is added.

In the course of a revision of cultures of *Sordaria* species maintained in the CBS Culture Collection, a checklist of all *Sordaria* species was compiled. The genus *Sordaria* is restricted to species with ostiolate ascomata, cylindrical asci and aseptate, elongate, cylindrical, ellipsoidal, or fusiform ascospores with a distinct gelatinous sheath. Species with ovate, broadly fusiform, or nearly spherical ascospores without sheath are classified in a separate genus *Asordaria* (von Arx & al., 1987).

Treatments on *Sordaria* have been published by Bayer (1924) for Central Europe, Cain (1934) for Ontario, Canada, Moreau (1953) for Europe, Boedijn (1962) for Indonesia, Cailleux (1972) for central parts of Africa, Lundqvist (1972) for nordic regions and by Barrasa & al. (1986) for Spain.

In this paper a new description of *Sordaria* sensu stricto is given, followed by a key to fourteen accepted species and a list of all species that have been referred to *Sordaria*.

S O R D A R I A Ces. & De Not.

Sordaria Ces. & De Not. in Comm. Soc. Crit. Ital. 1: 226. 1863. — Type species (lectotype): *S. fimicola* (Rob.) Ces. & De Not.

For synonyms see Lundqvist (1972).

Colonies expanding, dark, often with a pale or brown aerial mycelium, reverse brown or black; ascomata superficial, often embedded in the aerial mycelium, obovate, pyriform, or ampulliform, with a conical or cylindrical ostiolar beak; ascromatal wall pigmented, brown or nearly black, composed of 3–5 layers of rather large, slightly flattened cells, *textura angularis* in surface view, smooth or covered with pigmented hyphal appendages; asci cylindrical, with a short stalk, unitunicate, at the truncate apex with distinct, non amyloid, often ring-like thickenings, 8-spored; ascospores uniseriate, ellipsoidal, cylindrical, or fusiform with rounded ends, aseptate, brown or opaque when mature, with a distinct, often protuberant germ pore at base, with an amorphous, hyaline gelatinous sheath surrounding the spore except for the basal, porate part.

Coprophilous, soil- or seed-borne or isolated from plant debris.

Homothallic or heterothallic.

* Dept. of Biology, Fac. of Medicine, University of Barcelona, Reus, Spain.

** Centraalbureau voor Schimmelcultures, Baarn, The Netherlands.

The heterothallic species can be recognized by a more abundant, often pale aerial mycelium, on which aseptate, hyaline micro-conidia (spermatia) are formed by budding.

KEY TO THE SPECIES

1 a. Ascospores with a length/width ratio of 1.4–1.7	2
b. Ascospores with a length/width ratio of 1.8–3	9
2 a. Ascospores 15–24 × 9–13 µm	3
b. Ascospores broader	5
3 a. Homothallic; spermatia absent; ascomata pyriform, usually smooth	<i>S. fimicola</i>
b. Heterothallic; spermatia (aseptate, small conidia) present	4
4 a. Ascomata with a long beak, tomentose	<i>S. tomento-alba</i>
b. Ascomata pyriform, with a broad, short beak	<i>S. brevicollis</i> <i>S. sclerogenia</i> <i>S. thermophila</i>
5 a. Ascospores 18–23 × 12–15 µm, ellipsoidal-obovate	<i>S. lappae</i>
b. Ascospores larger	6
6 a. Ascospores 24–30 × 14–17 µm	<i>S. superba</i>
b. Ascospores 28–36 × 18–25 µm	7
7 a. Ascospores 22–25 µm broad	<i>S. capturae</i>
b. Ascospores 18–21 µm broad	8
8 a. Homothallic, spermatia absent	<i>S. macrospora</i>
b. Heterothallic, spermatia present	<i>S. heterothallis</i>
9 a. Ascospores with a rounded apex and an attenuated base	10
b. Ascospores attenuated at both ends, fusiform	11
10 a. Ascospores 21–26 × 9–12 µm, with a non-protuberant germ pore	<i>S. alcina</i>
b. Ascospores 25–29 × 12–14 µm, with a protuberant germ pore	<i>S. papillata</i>
11 a. Ascospores 25–31 × 12–14 µm	<i>S. baltica</i>
b. Ascospores 32–40 × 12–15 µm	<i>S. elongatispora</i>

CHECKLIST

List of species that, at one time or another, have been referred to *Sordaria* (*S.*).
Names accepted in *Sordaria* Ces. & De Not., emend. Guarro & v. Arx are printed
space d.

- S. alicina* Lundq. in Symb. bot. upsal. 20: 326. 1972.
S. allahabadensis Srivastava & al. in Mycopath. Mycol. appl. 30: 203. 1966 = *Podospora austro-americana* (Speg.) Mirza & Cain.
S. aloides Fuckel in Jb. nassau. Ver. Naturk. 27–28: 43. 1873 = *Podospora aloides* Fuckel (Mirza & Cain, 1969).
S. alpina Griff. in Mem. Torrey bot. Club 11: 49. 1901. — The type could not be studied.
S. amficornis Sacc., Syll. Fung. 1: 235. 1887 = *Podospora appendiculata* (Auersw.) Niessl (Mirza & Cain, 1969).
S. amphispheroides Ell. & Ev. in Am. Nat. 31: 340. 1897 = *Hypocopra amphispheroides* (Ell. & Ev.) Griff.

S. anomala (Griff.) Sacc. & D. Sacc. in Syll. Fung. 17: 603. 1905 = *Podospora anomala* (Griff.) Cain (Mirza & Cain, 1969).

S. anserina Wint. (Winter, 1873: 100) = *Podospora anserina* Niessl.

S. apiculifera Speg. in An. Mus. nac. Hist. nat. B. Aires 6: 251. 1899 = *Podospora apiculifera* (Speg.) Mirza & Cain (Mirza & Cain, 1969) or = *P. australis* (Speg.) Niessl (Lundqvist, 1972).

S. appendiculata Auersw. in Niessl in Verh. naturf. Ver. Brünn 10: 188. 1872 = *Podospora appendiculata* Auersw. ex Niessl (Mirza & Cain, 1969).

S. arachnoidea (Niessl) Sacc. & Syd. in Syll. Fung. 14: 492. 1899 = *Cercophora coprophila* (Fr.) Lundq. (Lundqvist, 1972).

S. araneosa Cain in Univ. Toronto Stud. (Biol. Ser.) 38: 52. 1934 = *Podospora araneosa* (Cain) Cain.

S. arctica Cain in Can. J. Bot. 35: 262. 1957 = *Asordaria arctica* (Cain) v. Arx & Guarro.

S. arenicola Grove in J. Bot. 66. 1930. — The type could not be studied.

S. argentina Speg. in An. Soc. ci. argent. 10: 16. 1880 = *S. fimicola* (Lundqvist, 1972).

S. argentinensis Speg. in An. Mus. nac. B. Aires 23: 49. 1912 = *Podospora argentinensis* (Speg.) Mirza & Cain (Mirza & Cain, 1969).

S. arizonensis (Griff.) Sacc. in Syll. Fung. 17: 601. 1905 = *Arnium arizone* Lundq. & Krug (Lundqvist, 1972).

S. australis (Speg.) Sacc., Syll. Fung. 1: 239. 1882 = *Podospora australis* (Speg.) Niessl (Mirza & Cain, 1969).

S. austro-americana (Speg.) Sacc., Syll. Fung. 1: 237. 1882 = *Podospora austro-americana* (Speg.) Mirza & Cain (Mirza & Cain, 1969).

S. baltica Lundq. in Symb. bot. upsal. 20: 238. 1972.

S. barbata E.C. Hans. in Vid. Meddr dansk naturh. Foren. 1876: 21. 1877 = *Arnium macrotheca* (Crouan) Lundq. (Lundqvist, 1972).

S. bicurvata Rick in Broteria 2: 135. 1933.

S. bisporula (Crouan) Cooke & Plowr. in Grevillea 7: 86. 1879 = *Trichodelitschia bisporula* (Crouan) Lundq. (Lundqvist, 1964a).

S. bolbitoni (Quél.) Wint. in Rabenh. Kryptog.-Fl. (2. Aufl.) 1, Pilze 2: 169. 1885.

— This is a doubtful species (Bayer, 1924).

S. bombardioides Auersw. ex Niessl in Verh. naturf. Ver. Brünn 10: 187. 1871 = *Bombardioidea bombardioides* (Auersw. ex Niessl) C. Moreau.

S. bosensis Das in Trans. Br. mycol. Soc. 45: 547. 1962 = *Areolospora bosensis* (Das) Hawksw. (Hawksworth, 1980).

S. botryosa Penz. & Sacc., Icon. Fung. Javan.: 9. 1904 = *Bombardia bombarda* (Batsch) J. Schroet. (Moreau, 1953).

S. bovilla (Cooke) Cooke in Grevillea 16: 55. 1887 = *Cercophora coprophila* (Fr.) Lundq. (Lundqvist, 1972).

S. brassicae (Kl.) Crouan, Fl. Finistère: 22. 1867 = *Arnium olerum* (Fr.) Lundq. & Krug apud Lundq. (Lundqvist, 1972).

S. brefeldii Zopf in Z. Naturw. 56: 539. 1883 = *Hypocopra brefeldii* (Zopf) Chen.

S. brevicaudata Speg. in An. Mus. nac. B. Aires 6: 252. 1899. — This is a nomen dubium (Lundqvist, 1972).

S. brevicollis Olive & Fantini in Am. J. Bot. 48: 124. 1961. — This species is a heterothallic relative of *S. fimicola*.

S. breviseta P. Karst. in Bidr. Känn. Finl. Nat. Folk 23: 52. 1873 = *Podospora appendiculata* (Auersw.) Niessl (Mirza & Cain, 1969) or = *P. pauciseta* (Ces.) Trav. (Lundqvist, 1972).

S. burkillii Mass. in Bull. misc. Inf. R. bot. Gdns Kew 1913: 105. 1913. — This is probably identical with *P. appendiculata* (Moreau, 1953).

S. byssiseda Petch in Ann. R. bot. Gdns Peradeniya 7: 302. 1922 = *Podospora* species (according to the description).

S. caerulea Petch in Ceylon J. Sci. (Bot.) 9: 319. 1915. — Probably a *Cercophora* species (Cain, 1934).

S. californica Plowr. in Grevillea 7: 72. 1878 = *Cercophora californica* (Plowr.) Lundq. (Lundqvist, 1972).

S. capturae Speg. in An. Soc. ci. argent. 9: 21. 1880. — Moreau (1953) considers this species to be a synonym of *S. papyricola*, but according to Lundqvist (1972) it is a good species.

S. carbonaria (Phill. & Plowr.) Sacc., Syll. Fung. 1: 233. 1882 = *Zopfiella carbonaria* (Phill. & Plowr.) v. Arx.

S. caricicola Ell. & Ev. in herb. = *S. fimicola* (Lundqvist, 1972).

S. caudata (Curr.) Sacc., Syll. Fung. 1: 236. 1882 = *Cercophora caudata* (Curr.) Lundq. (Hilber & Hilber, 1979).

S. caulincola Ces. in Atti R. Accad. Sci. Fis. Matem. 8(3): 23. 1879 = *Anthostomella caulincola* (Ces.) Sacc. (Moreau, 1953).

S. cervina Cain in Univ. Toronto Stud. (Biol. Ser.) 38: 36. 1934 = *P. cervina* Cain (Mirza & Cain, 1969).

S. cirrifera Speg. in An. Mus. nac. B. Aires 6: 253. 1899 = *Arnium cirriferum* (Speg.) Krug & Cain (Krug & Cain, 1972).

S. cirsii Crouan, Fl. Finistère: 22. 1867. — The type was not available for study.

S. citrina Petch in Ann. R. bot. Gdns Peradeniya 7: 301. 1922 = *Podospora* species (according to the description).

S. cladoniae Anzi in Atti Soc. ital. Sci. nat. 11: 179. 1868 = *Adelococcus cladoniae* (Anzi) Keissl. (Haworth, 1978).

S. clavariae Ces. & De Not. in Comm. Soc. critt. ital. 1: 226. 1863 = *Helminthosphaeria clavariae* (Tul.) Fuckel (von Arx & Müller, 1954).

S. clavata Pat. apud Pat. & Lagerh. in Bull. Soc. mycol. Fr. 11: 226. 1895. The description is not adequate, and the fungus has been excluded from *Sordaria* by Moreau (1953).

S. clypeata De Not., Sphaerici italicici 1: 22. 1863 = *Entosordaria clypeata* (De Not.) v. Arx & Müller (von Arx & Müller, 1954).

S. comata (Kirschst.) Bayer (Bayer, 1924: 71). — This is a doubtful species. It may represent a *Podospora* species (according to the description).

S. communis (Speg.) Sacc., Syll. Fung. 1: 231. 1882 = *Podospora communis* (Speg.) Sacc. (Mirza & Cain, 1969).

S. communis var. *brachyura* = *Podospora communis* (Speg.) Sacc. (Mirza & Cain, 1969).

S. communis var. *macrura* = *Podospora decipiens* (Winter) Niessl (Mirza & Cain, 1969).

S. communis var. *tetraspora* = *Podospora anserina* (Ces. in Rab.) Niessl (Mirza & Cain, 1969).

S. conoidea Cailleux in Bull. Soc. mycol. Fr. 87: 620. 1971 = *Asordaria conoidea* (Cailleux) v. Arx & Guarro.

S. consanguinea Ces. in Rab. F. eur. extraeur., Cent. XXVII; in Hedwigia 21: 10. — This is a doubtful species (Bayer, 1924).

S. coprophila (Fr.) Ces. & De Not. in Comm. Soc. critt. ital. 1: 226. 1863 = *Cercophora coprophila* (Fr.) Lundq. (Lundqvist, 1972).

S. coronifera Grove in J. Bot. 44: 185. 1916 = *Podospora aloides* (Grove) Mirza & Cain (Mirza & Cain, 1969).

S. culmigena Sacc. & Speg. in Michelia 1: 273. 1877 = *Arnium olerum* (Fr.) Lundq. & Krug (Moreau, 1953).

S. curreyi Auersw. ex Niessl in Verh. naturf. Ver. Brünn 10: 192. 1872 = *Arnium olerum* (Fr.) Lundq. & Krug (Moreau, 1953).

S. curvicolla Wint. in Hedwigia 10: 161. 1871 = *Podospora curvicolla* (Wint.) Niessl (Mirza & Cain, 1969).

S. curvispora Cain in Can. J. Res. (Ser. C) 26: 492. 1948 = *Podospora curvispora* (Cain) Cain (Mirza & Cain, 1969).

S. curvula de Bary, Morphol. Physiol. Pilze: 209. 1866 = *Podospora curvula* (de Bary) Niessl.

S. curvula var. *coronata* = *Podospora curvula* (de Bary) Niessl (Mirza & Cain, 1969) = *Schizothecium conicum* (Fuckel) Lundq. (Lundqvist, 1972).

S. curvula var. *coronifera* = *Podospora aloides* (Fuckel) Mirza & Cain (Mirza & Cain, 1969).

S. curvula var. *tetraspora* = *Schizothecium conicum* (Fuckel) Lundq. (Lundqvist, 1972).

S. decipiens Wint. ex Fuckel in Jb. nassau Ver. Naturk. 27-28: 44. 1873 = *Podospora decipiens* (Wint. ex Fuckel) Niessl (Mirza & Cain, 1969).

S. delitschioides Speg. in An. Mus. nac. B. Aires 6: 254. 1899 = *Cercophora sordarioides* (Speg.) Lundq. (Lundqvist, 1972).

S. destruens (Shear) Hawk, in Trans. Br. mycol. Soc. 34: 174. 1951. — This includes non-coprophilous relatives of *S. fimicola*.

S. discospora Auersw. ex Niessl in Verh. naturf. Ver. Brünn 10: 192. 1872 = *Coniochaeta discospora* (Auersw.) Cain (Cain, 1934).

S. discospora var. *major* Wint. (Winter, 1873: 20) = *Coniochaeta scatigena* (Berk. & Broome) Cain (Cain, 1934).

S. dubia E. C. Hans. in Vid. Meddr dansk naturh. Foren. 1876: 337. 1877 = *Podospora dubia* (E. C. Hans.) Cain (Cain, 1962) or *Schizothecium dubium* (Hans.) Lundq. (Lundqvist, 1972).

S. elephantina P. Henn. in Bot. Jb. (ed. Engler) 22: 77. 1895 = *Cercophora coprophila* (Fr.) Lundq. (Moreau, 1953).

S. ellisiana (Griff.) Sacc. in Syll. Fung. 17: 601. 1905 = *Podospora ellisiana* (Griff.) Mirza & Cain (Mirza & Cain, 1969).

S. elongatospora Barrasa & al. in Persoonia 13: 86. 1986.

S. eminens Cain in Univ. Toronto Stud. (Biol. Ser.) 38: 51. 1934 = *Podospora eminens* Cain (Mirza & Cain, 1969).

S. equina (Fr.) Cooke in Grevillea 17: 78. 1887 = *Podospora fimiseda* (Fuckel) Niessl.

S. equina (Berk. & Rav.) Mirza & Cain apud Udagawa & Horie (Kobayashi, 1971: 524). — From the description given in Kobayashi (1971), the fungus represents a relative of *S. fimicola* with ascospores showing a thin gelatinous sheath.

S. equorum (Fuckel) Wint. (Winter, 1873: 77) = *Hypocopra equorum* (Fuckel) Wint. (Bayer, 1924).

S. erecta (Speg.) Sacc., Syll. Fung. 1: 239. 1882 = *Podospora anserina* (Ces. in Rab.) Niessl (Mirza & Cain, 1969) = *Podospora pauciseta* (Ces.) Trav. (Lundqvist, 1972).

S. erostrata (Griff.) Sacc. in Syll. Fung. 17: 604. 1905 = *Zopfiella erostrata* (Griff.) Udagawa & Furuya (Udagawa & Furuya, 1974).

S. fermenti (Fuckel) Wint. (Winter, 1873: 85) = *S. humana* (Moreau, 1953).

S. fimbriata Bayer (Bayer, 1924: 29) = *Podospora fimbriata* (Bayer) Cain (Mirza & Cain, 1969).

S. fimetii (Pers.) Wint. (Winter, 1873: 78) = *Hypocopra fimetii* (Pers.) Fr. (Bayer, 1924).

S. fimicola (Rob.) Ces. & De Not. in Comm. Soc. critt. ital. 1: 226. 1863.

S. fimiseda Ces. & De Not. in Comm. Soc. critt. ital. 1: 226. 1863 = *Podospora fimicola* Ces. (Mirza & Cain, 1969).

S. friesii Niessl in Verh. naturf. Ver. Brünn 3: 112. 1864 = *Coniochaeta pulveracea* (Ehrh.) Munk (von Arx & Müller, 1954).

S. gagliardi De Not., Spher. ital. 21: 23. 1863. — This is a doubtful species.

S. garhwalensis Srivastava & Bhist. in Indian Phytopath. 36: 383. 1983. — The type was not available for study.

S. gigaspora Fuckel in Jb. nassau. Ver. Naturk. 29-30: 26. 1875. — A type is not available and the description is inadequate for recognition of the fungus.

S. globosa Mass. & Salm. in Ann. Bot. 15: 334. 1901 = *Podospora globosa* (Mass. & Salm.) Cain (Cain, 1962).

S. glutinans Cain in Univ. Toronto Stud. (Biol. Ser.) 38: 40. 1934 = *Podospora glutinans* (Cain) Cain (Mirza & Cain, 1969).

S. goundaensis Cailleux in Bull. Soc. mycol. Fr. 87: 620. 1971 = *Asordaria goundaensis* (Cailleux) v. Arx & Guarro.

S. grisea Ces. in Atti R. Accad. Sci. Fis. Matem. 8(3): 22. 1879. — This is a doubtful species and was excluded from *Sordaria* by Moreau (1953).

S. hansenii Oud. in Hedwigia 21: 123. 1882 = *Coniochaeta hansenii* (Oud.) Cain (Cain, 1934).

S. heterothallis Fields & Maniotis in Am. J. Bot. 50: 80. 1963. — This is a heterothallic relative of *S. macrospora*.

S. hippica (Berk. & Rav.) Ellis & Ev., N. Am. Pyrenom.: 127. 1892 = *S. equina* (Berk. & Rav.) Mirza & Cain apud Udagawa & Horie (Kobayashi, 1971).

S. hippica (Sacc.) Cooke in Grevillea 16: 55. 1887 = *S. fimicola* (Lundqvist, 1972).

S. hirsuta (Dang.) Sacc. & Trav. in Syll. Fung. 20: 807. 1911 = *Podospora hirsuta* Dang. (Mirza & Cain, 1969).

S. hirta E. C. Hans. in Vid. Meddr dansk naturh. Foren. 1876: 336. 1877 = *Arniump hirtum* (E. C. Hans.) Lundq. & Krug (Lundqvist, 1972).

S. hispidula Speg. in An. Mus. nac. B. Aires 6: 255. 1899 = *Podospora curvula* (de Bary) Niessl (Mirza & Cain, 1969) = *Podospora pseudominuta* Speg. (Mirza & Cain, 1969).

S. humana (Fuckel) Wint. in Bot. Ztg 30: 835. 1872 = *Asordaria humana* (Fuckel) v. Arx & Guarro.

S. hyalina Griff. in Mem. Torrey bot. Club 11: 48. 1901 = *Phomatospora hyalina* (Griff.) Cain (Cain, 1934) = *Phomatospora minutissima* (Crouan) Lundq. (Lundqvist, 1972).

S. hyalopilosa (Stratton) Cain in Univ. Toronto Stud. (Biol. Ser.) 38: 45. 1934 = *Podospora hyalopilosa* (Stratton) Cain (Mirza & Cain, 1969).

S. hydropila (Kirschst.) Sacc. in Syll. Fung. 24: 845. 1928 = *Cercophora hydropila* (Kirschst.) Lundq. (Lundqvist, 1972).

S. hypocoprooides Speg. in Boln Acad. Ci. Córdoba 11: 189. 1887 = *Hypocopra hypocoprooides* (Speg.) Krug & Cain (Krug & Cain, 1974b).

S. inaequalis Cain in Can. J. Res. (Ser. C.) 26: 489. 1948 = *Podospora inaequalis* (Cain) Cain (Mirza & Cain, 1969).

S. inaequilateralis Cain in Univ. Toronto Stud. (Biol. Ser.) 38: 29. 1934 = *Arniump inaequilateralis* (Cain) Lundq. & Krug (Lundqvist, 1972).

S. indica Srivastava & Tandon in Curr. Sci. 34: 354. 1966 = *Achaetomium globosum* Rai & Tewari (according to the description).

S. insignis E. C. Hans. in Vid. Meddr dansk naturh. Foren. 1876: 336. 1877 = *Strattonia insignis* (E. C. Hans.) Lundq. (Lundqvist, 1972).

S. iowana Ell. & Holw. in J. Mycol. 4: 65. 1888 = *S. fimicola* (Moreau, 1953).

S. kansensis (Griff.) Sacc. & D. Sacc. in Syll. Fung. 17: 602. 1905 = *Arniump kansense* (Griff.) Krug & Cain (Krug & Cain, 1971).

S. kilimandscharica A. Schm. in Jber. schles. Ges. vaterl. Kult.: 22. 1912 = *Podospora kilimandscharica* (A. Schm.) Cain (Cain, 1962).

S. lanuginosa (Preuss) Sacc., Fungi Veneti Ser. 6: 26. 1877 = *Arniump olerum* (Fr.) Lundq. & Krug (Lundqvist, 1972).

S. lanuginosa (Zopf) A. Schm. Verbreit. coproph. Pilze Schles.: 21. 1912 = *Strattonia zoppii* Lundq. (Lundqvist, 1972).

S. lappa Potebnia in Annls Mycol. 5: 13. 1907.

S. leporina Cain in Univ. Toronto Stud. (Biol. Ser.) 38: 30. 1934 = *Arniump leporinum* (Cain) Lundq. & Krug.

S. leucoplaca (Berk. & Rav.) Ell. & Ev., N. Am. Pyrenom.: 127. 1892 = *Coniochaeta leucoplaca* (Berk. & Rav.) Cain (Cain, 1934).

S. leucotricha Speg. in Michelia 1: 225. 1878 = *Zopfiella leucotricha* (Speg.) Malloch & Cain (Malloch & Cain, 1971).

S. lignicola Fuckel in Jb. nassau. Ver. Naturk. 25-26: 38. 1871 = *Cercophora cadata* (Curr.) Lundq. (Lundqvist, 1972).

S. linguiiformis Cain in Univ. Toronto Stud. (Biol. Ser.) 38: 42. 1934 = *Podospora linguiiformis* (Cain) Cain (Mirza & Cain, 1969) = *Schizothecium dubium* (Hans.) Lundq. (Lundqvist, 1972).

S. longicaudata (Griff.) Sacc. in Syll. Fung. 18: 601. 1905 = *Podospora longicaudata* (Griff.) Cain (Mirza & Cain, 1969).

S. longispora Batista & Pontual in Bol. Agr., Pernambuco 15: 39. 1948 = *Podospora appendiculata* (Auersw.) Niessl (Mirza & Cain, 1969).

S. lugubris Ces. & De Not. Comment. Soc. critt. ital. 1: 52. 1863 = *Anthostomella lugubris* (Rob. & Desm.) Sacc. (Moreau, 1953).

S. lutea Ell. & Ev. in J. Mycol. 3: 118. 1887 = *Cercophora lanuginosa* (Crouan) Lundq. (Lundqvist, 1972).

S. mabokensis Cailleux in Bull. Soc. mycol. Fr. 87: 620. 1971 = *Asordaria mabokensis* (Cailleux) v. Arx & Guarro.

S. macrospora Auersw. in Hedwigia 5: 192. 1866.

S. macrostoma Speg. in An. Mus. nac. B. Aires 6: 252. 1899 = *Podospora communis* (Speg.) Niessl (Mirza & Cain, 1969).

S. macrotheca (Crouan) Cooke in Grevillea 16: 55. 1887 = *Arniump macrotheca* (Crouan) Lundq. (Lundqvist, 1972).

S. macrura (Bayer, 1924: 71) = *Podospora australis* (Speg.) Niessl (Mirza & Cain, 1969).

S. malacotricha Auersw. ex Niessl in Verh. naturf. Ver. Brünn 10: 188. 1872 = *Coniochaeta malacotricha* (Auersw. ex Niessl) Trav.

S. maxima Niessl in Rab., Fungi Eur. exs. 1340. 1866 = *Bombardioidea stercoris* (DC.) Lundq. (Lundqvist, 1972).

S. merdaria (Fr.) Auersw. ex Wint. (Winter, 1873: 13) = *Hypocopra merdaria* Fries (Bayer, 1924).

S. microscopica De Not. in Comm. Soc. critt. ital. 2: 480. 1867 = *Rosellinia microscopica* (De Not.) Sacc. (Moreau, 1953).

S. microspora Ces. in Atti R. Accad. Sci. Fis. Matem. 8(3): 22. 1879 = *Rosellinia microspora* (Ces.) Sacc. (Moreau, 1953).

S. microspora Plowr. in Grevillea 6: 28. 1877 = *Coniochaeta leucoplaca* (Berk. & Rav.) Cain (Cain, 1934).

S. micrura (Speg.) Sacc., Syll. Fung. 1: 234. 1882 = *Podospora curvula* (de Bary) Niessl (Mirza & Cain, 1969).

S. millespora (A. Schm.) Cain in Univ. Toronto Stud. (Biol. Ser.) 38: 52. 1934 = *Podospora millespora* (Schmidt) Cain (Cain, 1962).

S. minima Sacc. & Speg. in Michelia 1: 373. 1878. — This is a doubtful species; Moreau (1953) excluded it from *Sordaria*; it probably is a species of *Coniochaeta*.

S. minor (Ell. & Ev.) Sacc. & Syd. in Syll. Fung. 14: 493. 1988 = *Podospora minor* Ell. & Ev. (Mirza & Cain, 1969).

S. minuta Fuckel in Jb. nassau. Ver. Naturk. 27-28: 44. 1873 = *Podospora vesticola* (Berk. & Br.) Mirza & Cain (Mirza & Cain, 1969).

S. minutella Cooke & Plowr. in Grevillea 7: 86. 1874 = *Trichodelitschia bisporula* (Crouan) Lundq. (Lundqvist, 1964a).

S. moravica (Niessl) Cooke in Grevillea 16: 56. 1887 = *Trichodelitschia bisporula* (Crouan) Lundq. (Lundqvist, 1964a).

S. multicaudata (Griff.) Sacc. in Syll. Fung. 17: 603. 1905 = *Arnium macrotheca* (Crouan) Lundq. (Lundqvist, 1972).

S. musae (Maire) Sacc. in Syll. Fung. 24: 847. 1928 = *Cercophora* spec. (according to the description).

S. myriaspore Crouan, Fl. Finistère: 22. 1867 = *Podospora myriaspore* (Crouan) Niessl (Lundqvist, 1972).

S. natalitia (Speg.) Sacc., Syll. Fung. 1: 231. 1882. — This is a doubtful species and was excluded from *Sordaria* by Moreau (1953).

S. neglecta E. C. Hans. in Vid. Meddr dansk naturh. Foren. 1876: 335. 1877 = *Arnium olerum* (Fr.) Lundq. & Krug (Lundqvist, 1972).

S. oblectans Ces. in Atti R. Accad. Sci. Fis. Matem. 8(3): 23. 1879 = *Rosellinia oblectans* (Ces.) Cooke (Moreau, 1953).

S. obliquata Ces. & De Not. in Comm. Soc. critt. ital. 1: 52. 1863 = *Rosellinia obliquata* (Somm.) Sacc. (Moreau, 1953).

S. occidentalis Batista & Pontual in Bol. Agr., Pernambuco 15: 38. 1948 = *Podospora communis* (Speg.) Niessl (Mirza & Cain, 1969).

S. ontariensis Cain in Univ. Toronto Stud. (Biol. Ser.) 38: 31. 1934 = *Arnium ontariense* (Cain) Krug & Cain (Krug & Cain, 1971).

S. oryzae Hara, Diseases of the riceplant: 132. 1918. — The type was not available and the description is not adequate for recognition of the fungus.

S. oryzeti Sacc. in N. Gior. bot. ital. 23: 201. 1916. — This is a doubtful species and has been excluded from *Sordaria* by Moreau (1953).

S. oryzina Hara, Diseases of the riceplant: 133. 1918. — The type was not available and the description is inadequate for recognition of the fungus.

S. ostiolata Ell. & Ev. in Bull. Torrey bot. Club 24: 458. 1897 = *S. fimicola* (Moreau, 1953).

S. pachydermatica Ces. in Atti R. Accad. Sci. Fis. Matem. 8(3): 22. 1879. Moreau (1953) includes this species in *Rosellinia* or *Coniochaeta*.

S. pachyspora Mout. in Bull. Soc. r. Bot. Belg. 39: 38. 1900. — This name has been excluded from *Sordaria* by Moreau (1953).

S. palmicola Auersw., Un. itin. Crypt., No. 58. 1866 = *Anthostomella palmicola* (Auersw.) Rab. (Pirozynski, 1972).

S. papillata Urries in Boln Soc. esp. Hist. nat. 32: 213. 1932.

S. papillosa Bayer (Bayer, 1924: 51) = *Sordaria humana* (Moreau, 1953).

S. papyricola Wint. (Winter, 1873: 18). — This is a nomen dubium (Lundqvist, 1972).

S. penicillata Ell. & Ev. in J. Mycol. 4: 78. 1888 = *Podospora pauciseta* (Ces.) Trav. (Lundqvist, 1972).

S. perfidiosa De Not. in Comm. Soc. critt. ital. 2: 481. 1867 = *Entosordaria perfidiosa* (De Not.) Hoehnel (von Arx & Müller, 1954).

S. perileuca Sacc. in Atti Accad. sci. ven.-trent.-istr. 10: 66. 1917 = *Cercophora sara-wacensis* (Ces.) Lundq. (Lundqvist, 1972).

S. perplexans Cain in Univ. Toronto Stud. (Biol. Ser.) 38: 50. 1934 = *Podospora perplexans* (Cain) Cain (Cain, 1962).

S. philocoproides Griff. in Mem. Torrey bot. Club 11: 54. 1901 = *Coniochaeta philo-coproides* (Griff.) Cain (Cain, 1934).

S. phyllogena (Sacc.) Niessl in Hedwigia 22: 155. 1883 = *S. humana*.

S. pilifera Bayer (Bayer, 1924: 62). — The type was not available. Probably a species with small ascospores ($10.5 \times 7 \mu\text{m}$) was studied.

S. pilosa Mout. in Bull. Soc. r. Bot. Belg. 25: 144. 1886 = *Podospora pilosa* (Mout.) Cain (Mirza & Cain, 1969) = *Cercophora elephantina* (P. Henn.) Lundq. (Lundqvist, 1972).

S. pilosa Petch in Ann. R. bot. Gdns Peradeniya 7: 302. 1922 = *Podospora* species (according to the description).

S. platyspora Plowr. in Grevillea 6: 28. 1877 = *Coniochaeta scatigena* (Berk. & Br.) Cain (Cain, 1962).

S. pleiospora Wint. in Hedwigia 10: 161. 1871 = *Podospora pleiospora* (Wint.) Niessl (Mirza & Cain, 1969).

S. polyspora Phill. & Plowr. in Grevillea 10: 73. 1881 = *Coniochaeta* sp. (Cain, 1934).

S. potentillae Gucewiz, Plant. non vasc.: 218. 1967. — The description and illustrations are incomplete. Probably a species of *Asordaria* with small ascospores ($14-15 \times 8.5-10 \mu\text{m}$) was studied.

S. prolifica Cailleux in Bull. Soc. mycol. Fr. 87: 620. 1971 = *Asordaria prolifica* (Cailleux) v. Arx & Guarro.

S. pseudominuta Speg. in Boln Acad. Ci. Córdoba 11: 57. 1887 = *Podospora curvula* (de Bary) Niessl (Mirza & Cain, 1969).

S. pumila Bayer (Bayer, 1924: 49). — The type was not available for study. The species is characterized by very small ascospores ($8-9 \times 5-7 \mu\text{m}$).

S. puntiformis Ces. in Atti R. Accad. Sci. Fis. Matem. 8(3): 23. 1879. — This is a doubtful species. Moreau (1953) excluded it from *Sordaria*.

S. pyriformis Bayer (Bayer, 1924: 113) = *Podospora pyriformis* (Bayer) Cain (Mirza & Cain, 1969).

S. rabenhorstii Niessl in Rab., Fungi eur. 1528. 1872 = *Fimetariella rabenhorstii* (Niessl) Lundq. (Lundqvist, 1946b).

S. rhynchophora De Not. in Comm. critt. ital. 2: 480. 1867 = *Ceratostoma notarisii* Sacc. (Moreau, 1953).

S. rotula (Cooke) Niessl in Hedwigia 22: 155. 1883 = *Apiosordaria rotula* (Cooke) v. Arx.

S. sarawacensis Ces. in Atti R. Accad. Sci. Fis. Matem. 8(3): 22. 1879. — This is a doubtful species.

S. sclerogenia Fields & Grear. in Mycologia 58: 524. 1966. — This is a heterothallic relative of *S. fimicola* with a spermatrical state.

S. seminuda Griff. in Mem. Torrey bot. Club 11: 50. 1901 = *Podospora seminuda* (Griff.) Mirza & Cain (Mirza & Cain, 1969).

S. septospora Cain in Univ. Toronto Stud. (Biol. Ser.) 38: 24. 1934. — The type could not be examined.

S. serignanensis (Fabre) Cooke in Grevillea 16: 55. 1887 = *Bombardioidea serignanensis* (Fabre) Lundq.

S. setosa Wint. (Winter, 1873: 97) = *Podospora setosa* (Wint.) Niessl (Mirza & Cain, 1969).

S. sibutii Cailleux in Bull. Soc. mycol. Fr. 87: 620. 1971 = *Asordaria sibutii* (Cailleux) v. Arx & Guarro.

S. similis E. C. Hans. in Vid. Meddr dansk Naturh. Foren 1876: 336. 1877 = *Podospora similis* (E. C. Hans.) Niessl (Mirza & Cain, 1969).

S. socia De Not. in Comm. Soc. critt. ital. 2: 480. 1867 = *Rosellinia socia* (De Not.) Sacc. (Moreau, 1953).

S. sordariooides (Speg.) Cooke in Grevillea 16: 56. 1887 = *Cercophora sordariooides* (Speg.) Lundq. (Lundqvist, 1972).

S. sphaerospora Ell. & Ev., N. Am. Pyrenom.: 128. 1892 = *Asordaria humana* (Moreau, 1953).

S. squamulosa Crouan, Fl. Finistère: 22. 1867 = *Schizothecium squamulosum* (Crouan) Lundq.

S. stercaria (Sow.) Oud., Rév. Champ. Pays Bas: 159. 1897 = *Hypocopra stercaria* (Sow.) Fuckel.

S. striata Ell. & Ev. in J. Mycol. 4: 67. 1888 = *Cercophora striata* (Ell. & Ev.) Lundq. (Lundqvist, 1972).

S. superba De Not. in Comm. Soc. crit. ital. 2: 479. 1867.

S. superior (Griff.) Sacc. & D. Sacc. in Syll. Fung. 17: 603. 1905 = *Podospora appendiculata* (Auersw.) Niessl (Mirza & Cain, 1969).

S. sylvatica Daszewska in Bull. Soc. bot. Genève 4: 294. 1912 = *S. fimicola* (Moreau, 1953).

S. taediosa Speg. in An. Mus. nac. B. Aires 6: 255. 1899. — This is a nomen dubium (Lundqvist, 1972).

S. taenioides (Griff.) Sacc. & D. Sacc. in Syll. Fung. 17: 602. 1905 = *Podospora australis* (Speg.) Niessl (Mirza & Cain, 1969).

S. tetraspora Wint. in Hedwigia 10: 161. 1871 = *Podospora tetraspora* (Wint.) Cain (Mirza & Cain, 1969).

S. texanica (Rehm) Doguet in Botaniste 39: 222. 1955. — The type could not be examined.

S. thermophilila Fields in Mycologia 60: 1117. 1968. — This is a heterothallic relative of *S. fimicola* with thermophilic requirements.

S. tibiodiana Penz. & Sacc. in Malpighia 11: 395. 1897. — The type could not be examined. Moreau (1953) excluded the species from *Sordaria*.

S. tomento-alba Cailleux in Bull. Soc. mycol. Fr. 87: 620. 1971. — This species is a heterothallic relative of *S. fimicola*.

S. tomentosa (Speg.) Sacc., Syll. Fung. 1: 236. 1882 = *Arnium tomentosum* (Speg.) Lundq. & Krug (Lundqvist, 1972).

S. ustorum Mout. in Bull. Soc. r. Bot. Belg. 36: 11. 1897 = *Apiosordaria rotula* (Cooke) v. Arx.

S. uvicola Viala & Marsais in Annls Ins. natn. Rech. agron., Paris (C) 2: 68. 1927 = *S. destruens* (Shear) Hawk. (Lundqvist, 1972) = *S. fimicola* (Cain, 1950).

S. vagans De Not. in Comm. Soc. critt. ital. 2: 479. 1867. — The type was not available for examination. Its description is inadequate for the recognition of the fungus.

S. valoides (Peck) Sacc., Syll. Fung. 1: 235. 1882 = *Podospora appendiculata* (Auersw.) Niessl (Mirza & Cain, 1969).

S. verruculosa (Jensen) Sacc. in Syll. Fung. 24: 845. 1928 = *Apiosordaria verruculosa* (Jensen) v. Arx & Gams.

S. vesticola (Berk. & Br.) Höhn. in Annls Mycol. 16: 45. 1918 = *Podospora vesticola* (Berk. & Br.) Mirza & Cain (Mirza & Cain, 1969).

S. vestita (Zopf) Sacc. & D. Sacc. in Syll. Fung. 18: 604. 1905 = *Podospora communis* (Speg.) Niessl (Mirza & Cain, 1969).

S. violacea Ell. & Ev. in Am. Nat. 31: 340. 1897 = *Podosordaria violacea* (Ell. & Ev.) Krug & Cain (Krug & Cain, 1974a).

S. vratislaviensis A. Schm., Verbreit. coproph. Pilze Schles.: 22. 1912 = *Podospora fimbriata* (Bayer) Cain (Mirza & Cain, 1969).

S. wiesneri Zuk. in Verh. zool.-bot. Ges. Wien 37: 41. 1887 = *Asordaria humana* (Moreau, 1953).

S. winteri P. Karst. in Bidr. Känn. Finl. Nat. Folk 23: 151. 1873 = *Podospora appendiculata* (Auersw.) Niessl (Mirza & Cain, 1969).

S. winteri Oud. in Hedwigia 21: 123. 1884 = *S. fimicola* (Moreau, 1953).

S. zygospora Speg. in Michelia 1: 227. 1878 = *Zygopleurage zygospora* (Speg.) Boed. (Moreau, 1953).

REFERENCES

- ARX, J. A. von, GUARRO, J. & AA, H. A. van der (1987). *Asordaria*, a new genus of the Sordariaceae, and a new species of *Melanocarpus*. In Persoonia 13: 263–272.
 ARX, J. A. von & MÜLLER, E. (1954). Die Gattungen der amerosporen Pyrenomyceten. In Beitr. Krypt. Fl. Schweiz 11(1): 1–434.
 BARRASA, J. M., LUNDQVIST, N. & MORENO, G. (1986). Notes on the genus *Sordaria* in Spain. In Persoonia 13: 83–88.

- BAYER, A. (1924). Monographiká studie středoevropských druhů celedi Sordariaceae. In Práce Moravské Prírod. Spolec, Brno 1: 29–185.
- BOEDIJN, K. B. (1962). The Sordariaceae of Indonesia. In Persoonia 2: 305–320.
- CAILLEUX, R. (1972). Recherches sur la mycoflore Centrafricaine. Les genres *Sordaria*, *Gelsinospora*, *Bombardia*. Ecologie. In Bull. Soc. mycol. Fr. 87: 461–626 ('1971').
- CAIN, R. F. (1934). Studies of coprophilous Sphaeriales in Ontario. In Univ. Toronto Stud. (Biol. Ser.) 38: 1–126.
- (1950). Studies of coprophilous Ascomycetes I. *Gelasinospora*. In Can. J. Res. (C) 28: 566–576.
- (1962). Studies of coprophilous Ascomycetes VIII. New species of *Podospora*. In Can. J. Bot. 40: 447–490.
- HAWKSWORTH, D. L. (1978). Notes on British lichenicolous fungi II. In Notes R. bot. Gdn Edinb. 36: 181–197.
- (1980). *Sordaria bosensis*, an earlier epithet for *Areolospora terrophila*. In Norwegian J. Bot. 27: 97–100.
- HILBER, R. & HILBER, O. (1979). Einige Anmerkungen zu der Gattung *Cercophora* (Lasiosphaeriaceae). In Z. Mykol. 45: 209–233.
- KOBAYASHI, Y. (1971). Mycological reports from New Guinea and the Solomon Islands (1–11). In Bull. natn. Sci. Mus., Tokyo 14: 367–551.
- KRUG, J. C. & CAIN, R. F. (1972). Additions to the genus *Arnium*. In Can. J. Bot. 50: 367–373.
- & — (1974a). A preliminary treatment of the genus *Podosordaria*. In Can. J. Bot. 52: 589–605.
- & — (1974b). New species of *Hypocopra* (Xylariales). In Can. J. Bot. 52: 809–843.
- LUNDQVIST, N. (1964a). The genus *Trichodelitschia* in Sweden. In Svensk bot. Tidskr. 58: 267–272.
- (1964b). *Fimetariella*, a new genus of coprophilous Pyrenomycetes. In Bot. Notiser 117: 248.
- (1972). Nordic Sordariaceae s.lat. In Symb. bot. upsal. 20(1): 1–374.
- MALLOCH, D. & CAIN, R. F. (1971). New cleistothelial Sordariaceae and a new family, Coniochaetaceae. In Can. J. Bot. 49: 869–880.
- MIRZA, J. H. & CAIN, R. F. (1969). Revision of the genus *Podospora*. In Can. J. Bot. 47: 1999–2048.
- MOREAU, C. (1953). Les genres *Sordaria* et *Pleurage*. In Encycl. Mycol. 25: 1–330.
- PIROZYNSKI, K. A. (1972). Microfungi of Tanzania. In Mycol. Pap. 129: 1–64.
- UDAGAWA, S. & FURUYA, K. (1974). Notes on some Japanese Ascomycetes XII. In Trans. mycol. Soc. Japan 15: 206–214.
- WINTER, G. (1873). Die deutsche Sordarien. In Abh. naturf. Ges. Halle 13: 65–107 (1877 as separate issue).