

RESURRECTION OF HANCEA AND LECTOTYPIFICATION OF ADISCA (EUPHORBIACEAE)

S.E.C. SIERRA, K.K.M. KULJU, J.F. VELDKAMP & P.C. VAN WELZEN

Nationaal Herbarium Nederland, Universiteit Leiden branch, P.O. Box 9514,
2300 RA Leiden, The Netherlands; e-mail: sierra@nhn.leidenuniv.nl

SUMMARY

Based on phylogenetic studies with DNA sequence data, the formerly monotypic genus *Cordemoya* was recently expanded with 16 additional species previously assigned to *Mallotus*. However, the earliest legitimate generic name available is *Hancea*, and, therefore, *Cordemoya* is here reduced to *Hancea* (20 new combinations are proposed). Subsequently, the subgenus and section *Diplochlamys* from Asia are renamed to *Hancea*. Additionally, the genus *Adisca* is lectotypified with *Adisca floribundus* (= *Mallotus floribundus*).

Key words: Euphorbiaceae, *Adisca*, *Cordemoya*, *Diplochlamys*, *Hancea*, *Mallotus*, Madagascar, Malesia, Mascarenes.

INTRODUCTION

Based on a well-supported clade obtained with DNA sequence data (Kulju et al., 2007, f. 2), the genus *Cordemoya* Baill. (formerly monotypic with *C. integrifolia* (Willd.) Baill. from the Mascarene Islands as single species) was expanded by Sierra et al. (2006) with sixteen additional species previously assigned to *Mallotus* Lour. Pollen and leaf anatomical studies revealed characters that are important for the delimitation of both genera. *Cordemoya* can be distinguished by the presence of pollen ornamentation with scabrae and by capitate glandular hairs with multicellular stalks and sessile peltate-stellate hairs with a central cell, while *Mallotus* is recognized by the presence of a perforate/micro-reticulate pollen ornamentation with scabrae and by spherical to disc-shaped multicellular glandular hairs. The rank of subgenus was assigned to the two strongly supported and geographically distinct *Cordemoya* clades present in the phylogeny of Kulju et al. (2007): subg. *Cordemoya* from Madagascar and the Mascarenes and subg. *Diplochlamys* (Müll.Arg.) S.E.C. Sierra, Kulju & Welzen from Asia; the latter was divided into two sections: *Diplochlamys* (former *Mallotus* sect. *Hancea* (Seem.) Pax & K. Hoffm.) and *Oliganthae* (Airy Shaw) S.E.C. Sierra, Kulju & Welzen. For a complete synonymy list of the species of subg. *Hancea* see Slik & Welzen (2001). For subg. *Hancea* sect. *Oliganthae* and subg. *Cordemoya* see Sierra et al. (2006).

The epithet '*Hancea*' was usually associated with use on the sectional level within *Mallotus*. However, its generic (and subgeneric) status must also be resurrected, as it is the earliest legitimate generic name available for the basal clade in the phylogeny by Kulju et al. (2007): *Hancea* Seem. was published in 1857 while *Cordemoya* was

published in 1861. There is even an older name, the first generic name is *Boutonia* Bojer (1837), however it was invalidly described and its later validation by Bojer in 1846 created a later homonym.

Furthermore, among the species recognized in the circumscription of *Cordemoya* by Sierra et al. (2006), *Cordemoya subpeltata* (Blume) M. Aparicio has as synonym *Adisca subpeltata* Blume. Until now the genus *Adisca* Blume was not yet lectotypified; one could decide to select *C. subpeltata* as the type of *Adisca*, and it would become the oldest available generic name for *Cordemoya*. However, we opt to appoint *Adisca floribunda* Blume as the lectotype of *Adisca*, as was implicitly done by Slik & Welzen (2001), who included *Adisca* in the synonymy of *Mallotus* sect. *Stylanthus* (Rchb.f. & Zoll.) Pax & K. Hoffm. (from the species included in *Adisca* by Blume, Airy Shaw (1968: 388) selected *A. floribunda* as the type of *Mallotus* sect. *Stylanthus*). From 1 January 2001, the International Code of Botanical Nomenclature (McNeill et al., 2006) states that one should mention “here designated or an equivalent” (Art. 7.11) and this information is lacking in the publication of Slik et al. (2001).

NEW COMBINATIONS IN HANCEA

HANCEA

- Hancea* Seem. (1857) 409. — *Mallotus* Lour. sect. *Hancea* (Seem.) Pax & K. Hoffm. (1914) 199. — Type: *Hancea hookeriana* Seem.
- Cordemoya* Baill. (1861) 255. — *Boutonia* Bojer (1837) 282, nom. nud.; (1846) 151, non DC. (1838). — *Mallotus* Lour. sect. *Cordemoya* (Baill.) Müll.Arg. (1865) 186. — *Echinus* Lour. sect. *Cordemoya* (Baill.) Baill. (1891) 977. — *Cordemoya* Baill. subg. *Cordemoya*: S.E.C. Sierra, Kulju & Welzen in Sierra et al. (2006) 526. — Type: *Boutonia mascareinensis* Bojer [= *Hancea integrifolia* (Willd.) S.E.C. Sierra, Kulju & Welzen].
- Diplochlamys* Müll. Arg. (1864) 539; (1866) 1023. — *Mallotus* Lour. sect. *Diplochlamys* (Müll. Arg.) Baill. ex T. Durand (1888) 370; Müll. Arg. ex Pax (1890) 55, isonym. — *Cordemoya* Baill. subg. *Diplochlamys* (Müll. Arg.) S.E.C. Sierra, Kulju & Welzen in Sierra et al. (2006) 532. — *Cordemoya* Baill. sect. *Diplochlamys* (Müll. Arg.) S.E.C. Sierra, Kulju & Welzen in Sierra et al. (2006) 533. — Type: *Diplochlamys griffithianus* Müll. Arg. [= *Hancea griffithiana* (Müll. Arg.) S.E.C. Sierra, Kulju & Welzen].
- Deuteromallotus* Pax & K. Hoffm. (1914) 212. — Type: *Deuteromallotus acuminatus* (Baill.) Pax & K. Hoffm. [= *Hancea acuminata* (Baill.) S.E.C. Sierra, Kulju & Welzen].
- Mallotus* Lour. sect. *Oliganthae* Airy Shaw (1968) 389. — *Cordemoya* Baill. sect. *Oliganthae* S.E.C. Sierra, Kulju & Welzen in Sierra et al. (2006) 535. — Type: *Mallotus subpeltatus* Blume [= *Hancea subpeltata* (Blume) S.E.C. Sierra, Kulju & Welzen].

Subgenus *Cordemoya* (Baill.) S.E.C. Sierra, Kulju & Welzen, *comb. nov.*

- Cordemoya* Baill. (1861) 255. — *Boutonia* Bojer (1837) 282, nom. nud.; (1846) 151, non DC. (1838). — *Mallotus* Lour. sect. *Cordemoya* (Baill.) Müll.Arg. (1865) 186. — *Echinus* Lour. sect. *Cordemoya* (Baill.) Baill. (1891) 977. — *Cordemoya* Baill. subg. *Cordemoya*: S.E.C. Sierra, Kulju & Welzen in Sierra et al. (2006) 526. — Type: *Boutonia mascareinensis* Bojer [= *Hancea integrifolia* (Willd.) S.E.C. Sierra, Kulju & Welzen].
- Deuteromallotus* Pax & K. Hoffm. (1914) 212. — Type: *Deuteromallotus acuminatus* (Baill.) Pax & K. Hoffm. [= *Hancea acuminata* (Baill.) S.E.C. Sierra, Kulju & Welzen].

1. *Hancea acuminata* (Baill.) S.E.C. Sierra, Kulju & Welzen, *comb. nov.*

Boutonia acuminata Baill. (1858) 401. — Lectotype (McPherson, 1995): *Chapelier s.n.* (P), Madagascar.

2. *Hancea capuronii* (Léandri) S.E.C. Sierra, Kulju & Welzen, *comb. nov.*

Deuteromallotus capuronii Léandri (1957, '1956') 605. — Type: *SF (MAD)* [Capuron] 8713 (holo P; iso MO), Madagascar, Minombo, Amboditavolo.

3. *Hancea integrifolia* (Willd.) S.E.C. Sierra, Kulju & Welzen, *comb. nov.*

Ricinus integrifolius Willd. (1805) 567. — Lectotype (Coode, 1982): *Unknown s.n. (Herb. Willd. 17934, B*, here limited to the sheet with Radlkofer's notes (IDC microfiche no. 1261)), Mauritius.

4. *Hancea spinulosa* (McPherson) S.E.C. Sierra, Kulju & Welzen, *comb. nov.*

Mallotus spinulosus McPherson (1995) 170. — Type: *Schatz & Lowry 1301* (holo MO; iso G, K, P, S, WAG), Madagascar, Tamatave, Ambila-Lemaitso.

Subgenus *Hancea*

Section *Hancea* Seem.

Hancea Seem. (1857) 409. — *Mallotus* Lour. sect. *Hancea* (Seem.) Pax & K. Hoffm. (1914) 199. — Type: *Hancea hookeriana* Seem.

Diplochlamys Müll.Arg. (1864) 539; (1866) 1023. — *Mallotus* Lour. sect. *Diplochlamys* (Müll. Arg.) Baill. ex T. Durand (1888) 370; Müll.Arg. ex Pax (1890) 55, isonym. — *Cordemoya* Baill. subg. *Cordemoya*: S.E.C. Sierra, Kulju & Welzen in Sierra et al. (2006) 532. — *Cordemoya* Baill. sect. *Diplochlamys* (Müll.Arg.) S.E.C. Sierra, Kulju & Welzen in Sierra et al. (2006) 533. — Type: *Diplochlamys griffithianus* Müll.Arg. [= *Hancea griffithiana* (Müll.Arg.) S.E.C. Sierra, Kulju & Welzen].

5. *Hancea cordatifolia* (Slik) S.E.C. Sierra, Kulju & Welzen, *comb. nov.*

Mallotus cordatifolius Slik (1998) 225. — Type: *PNH (Gutierrez et al.) 117545* (holo L), Philippines, Samar Island, Mt Sohoon.

6. *Hancea eucausta* (Airy Shaw) S.E.C. Sierra, Kulju & Welzen, *comb. nov.*

Mallotus eucaustus Airy Shaw (1969) 80. — Type: *S (Sibat ak Luang) 23630* (holo K; iso L), Borneo, Sarawak, Third Division, Bukit Iju, Ulu Arip, Balingian.

7. *Hancea grandistipularis* (Slik) S.E.C. Sierra, Kulju & Welzen, *comb. nov.*

Mallotus grandistipularis Slik (1998) 227. — Type: *Burley, Tukirin et al. 2021* (holo L; iso CANB, K, US), Indonesia, Sumatra, Riau Province, Tigapulu, 5 km W of Talanglakat.

8. *Hancea griffithiana* (Müll. Arg.) S.E.C. Sierra, Kulju & Welzen, *comb. nov.*

Diplochlamys griffithianus Müll.Arg. (1864) 539. — Type: *Griffith KD 4961* (holo K).

9. *Hancea hirsuta* (Elmer) S.E.C. Sierra, Kulju & Welzen, *comb. nov.*

Mallotus hirsutus Elmer (1915) 2648. — Type: *Elmer 13480* (holo PNH?; iso K, L, NY), Philippines, Mindanao, Agusan Province, Cabadbaran (Mt Urdaneta).

10. *Hancea hookeriana* Seem.

Hancea hookeriana Seem. (1857) 409. — Type: *Champion s.n.* (holo K), Hong Kong.

11. *Hancea kingii* (Hook.f.) S.E.C. Sierra, Kulju & Welzen, *comb. nov.*

Mallotus kingii Hook.f. (1887) 439. — Type: *King's collector 7414* (holo K; iso G, SING), Malaya, Peak.

12. *Hancea longistyla* (Merr.) S.E.C. Sierra, Kulju & Welzen, *comb. nov.*

Mallotus longistylus Merr. (1920) 560. — Type: *BS (Ramos & Pascasio) 35173* (holo PNH†; iso K), Philippines.

13. *Hancea papuana* (J.J. Sm.) S.E.C. Sierra, Kulju & Welzen, *comb. nov.*

Mallotus hookerianus (Seem.) Müll.Arg. var. *papuanus* J.J. Sm. (1912) 787, t. 137. — Type: *Gjellerup 316* (holo L; iso A, BO, K, P, U), Irian Jaya, Bivak Hollandia.

14. *Hancea penangensis* (Müll.Arg.) S.E.C. Sierra, Kulju & Welzen, *comb. nov.*

Mallotus penangensis Müll.Arg. (1865) 186. — Type: *Wallich 8576* (holo G; iso A, K, P), Malaysia, Pulau Penang.

15. *Hancea stipularis* (Meijer ex Airy Shaw) S.E.C. Sierra, Kulju & Welzen, *comb. nov.*

Mallotus stipularis Meijer ex Airy Shaw (1968) 398. — Type: *Lörzing 14746* (holo K; iso L), Sumatra, N Sibajak, W of Bandarbaru.

16. *Hancea wenzeliana* (Slik) S.E.C. Sierra, Kulju & Welzen, *comb. nov.*

Mallotus wenzelianus Slik (1998) 229. — Type: *Wenzel 2697* (holo G; iso A, UC), Philippines, Mindanao, Surigao Province, Jubud.

Section *Oliganthae* (Airy Shaw) S.E.C. Sierra, Kulju & Welzen, *comb. nov.*

Mallotus Lour. sect. *Oliganthae* Airy Shaw (1968) 389. — *Cordemoya* Baill. sect. *Oliganthae* S.E.C. Sierra, Kulju & Welzen in Sierra et al. (2006) 535. — Type: *Mallotus subpeltatus* Blume [= *Hancea subpeltata* (Blume) S.E.C. Sierra, Kulju & Welzen].

17. *Hancea subpeltata* (Blume) M. Aparicio, *comb. nov.*

Adisca subpeltata Blume (1826) 610. — Lectotype (Sierra et al., 2006): *Blume 1510* (holo L), Java, Krawong.

LECTOTYPIIFICATION OF ADISCA

Adisca Blume (1826) 609; Slik & Welzen (2001) 31. — Lectotype (selected here): *Adisca floribunda* Blume [= *Mallotus floribundus* (Blume) Müll. Arg.].

ACKNOWLEDGEMENTS

Special thanks are due to D.J. Mabberley for opening our eyes and to the comments of the anonymous reviewer.

REFERENCES

- Airy Shaw, H.K. 1968. Malesian and other Asiatic Euphorbiaceae. *Kew Bull.* 21: 379–400.
- Airy Shaw, H.K. 1969. Malesian and other Asiatic Euphorbiaceae. *Kew Bull.* 23: 80–82.
- Baillon, H.E. 1858. Étude générale du groupe des Euphorbiacées. Masson, Paris.
- Baillon, H.E. 1861. Species Euphorbiacearum. Euphorbiacées africaines. *Adansonia* 1: 255–256.
- Baillon, H.E. 1891. Liste des plantes de Madagascar. *Bull. Mens. Soc. Linn. Paris* 2: 977–978.
- Blume, C.L. 1826. *Adisca*. In: *Bijdragen tot de Flora van Nederlandsch-Indië* 11: 609–611. Lands Drukkerij, Batavia.
- Bojer, W. 1837. *Hortus Mauritianus*. Aime Mararot & Co., Mauritius.
- Bojer, W. 1846. *Boutonia*. In: L. Bouton, *Trav. Soc. Hist. Nat. Maurice 1842–1846*: 151–153.
- Coode, M.J.E. 1982. Euphorbiaceae. In: *Flora des Mascareignes* 160: 50–51. The Sugar Industries Research Institute, Mauritius. L'Office de la Recherche Scientifique et Technique Outre-mer, Paris; The Royal Botanical Gardens, Kew.
- De Candolle, A.P. 1838. Revue sommaire de la famille des Bignoniacées. *Biblioth. Universelle Genève* II, 17: 134, reprinted in *Ann. Sci. Nat., Bot.* II, 11 (1839): 296.
- Durand, T.A. 1888. *Index generum phanerogamorum*: 370. Durand, Brussel.
- Elmer, A.D.E. 1915. Two hundred twenty six new species. *Leaflet Philipp. Bot.* 7: 2648.
- Hooker, J.D. 1887. *Flora of British India* 5. Reeve & Co., London.
- Kulju, K.K.M., S.E.C. Sierra, S.G.A. Draisma, R. Samuel & P.C. van Welzen. 2007. Molecular phylogeny of *Macaranga*, *Mallotus*, and related genera (Euphorbiaceae s.s.): insights from plastid and nuclear DNA sequence data. *Amer. J. Bot.* 94: 1726–1743.
- Léandri, J. 1957 ('1956'). Euphorbiaceae malgaches nouvelles recoltées par M.R. Capuron. *Bull. Soc. Bot. France* 103: 605–606.
- McNeill, J., F.R. Barrie, H.M. Burdet, V. Demoulin, D.L. Hawksworth, K. Marhold, D.H. Nicolson, J. Prado, P.C. Silva, J.E. Skog, J.H. Wiersema & N.J. Turland (eds.). 2006. International Code of Botanical Nomenclature (Vienna Code) adopted by the Seventeenth International Botanical Congress Vienna, Austria, July 2005. *Regnum Veg.* 146.
- McPherson, G.D. 1995. On *Mallotus* and *Deuteromallotus* (Euphorbiaceae) in Madagascar. *Bull. Mus. Natl. Hist. Nat., B, Adansonia* IV, 17: 170.
- Merrill, E.D. 1920. Notes on Philippine Euphorbiaceae, III. *Philipp. J. Sci.* 16: 559–561.
- Müller Argoviensis, J. 1864. Neue Euphorbiaceen des Herbarium Hooker in Kew. *Flora* 47: 539.
- Müller Argoviensis, J. 1865. Euphorbiaceae. *Linnaea* 34: 184–197.
- Müller Argoviensis, J. 1866. Euphorbiaceae tribus Acalypheae. In: A. de Candolle (ed.), *Prodr.* 15, 2: 956–958, 1023. Masson & Filii, Paris.
- Pax, F.A. 1890. Euphorbiaceae. In: A. Engler, *Die natürlichen Pflanzenfamilien* 3, 5: 53–55. Engelmann, Leipzig.
- Pax, F.A. & K. Hoffmann. 1914. Euphorbiaceae–Acalypheae–Mercurialinae. In: A. Engler (ed.), *Das Pflanzenreich* IV.147.vii: 145–212, 394–397. Engelmann, Leipzig.
- Seemann, B.C. 1857. *Flora of the Island of Hong Kong*. The Botany of the Voyage of H.M.S. Herald: 409–410. Reeve & Co., London.

- Sierra, S.E.C., M. Aparicio, K.K.M. Kulju, Ž. Fišer, P.C. van Welzen & R.W.J.M. van der Ham. 2006. Re-shaping *Mallotus* [part 1]: Expanded circumscription and revision of the genus *Cordemoya* (Euphorbiaceae). *Blumea* 51: 519–540.
- Slik, J.W.F. 1998. Three new Malesian species of *Mallotus* section *Hancea* (Euphorbiaceae). *Blumea* 43: 225–232.
- Slik, J.W.F. & P.C. van Welzen. 2001. A taxonomic revision of *Mallotus* sections *Hancea* and *Stylanthus* (Euphorbiaceae). *Blumea* 46: 3–66.
- Smith, J.J. 1912. Euphorbiaceae. *Nova Guinea* 8, 4: 779–796, t. 130–142.
- Willdenow, C.L. 1805. *Species Plantarum*, ed. 4, 1. Nauk, Berlin.