

***Heliconoides linneensis* sp. nov., a new holoplanktonic gastropod
(Mollusca, Thecosomata) from the Late Oligocene
of the Aquitaine Basin (France, Landes)**

Arie W. Janssen

Janssen, A.W. *Heliconoides linneensis* sp. nov., a new holoplanktonic gastropod (Mollusca, Thecosomata) from the Late Oligocene of the Aquitaine Basin (France, Landes).

Zool. Med. Leiden 82 (9): 1.i.2008: 69-72, figs 1-6.— ISSN 0024-0672.

A.W. Janssen, National Museum of Natural History, P.O. Box 9517, 2300 RA Leiden, The Netherlands.

Present address: 12 Triq tal'Hamrija, Xewkija XWK 9033, Gozo, Malta (ariewjanssen@waldonet.net.mt).

Key words: Mollusca, Gastropoda, Thecosomata, pteropod, new species, Oligocene, France.

The gastropod *Heliconoides linneensis* sp. nov. (Mollusca, Thecosomata) is described from Late Oligocene (Chattian) deposits of several localities in the southwestern Aquitaine Basin (France, Landes).

Introduction

Thecosomatous gastropods (so-called 'pteropods') from the Paleogene and Neogene of the Aquitaine Basin, SW France, have been described in a large number of papers, the most important of which are Daudin (1800), Grateloup (1840-1847), Benoist (1873, 1889), Peyrot (1932, 1933), Curry (1982), Lozouet & Maestrati (1982) and Lozouet & Le Renard (2007). Research and extensive field work during the last 25 years, however, have also yielded a large number of species new for the basin, some of which are new to science. One of these latter, occurring in Late Oligocene deposits, is introduced in the present paper, anticipating a more extensive description of the assemblages (Cahuzac & Janssen, in prep.).

Abbreviations

- BRS collection B.G. Roest, Silvolde, The Netherlands.
MNHN Muséum national d'Histoire naturelle, Laboratoire de Biologie des Invertébrés et de Malacologie, Paris, France. Unofficial registration numbers (PL) refer to the Pierre Lozouet et al. collection.
RGM Nationaal Natuurhistorisch Museum Naturalis, Palaeontology Department (formerly Rijksmuseum van Geologie en Mineralogie), Leiden, The Netherlands (RGM registration numbers).

Systematic palaeontology

Phylum Mollusca Linné, 1758
Class Gastropoda Cuvier, 1797
Superorder Heterobranchia Burmeister, 1837
Order Euthyneura Spengel, 1881
Suborder Thecosomata de Blainville, 1824
Infraorder Euthecosomata Meisenheimer, 1905
Superfamily Limacinoidea Gray, 1847
Family Limacinidae Gray, 1847
Genus *Heliconoides* d'Orbigny, 1836

Type species.— *Heliconoides inflata* (d'Orbigny, 1836) (Recent).

Heliconoides linneensis sp. nov. (figs 1-7)

Type material.— Holotype RGM 541 416 (Fig. 1a-b).

Type locality.— St. Etienne-d'Orthe, outcrop in Ruisseau de l'Église, east of Lartigaou, France (Landes), coordinates $x = 316.500$, $y = 149.150$.


Stratum typicum.— Marnes de St. Etienne-d'Orthe (Oligocene, Chattian), zone NP25 (Steurbaut, in Zorn & Janssen, 1993, p. 61).

Derivatio nominis.— The species is named after the Société linnéenne de Bordeaux, which in the course of time published numerous papers on geology and palaeontology of the Aquitaine Basin. The Société, of course, is named after Carl von Linné.


Diagnosis.— *Heliconoides* of near spherical shape, $3\frac{1}{4}$ moderately convex whorls, tangents along the spira convex, body whorl with oblique reinforcement rim slightly behind the apertural margin.

Description.— The shell is sinistral, small, slightly higher than wide (height of holotype 0.92 mm, width 0.86 mm) and nearly spherical in shape. The protoconch is not clearly separated from the teleoconch. The nucleus has a diameter of *c.* 45 μm and has a granulated to honey-comb micro-ornament (Fig. 7b). There are somewhat more than three, rarely three and a half, moderately convex whorls, together forming a low cone with convex tangents. The body whorl is gradually rounded and has a narrow umbilicus. Close to the apertural margin an obliquely vertical, slightly curved swelling in the shell wall forms a reinforcement rim. At this place the shell wall has a micro-ornament of irregularly arranged pustules (Fig. 6b). The shell wall in front of this rim is very fragile, and broken in practically all specimens.

Paratypes.— All specimens from the Late Oligocene of the Aquitaine Basin, further details on these localities will be specified in Cahuzac & Janssen (in prep.). Belus (Marcon): MNHN PL 6594/17 + 1 (Fig. 2a-b); Marcon (Sas): MNHN unnumbered/2 fragments; Orthevielle (Tauzia): MNHN PL 5497/1 fragment; Peyrehorade (Peyrère): MNHN PL 6626/1 (Fig. 3a-c), MNHN PL 6626/5; RGM 541 403/2, leg. A.W. Janssen; Peyrehorade (Tauziède): MNHN PL 5897/5, 1 fragment; St. Etienne-d'Orthe (Église A): MNHN PL 6593/8 + 10 damaged; St. Etienne-d'Orthe (Église B): MNHN PL 6627/18; St. Etienne-d'Orthe (Église C): MNHN unnumbered/2 fragments, leg. Lozouet ?; St Etienne d'Orthe



Figs 1-5. *Heliconoides linneensis* sp. nov., holotype (1) and paratypes (2-5). 1a-b: St. Etienne-d'Orthe, Lartigaou, RGM 541 416; 2a-b: Belus (Marcon): MNHN PL 6594; 3a-c: Peyrehorade (Peyrère): MNHN PL 6626; 4a-c: St. Paul-les-Dax (Estoti): MNHN PL 6597; 5a-b: St. Paul-les-Dax (Lestrilles): MNHN PL 6614. a: apertural, b: lateral, c: apical views.



Figs 6-7. *Heliconoides linneensis* sp. nov., paratypes. 6a-b: St. Etienne-d'Orthe (Lartigaou), RGM 541 417; a: lateral view, b: micro-ornament just behind apertural reinforcement; 7: same locality, RGM 541 418; a: apical view, b: micro-ornament on nucleus. Bar length 100 μ m (figs. 6a, 7a), 10 μ m (figs. 6b, 7b)

(Hondelatte): RGM 541 457/2 fragments, leg. A./W. Janssen; St. Etienne-d'Orthe (Lartigaou): BRS unnumbered/11; RGM 541 417/1 (Fig. 6a-b), RGM 541 418/1 (Fig. 7), RGM 541 419/88, leg. A.W. Janssen, 1984-1990, RGM 541 449/3, leg./don. A.C. Janse; St. Etienne-d'Orthe (Verdun): MNHN PL 6592/6 + 27; St. Paul-les-Dax (Bezoye): MNHN PL 6595/3 juveniles; St. Paul-les-Dax (Estoti): MNHN PL 6597/1 (Fig. 4a-c); St. Paul-les-Dax (Lestrilles): MNHN PL 6614/1 (Fig. 5a-b).

Distribution.— Exclusively known from Late Oligocene deposits (' Chattian Adour') in the Aquitaine Basin.

Discussion

A closely related species is *Heliconoides vanderweideni* Janssen, 2004 (p. 14, pl. 3, figs. 17a-b, 18a-c), introduced from the Aquitanian Lower Globigerina Limestone Formation of the Maltese Archipelago. It differs from the present species in being more globular, with more convex whorls and the apertural rim is interrupted at 2/3rd of its height.

Acknowledgements

The author is grateful to Dr Pierre Lozouet (Muséum national d'Histoire naturelle, Laboratoire de Biologie des Invertébrés et de Malacologie, Paris, France), Messrs Anton C. Janse (Brielle, The Netherlands) and Ben G. Roest (Silvolde, The Netherlands) for making samples available. Mr Frank P. Wesselingh critically read the manuscript.

References

- Benoist, E.-A., 1873. Catalogue synonymique et raisonné des testacés fossiles recueillis dans les faluns miocènes des communes de La Brède et de Saucats.— Actes de la Société linnéenne de Bordeaux 29(3)9: 5-78, 265-459.
- Benoist, E.-A., 1889. Coquilles fossiles des terrains tertiaires moyens du sud-ouest de la France. Description des céphalopodes, ptéropodes et gastropodes opisthobranches (Acteonidae).— Actes de la Société linnéenne de Bordeaux 42: 11-84.
- Cahuzac, B. & A.W. Janssen, in prep. Eocene to Miocene pteropods (Gastropoda, Euthecosomata) from the Aquitaine Basin, SW France.
- Curry, D., 1982. Ptéropodes éocènes de la tuilerie de Gan (Pyrénées-Atlantiques) et de quelques autres localités du SW de la France.— Cahiers de Micropaléontologie 4 (1981): 35-44 (usually cited as 1981).
- Grateloup, J.P.S. de, 1840-1847. Conchyliologie fossile des terrains tertiaires du Bassin de l'Adour (environs de Dax), 1. Univalves. Atlas. Bordeaux (Lafargue): i-xviii, pls 1-45 with explanations (1840); 1-12, pls 46-48 (suppl.) (1847).
- Janssen, A.W., 2004. Fossils from the Lower Globigerina Limestone Formation at Wardija, Gozo (Miocene, Aquitanian), with a description of some new pteropod species (Mollusca, Gastropoda).— The Central Mediterranean Naturalist 4: 1-33, 5 figs., 4 pls.
- Lozouet, P. & J. Le Renard, 2007. *Praecuvierina lura* (holoplanktonic Mollusca: Gastropoda Euthecosomata) a biostratigraphic marker of the Middle Eocene of Aquitaine Basin.— *Cossmanniana* 11: 1-5, 1 pl.
- Lozouet, P. & P. Maestrati, 1982. Nouvelles espèces de mollusques de l'Oligocène (Stampien) pour les bassins de Paris et d'Aquitaine.— *Archiv für Molluskenkunde* 112: 163-189.
- Peyrot, A., 1932. Conchologie néogénique de l'Aquitaine.— *Ouvrages couronnés de l'Académie des Sciences et Belles-Lettres de Bordeaux* 6: 295-541.
- Peyrot, A., 1933. Conchologie néogénique de l'Aquitaine, 6. Gastropodes (suite).— *Actes de la Société linnéenne de Bordeaux* 85: 5-71.
- Zorn, I. & A.W. Janssen, 1993. *Vaginella tricuspida* sp. nov. from the Late Oligocene of Hungary and France.— *Contributions to Tertiary and Quaternary Geology*, 29: 61-71, 5 pl.

Received: 18.x.2007

Accepted: 1.xi.2007

Edited: J. van Tol