

BULLETIN

ZOÖLOGISCH MUSEUM

UNIVERSITEIT VAN AMSTERDAM

Vol. 14 No. 8 1995

REVISED CATALOGUE OF THE TYPE SPECIMENS OF RECENT AMPHIBIANS AND REPTILES IN THE "ZOÖLOGISCH MUSEUM" UNIVERSITY OF AMSTERDAM THE NETHERLANDS

L. van Tuijl

INTRODUCTION

This is a revision of the catalogue issued in 1966 (Daan & Hillenius) of type specimens of amphibians and reptiles in the collections of the "Zoölogisch Museum Amsterdam" (ZMA), also named: Institute for Systematics and Population Biology, of the University of Amsterdam. These include 51 holotypes, 1 neotype, 20 lectotypes, 84 syntypes, 63 paralectotypes, 187 paratypes; only the type specimens in ZMA collection are recorded.

In many cases, type specimens have been sent to other museums: AMS, BMNH, EHT, FMNH, MCZ, NMBA, NMW, RMNH, UNSM, and ZSM.

The following type specimens are missing in the ZMA collection:

Amphibia: *Chaperina basipalmata* (syntype), *Choeophryne proboscidea* (holotype), *Dyscophina volzi* (2 paralectotypes), *Metopostira macra* (holotype), *Nectophryne sumatrana* (2 paralectotypes), *Rana debussy* (holotype), *Rana fallax* (holotype), *Rana pantherina* (holotype);

Reptilia: *Ablepharus boutonii furcata* (syntype), *Ablepharus burnetti* (2 syntypes), *Gehyra interstitialis* (holotype), *Lygosoma florense* (2 syntypes), *Lygosoma minutum rotundirostrum* (syntype), *Lygosoma mivarti obscurum* (6 syntypes) *Lygosoma verreauxii biunguimaculata* (3 syntypes).

Curators of the Amphibia / Reptilia collection of the ZMA, (Zoölogisch Museum Amsterdam) were: Prof. Max (C. W.) Weber from 1883 to 1922, Dr. P. N. van Kampen from 1900 to 1905, Dr. Nelly (P. J.) de Rooy from 1907 to 1922, Prof. L. F. de Beaufort from 1922 to 1949, and Dr. D. Hillenius from 1954 to 1987. At the moment the collections are curated by the author with assistance of Dr. H. Nijssen.

Whenever possible, the remaining type specimens in other collections are mentioned. The number of specimens is given in parentheses after the registration number.

- AMNH American Museum of Natural History, New York.
- AM Australian Museum, Sydney.
- ANSP Academy of Natural Sciences, Philadelphia.
- BMNH Natural History Museum, London.
- CAS California Academy of Sciences, Los Angeles.
- CM Carnegie Museum, Pittsburgh.
- EHT Edward H. Taylor private collection.
- FMNH Field Museum of Natural History, Chicago.
- IRSNB Institut Royal des Sciences Naturelles de Belgique, Brussels.
- J University of Ljubljana, Slovenia, Biotechnical Faculty, Dept. of Biology.
- KU Kansas University, Museum of Natural History, Lawrence.
- LACM Los Angeles County Museum of Natural History.
- LDG Lescure Jean, private collection, Paris.
- LSUMZ Louisiana State University, Museum of Zoology, New Orleans.
- MCZ Museum of Comparative Zoology, Cambridge, Massachusetts.
- MNCN Museo Nacional de Ciencias Naturales, Madrid.
- MNCR Museo Nacional Costa Rica, San José.
- MNHN Muséum National d'Histoire Naturelle, Paris.
- MVZ Museum of Vertebrate Zoology, University of California at Berkeley.
- MZB Museum Zoologicum, Bogor.
- NMBA Naturhistorisches Museum, Basel.
- NMS National Museum, Singapore.
- NMW Naturhistorisches Museum, Wien.
- PNM Philippine National Museum, Manila.
- RMNH Nationaal Natuurhistorisch Museum, Leiden.
- SMF Senckenberg Museum, Frankfurt am Main.
- SMK Sarawak Museum Kuching, Sarawak.
- UINMH University of Illinois, Museum of Natural History, Urbana.
- UMMZ University of Michigan, Museum of Zoology, Ann Arbor.
- USC University of South Carolina, Columbia.
- USNM United States National Museum of Natural History, Washington D.C.
- WCAB Werner C. A. Bokermann, Sao Paulo.
- ZMA Zoologisch Museum, Amsterdam.
- ZMB Zoologisches Museum, Berlin.
- ZSM Zoologische Staatssammlung, München.

ANURA

- Alytes dickhilleni* Arntzen & Garcia Paris, 1995: 26
Holotype: MNCN 16662, Canillas de Aceituno, Sierra Tejada, Provincia de Malaga, Spain, coll. J. W. Arntzen & M. Garcia Paris, 16.V.1992.
Paratypes: ZMA 8156 (4), 18.V.1992; ZMA 9227 (1), 18.V.1992; ZMA 9228 (2), 18.V.1992, from type locality; ZMA 9229 (1) Baranco de Los Cazadores, Sierra Almijara, Nerja, Prov. Malaga, Spain, coll. A. Antunez, VI. 1991. Other paratypes in MNCN.
- Alytes obstetricans almogavarii* Arntzen & Garcia Paris, 1995: 29
Holotype: MNCN 16663, Rasos de Peguera, Berga, Provincia de Barcelona, Spain, coll. M. Garcia Paris & J. W. Arntzen, 6.IX.1992.
Paratypes: ZMA 9232 (6), from type locality. Other paratypes in MNCN.
- Asterophrys minima* Parker, 1934: 67
Holotype: ZMA 5818, female, Went Mountains, West New Guinea, altitude 1000-1360 m, coll. H. A. Lorentz, 12.X.1909.
Paratype: ZMA 5817, Hellwig Mountains, West New Guinea, altitude 2600 m, coll. H. A. Lorentz, 13.XI.1909.
- Austrochaperina robusta* Fry, 1912: 89
Holotype: AMS R5295, Russell River, N. E. Queensland, Australia, coll. D. H. Fry.
Paratype: ZMA 5877 (1), from type locality. Other paratypes in AMS.
- Boophis hillenii* Blommers-Schlösser, 1979b: 299
Holotype: ZMA 7123a, Perinet, Malagasy, altitude 900 m, coll. R. Blommers-Schlösser, 24.XII.1972.
Paratype: ZMA 7123b, from type locality.
- Boophis mandraka* Blommers-Schlösser, 1979b: 267
Holotype: ZMA 7119a, Mandraka, Malagasy, coll. R. Blommers-Schlösser, 4.III.1973.
Paratypes: ZMA 7119b (6), ZMA 7118 (6), from type locality.
- Boophis reticulatus* Blommers-Schlösser, 1979b: 294
Holotype: ZMA 7101a, Perinet, Malagasy, altitude 1100 m, coll. R. Blommers-Schlösser, 13.XI.1972.

- Paratypes: ZMA 7101b (8), from type locality.
Boophis viridis Blommers-Schlösser, 1979b: 272
 Holotype: ZMA 7100a, Perinet, Malagasy, altitude 900 m, coll. R. Blommers-Schlösser, 14.XI.1972.
- Paratypes: ZMA 7100b (2), from type locality.
Bufo mcgregori Taylor, 1922b: 182
 Holotype: CAS 61839, Pasananka, Zamboanga, Mindanao Island, coll. E. H. Taylor, 30.IX.1920. destroyed in World War II, Brown & Alcalá (1980: 29).
- Paratypes: ZMA 5758-59 (2), from type locality. Other paratypes in EHT, MCZ, and NMBA.
- Chaperina basipalmata* Van Kampen, 1906: 169
 Syntypes: ZMA 5875-76 (2), Timena River, West New Guinea, coll. Dutch New Guinea Expedition, 6-7.IV.1903; ZMA (1), from Tawarin, West New Guinea, could not be found in the collection. Other syntype in RMNH.
- Chaperina ceratophthalmus* Van Kampen, 1909: 43
 Syntypes: ZMA 5777 (1) male, Geitenkamp, Lorentz river, West New Guinea, coll. H. A. Lorentz, 2.VII.1907; ZMA 5778 (1) female; ZMA 5779 (1), Resi peak, West New Guinea, coll. H. A. Lorentz, 20.VIII & 9.IX.1907.
- Chaperina punctata* Van Kampen, 1913b: 463
 Syntypes: ZMA 5747-50 (4), ZMA 5751-53 (3), Went Mountains, West New Guinea, altitude 800-1050 m, coll. H. A. Lorentz, 6.IX.1909; 11-12.X.1909; ZMA 5754 (1), Hellwig Mountains, West New Guinea, altitude 2500 m, coll. H. A. Lorentz, X.1909. Other syntypes in AMS, BMNH, FMNH, and MCZ.
- Choerophryne proboscidea* Van Kampen, 1914: 376
 Holotype: ZMA unregistered, Njao, West New Guinea, coll. P. N. van Kampen, 1910, could not be found in the collection.
- Cophixalus crucifer* Van Kampen, 1913b: 462
 Syntypes: ZMA 5819-20 (2), Went Mountains, West New Guinea, altitude 800 and 1050 m, coll. H. A. Lorentz, 11-12.X.1909.
- Cornufer beauforti* Van Kampen, 1913a: 91
 Syntypes: ZMA 5156 (1), ZMA 5755 (1), Majalibit bay, Waigeu, West New Guinea, coll. L. F. de Beaufort, 1910; ZMA 5756-57 (2), Bajon, Waigeu, West New Guinea, coll. L.F. de Beaufort, 9.II.1910. Other syntypes in, BMNH, FMNH, MCZ, NMBA, and ZSM.
- Dyscophina volzi* Van Kampen, 1905: 708
 Lectotype, designated by Daan & Hillenius (1966: 120)
 ZMA 5871, male, Tandjong Laut, Sumatra, Indonesia, coll. W. Volz; ZMA unregistered, 2 specimens, could not be found in the collection.
- Eleutherodactylus zeuctotylus* Lynch & Hoogmoed, 1977: 432
 Holotype: RMNH 17701, male, Vier Gebroeders Mountain, Sipaliwini, Nickerie District, Surinam, coll. M. S. Hoogmoed, 7.II.1970.
- Paratype: ZMA 6293 (1), Linker Coppename Rivier, Surinam, coll. H. Nijssen, 15.V.1967. Other paratypes in, AMNH, BMNH, FMNH, KU, LG, LDG, MCZ, RMNH, UNSM, and WCAB.
- Hyla papua* Van Kampen, 1909: 33
 Holotype: ZMA 5679, Bivak Island, Lorentz River, West New Guinea, coll. H. A. Lorentz, 30.VI.1907. The holotype was referred to by Tyler (1964: 266; 1968: 94) as "AZM 5679", acronym incorrect.
- Hyla proboscidea* Brongersma, 1933c: 267
 Holotype: ZMA 5710, Upper Gran Rio, Surinam, coll. J. H. A. Tresling, 27.IX.1908.
- Hyla rhacophorus* Van Kampen, 1909: 32
 Holotype: ZMA 5323, Etna Bay, West New Guinea, coll. J. W. R. Koch, 1904-05. The holotype was referred to by Tyler (1965: 99, acronym only; 1968: 87) as "AZM 5823", both acronym and registration number are incorrect.
- Hyla sanguinolenta* Van Kampen, 1909: 33
 Lectotype, designated by Daan & Hillenius (1966: 121)
 ZMA 5676, Sabang, Lorentz River, West New Guinea, coll. H. A. Lorentz, 30.VI.1907.
- Paralectotypes: ZMA 5674 (1), Bivak Island, Lorentz River, West New Guinea, coll. H. A. Lorentz, 3.VIII.1907; ZMA 5675 (1), Lorentz River, West New Guinea, coll. H. A. Lorentz, 20.V.1907. The lectotype and paralectotypes were referred to by Tyler (1968: 164) as "AZM 5676", "AZM 5674", "AZM 5675", acronym incorrect. The "cotype" recorded by Taylor (1944: 177) and by Marx (1976: 53) is not a syntype, being collected in 1909, rather than in 1907.
- Hylophorbus variegatus* Van Kampen, 1923: 138
 Holotype: ZMA 5706, Digul River, South West New Guinea, coll. Dutch New Guinea Expedition.
- Ixalus jacobsoni* Van Kampen, 1912: 78
 Holotype: ZMA 5709, Gunung Ungaran, near Semarang, Java, Indonesia, coll. E. R. Jacobson, 1910.
- Kaloula rigida* Taylor, 1922b: 176
 Holotype: CAS 61475, Balbalan, Kalinga, Mountain Province, Luzon, Philippines, coll. E. H. Taylor, 28-30.IV.1920.
- Paratype: ZMA 5718 (1), from type locality. Other paratypes in AMNH, BMNH, EHT, FMNH, MCZ, NMBA, and UMMZ.
- Lechriodus platyceps* Parker, 1940: 28
 Holotype: BMNH 76.7.18.6, female, Arfak Mountains, West New Guinea, coll. A.A. de Bruijn.
- Paratype: ZMA 5799, juv., Upper Tor River, North West New Guinea, coll. K. Gjellerup, 2.XI.1911. Other paratypes in BMNH and RMNH.
- Mantella pulchra* Parker, 1925: 393
 Holotype: BMNH, Antsihanaka, Malagasy, coll. W. F. H. Rosenberg.
- Paratypes: ZMA 5809-10 (2), from type locality. Other

- paratype in MCZ.
- Mantidactylus flavobrunneus* Blommers-Schlösser, 1979a: 54
 Holotype: ZMA 7172a, Moramanga, Anosibe, Malagasy, coll. R. Blommers-Schlösser, 25.VIII.1971.
 Paratypes: ZMA 7172b (1), from type locality; ZMA 7173 (1), Perinet, Malagasy, coll. R. Blommers-Schlösser, VI.1971.
- Mantidactylus punctatus* Blommers-Schlösser, 1979a: 51
 Holotype: ZMA 7171a, Tampoketsa d'Ankazobe, Malagasy, altitude 1600 m, coll. R. Blommers-Schlösser, 30.III.1972.
 Paratypes: ZMA 7171b (1), from type locality; ZMA 7170 (14), Tampoketsa d'Ankazobe, Malagasy, altitude 1600 m, coll. R. Blommers-Schlösser, 12.IX.1972.
- Metopostira macra* Van Kampen, 1906: 167
 Holotype: ZMA unregistered, collected "am Moso und Tami", North West New Guinea, coll. Dutch New Guinea Expedition, 1903, could not be found in the collection.
- Micrixalus diminutiva* Taylor, 1922c: 267
 Holotype: CAS 61842, Pasananka, Zamboanga, Mindanao Island, Philippines, coll. E. H. Taylor, 10.XI.1920.
 Paratypes: ZMA 5780-81 (2), Philippines, coll. E. H. Taylor, 10-XI-1920. Other paratypes in EHT, MCZ, NMBA, and UMMZ.
- Microhyla niasensis* Van Kampen, 1915a: 279
 Holotype: ZMA 5253, Gunung Sitoli, Nias, near Sumatra, Indonesia, coll. J. P. Kleiweg de Zwaan, 1910.
- Microhyla superciliaris* Parker, 1928: 486
 Holotype: BMNH 1904.7.19.3, Batu Caves, Kuala Lumpur, Malay Peninsula.
 paratype: ZMA 5255 (1), Deli, Sumatra, Indonesia, coll. L. Ph. de Bussy.
- Nectophryne sumatrana* Van Kampen, 1909b: 19
 Lectotype, designated by Daan & Hillenius (1966: 122), ZMA 5177, Bandar Baru, Battak Mountains, Deli, Sumatra, Indonesia, coll. L. Ph. de Bussy.
 Paralectotypes: ZMA unregistered (2), could not be found in the collection.
- Oreophryne flava* Parker, 1934: 168
 Holotype: ZMA 5823, Kloofbivak, Lorentz River, West New Guinea, coll. G. M. Versteeg, 14.X.1912.
- Oxydactyla brevicrus* Van Kampen, 1913b: 465
 Lectotype, designated by Daan & Hillenius (1966: 122), ZMA 5714, Hellwig Mountains, West New Guinea, altitude 2500 m, coll. A. H. Lorentz, X.1909.
 Paralectotypes: ZMA 5715-16 (2), Wichmann Mountains, West New Guinea, altitude 3000 m, coll. H. A. Lorentz, XI.1909.
- Phanerotis novaeguineae* Van Kampen, 1909: 36
 Syntype: ZMA 5874 (1), Merauke, Southwest New Guinea, coll. J. W. R. Koch, 1904-05. Other syntypes in AMS and FMNH.
- Philautus polillensis* Taylor, 1922b: 171
 Holotype: CAS 62250, South Polillo, Philippines, coll. E. H. Taylor, 12-VII-1920.
 Paratype: ZMA 5719 (1), from type locality. Other paratypes in EHT and MCZ.
- Philautus williamsi* Taylor, 1922b: 167
 Holotype: CAS 62253, Polillo Island, Philippines, coll. E. H. Taylor, 12.VIII.1920.
 Paratype: ZMA 5721 (1), between Polillo and Bislian, Luzon, Philippines, coll. E. H. Taylor, 12.VIII.1920. Other paratypes in EHT and MCZ.
- Polypedates linki* Taylor, 1922c: 276
 Holotype: CAS 60684, Jolo, Jolo Island, Sulu Archipelago, Philippines, coll. E. H. Taylor, 10.XI.1920.
 Paratypes: ZMA 5783-84 (2), from type locality. Other paratypes in EHT and MCZ.
- Rana acanthi* Taylor, 1923: 523
 Holotype: CAS 62577, Busuanga Island, Calamian Islands, Philippines, coll. E. H. Taylor.
 Paratypes: ZMA 5785-86 (2), from type locality. Other paratypes in EHT and MCZ.
- Rana debussyi* Van Kampen, 1909b: 23
 Holotype: ZMA unregistered, Bandar Baru, Batak Mountains, Deli, Sumatra, Indonesia, coll. L. Ph. de Bussy, could not be found in the collection.
- Rana fallax* Van Kampen, 1913b: 459
 Holotype: ZMA unregistered, female, Majalibit Bay, Waigeu, West New Guinea, coll. L. F. de Beaufort, 1910, could not be found in the collection.
- Rana grisea* Van Kampen, 1913b: 460
 Holotype: ZMA 5704, female, Went Mountains, West New Guinea, altitude 1360 m, coll. H. A. Lorentz, 13.X.1909. Menzies (1987: 403) recorded ZMA 5796 as a paratype of *Rana grisea*. This species, however, was described from the holotype only.
- Rana guerreroi* Taylor, 1920: 255
 Holotype: probably in Carnegie Museum, EHT 881, Baguio, Mountain Province, Luzon, Philippines, coll. E. H. Taylor, 1.VI.1915.
 Paratype: ZMA 5868 (1), from type locality. Other paratypes in CM and MCZ.
- Rana igorata* Taylor, 1922c: 260
 Holotype: CAS 61484, Balbalan, Kalinga, Subprovince, Northern Luzon, Philippines, coll. E. H. Taylor, 28.IV.1920.
 Paratype: ZMA 5722 (1), from type locality. Other paratypes in EHT and MCZ.
- Rana microtypanum* Van Kampen, 1907: 386
 Syntypes: ZMA 5797 (5), Loka, near Bonthain, Cele-

- bes, Indonesia, coll. M. Weber, 1888. Other syntype in FMNH.
- Rana novaeguineae* Van Kampen, 1909: 37
Lectotype, designated by Daan & Hillenius (1966: 124), ZMA 5761, Sabang, Lorentz River, West New Guinea, coll. H. A. Lorentz, 30.VI.1907.
Paralectotypes: ZMA 5760 (1), ZMA 5762 (1), ZMA 5763-73 (11), from type locality; ZMA 5774 (1), Bivak Island, Lorentz River, West New Guinea, coll. H. A. Lorentz, 24.VII.1907; ZMA 5775-76 (2), Lorentz River, West New Guinea, coll. H. A. Lorentz, 15.IV.1909, Other paralectotypes in EHT and ZSM.
- Rana pantherina* Van Kampen, 1909b: 22
Holotype: ZMA unregistered, Badar Baru, Battak Mountains, Deli, Sumatra, Indonesia, coll. L. Ph. de Bussy, could not be found in the collection.
- Rana tigrina angustopalmata* Van Kampen, 1907: 389
Syntypes: ZMA 5729-31 (3), Waingapu, Sumba, Indonesia, coll. Siboga Expedition, IV.1899; ZMA 5745-46 (2), Saleyer, Indonesia coll. M. Weber, 1888; ZMA 5744 (1), Makassar, Celebes, Indonesia, coll. Siboga Exp. 1899-1900; ZMA 5732 (1), Tempe, Celebes, Indonesia, coll. M. Weber, 1888; ZMA 5733-37 (5), Lalalupa River near Tempe, Celebes, Indonesia, coll. M. Weber, 1888; ZMA 5738 (1), Pampanua, Celebes, Indonesia, coll. M. Weber, 1888; ZMA 5739-40 (2), Katjang, Celebes, Indonesia, coll. M. Weber, 1888; ZMA 5741 (1), Maros, Celebes, Indonesia, coll. M. Weber, 1888; ZMA 5742-43 (2), Pare-pare, Celebes, Indonesia, coll. M. Weber, 1888. Other syntypes in EHT.
- Rana waigeensis* Van Kampen, 1913a: 90
Holotype: ZMA 5782, along Rabiai River, Waigeu, West New Guinea, coll. L. F. de Beaufort, 31.XII.1909. The holotype is referred to by Menzies (1987: 389) as ZMA 782; this number is incorrect.
- Rana yakani* Taylor, 1922c: 262
Holotype: CAS 60135, Abung-abung, Basilan, Philippines, coll. E. H. Taylor, 22.X.1920.
Paratypes: ZMA 5723-24 (2), from type locality. Other paratypes in FMNH, MCZ, NMBA, UIMNH, UMMZ, and UNSM.
- Rhacophorus anodon* Van Kampen, 1907: 400
Holotype: ZMA 5707, female, cleft near Ajer Mantjur, Kaju tanam, Sumatra, Indonesia, coll. M. Weber, 1888.
- Rhacophorus bifasciatus* Van Kampen, 1923: 257
Syntypes: ZMA 5872-73, Sinlak, Deras, Kurintji, Sumatra, Indonesia, coll. E. R. Jacobson, IX.1915. Other syntypes in RMNH.
- Sphenophryne albopunctata* Van Kampen, 1909: 42
Syntypes: ZMA 5821-22, Sabang, Lorentz River, West New Guinea, coll. H.A. Lorentz, 13-24.VI.1907.
Xenorhina bidens Van Kampen, 1909: 39
Holotype: ZMA 5705, Digul River, Southwest New Guinea, coll. J. W. R. Koch, 1904-05.
- Xenorhina gigantea* Van Kampen, 1915b: 40
Lectotype, designated by Daan & Hillenius (1966: 126), ZMA 5702, Bijkenkorfpad, Lorentz River, West New Guinea, coll. G. M. Versteeg, 13.I.1913.
Paralectotype: ZMA 5703 (1), from type locality.
Xenorhina macrops Van Kampen, 1913: 460
Lectotype, designated by Daan & Hillenius (1966: 126), ZMA 5725, peak in Hellwig Mountains, West New Guinea, altitude 2200 m, coll. H. A. Lorentz, 4.IX.1907.
Paralectotypes: ZMA 5726 (1), from type locality; ZMA 5727-28 (2), Hellwig Mountains, West New Guinea, altitude 2500-2600 m, coll. H. A. Lorentz, X-XI.1909.
- Xenorhina ocellata* Van Kampen, 1913b: 461
Syntypes: ZMA 5815-16 (2), FMNH (1), Hellwig Mountains, West New Guinea, altitude 2500-2600 m, coll. H. A. Lorentz, XI.1909.

CAUDATA

- Proteus anguinus parkelj* Sket & Arntzen, 1994: 33-53
Holotype: J8, Na trati, Jelsevnik near Crnomelj, Slovenia.
Paratype: ZMA 9239 (1), from type locality. Other paratypes in J.
- Molge pyrrhogaster* Boie, 1826: 215
Lectotype, designated by Hoogmoed, (1978: 99), ZMA 7286, Japan.
- Oedipina pseudouniformis* Brame, 1968: 25
Holotype: LACM 1729, Cienaga Colorado, Provincia Cartago, Costa Rica, coll. A. G. Kluge, R.J. Lavenberg, A.H. Brame, 11.VI.1959.
Paratypes: ZMA 9031 (2), Provincia de Limón, Zent, Costa Rica, coll. E. R. Dunn, 1920. Other paratypes in AMNH, FMNH, KU, LACM, LSUMZ, MCZ, MNCR, MVZ, UMMZ, USC, and USNM.

GYMNOPHIONA

- Ichthyophis billitonensis* Taylor, 1965: 294
Holotype: ZMA 5209, Billiton Island, Indonesia, coll. Clemtuar, VI.1914.

REPTILIA, LACERTILIA

Ablepharus boutonii furcata Weber, 1890: 174

Syntypes: ZMA 10815 (1), ZMA 10831 (1), Endeh, Flores, Indonesia, coll. M. Weber, 1888; ZMA 10830 (2), Sikka, Flores, Indonesia, coll. M. Weber, 1888; ZMA unregistered, Sikka, Flores, Indonesia, could not be found in the collection.

Ablepharus boutonii intemedius De Jong, 1926a: 93

Lectotype, designated by Mertens (1931: 149), ZMA 10972, Rana, Buru, Indonesia coll. L. J. Toxopeus, 1921.

Paralectotype: ZMA 10973 (1), from type locality.

Ablepharus boutonii keiensis Roux, 1910: 240

Lectotype, designated by Mertens (1931: 147), NMBA 6203, Elat, Great Kei, Indonesia, coll. H. Merton & J. Roux, 4.VI.1908.

Paralectotype: ZMA 10836 (1), from type locality. Other paralectotypes in MCZ, NMBA, and SMF.

Ablepharus bumetti Oudemans, 1894: 144

Syntypes: ZMA 11345-46 (2), Burnett river, Queensland, Australia, coll. R. Semon, 1893; ZMA unregistered (2), from type locality, could not be found in the collection.

Agama stellio cyprica Daan, 1967: 129

Holotype: BMNH 1930.10.5.6. Limasol, Cyprus, coll. R. L. Cheverton, 1930.

Paratypes: ZMA 13073 (1), Nikosia, Cyprus, coll. H. Schnurrenberger, 1962; ZMA 13746 (1), ZMA 13749 (1), Nikosia, Cyprus, coll. Monniot, 1962; ZMA 13747 (1), Temple de Vouni, Cyprus, coll. Monniot, 1962; ZMA 13748 (1), Salamis, Cyprus, coll. Biguet, 1962, Other paratypes in BMNH and RMNH.

Ameiva aguilina Garman, 1888: 3

Syntype: ZMA 18159 (1), Kingston, St. Vincent, coll. S. Garman, 1879.

Other syntypes in ANSP, MCZ, and NMW.

Anolis bonairensis blanquillanus Wagenaar Hummelinck, 1940: 79

Holotype: RMNH 11761, El Jaque, Isla Blanquilla, Venezuela, coll. P. Wagenaar Hummelinck, 22.VII.1936.

Paratypes: ZMA 10977-79 (3), Plantio del Jaque, Blanquilla, Venezuela, coll. P. Wagenaar Hummelinck, 22.VII.1936. Other paratypes in RMNH.

Anolis conspersus Garman, 1887: 273

Syntype: ZMA 18707 (1), Grand Cayman, Cayman Islands, coll. C. J. Maynard, 1885. Other syntypes in ANSP, MCZ, NMW, UMMZ, and USNM.

Bachia intermedia Noble, 1921b: 142

Holotype: MCZ 14630, Perico, Peru, coll. G. K. Noble, IX.1916.

Paratypes: ZMA 11438-441 (4), Bellavista, Peru, coll. G.

K. Noble, 23.IX.1916. Other paratypes in FMNH, MCZ, NMW, UIMNH, UMMZ, and UNSM.

Brachymeles pathfinderi Taylor, 1925: 104

Holotype: MCZ 26581, Glan, Cotobato Province, Mindanao, Philippines, coll. E. H. Taylor, IV.1923.

Paratype: ZMA 11243 (1), Cotobato Province, Mindanao, Philippines, coll. E. H. Taylor, IV.1923. Other paratypes in BMNH, EHT, and MCZ.

Brachymeles vermis Taylor, 1918: 255

Holotype: PNM 1980, Bubuan, Sulu Archipelago, Philippines, destroyed in World War II, Brown & Alcalá (1980: 29).

Paratypes: ZMA 11008-009 (2), Sulu, Philippines, coll. E. H. Taylor, 1917. Other paratypes in MCZ, PNM or EHT.

Chamaeleon incornutus Loveridge, 1932: 380

Holotype: MCZ 31350, male, Madehani, Ukinga Mountains, Tanganyika, coll. A. Loveridge, 14.II.1930.

Paratypes: ZMA 10249 (1) male, ZMA 10984 (1) female, Nyamwanga, Poroto Mountains, Tanganyika, coll. A. Loveridge, 17.III.1930. Other paratypes in FMNH, MCZ, NMW, and UINMH.

Chamaeleon laterispinnis Loveridge, 1932: 381

Holotype: MCZ 31386, male, Kigogo, near Mufindi, Uzungwe Mountains, Tanganyika, coll. A. Loveridge, 23.I.1930.

Paratype: ZMA 10250 (1) male, Kigogo, near Mufindi, Uzungwe Mountains, Tanganyika, coll. A. Loveridge, 13.I.1930. Other paratype in MCZ.

Dicrodon barbouri Noble, 1921: 108

Holotype: MCZ 17972, Sullana, Piura, Chira River, Peru, coll. G. K. Noble, 28.VII.1916.

Paratypes: ZMA 18166 (2), from type locality. Other paratypes in MCZ, NMW, and UMMZ

Draco buruensis De Jong, 1926a: 89

Syntypes: ZMA 10931 (1) male, Leksula, Buru, Indonesia, coll. L. J. Toxopeus, 1921; ZMA 10932-933 (2) males, Ehu, Buru, Indonesia, coll. L. J. Toxopeus, 1921.

Draco fimbriatus hennigi Musters, 1983: 27

Holotype: RMNH 19973, Soekamandi, Java, Indonesia, coll. F. Kopstein, II-1936.

Paratypes: ZMA 12103 (2), surroundings Buitenzorg, Java, Indonesia, coll. H. Boschma. Other paratypes in RMNH and SMF.

Draco quinquefasciatus longibarba Hennig, 1936: 193

Holotype: NMS, male, Baram, Sarawak, Borneo, Indonesia.

Paratypes: ZMA 11022-023 (2), male and female, Samarinda, Borneo, Indonesia, coll. H. A. Lorentz, 24.V.1909. Other paratypes in NMS and RMNH.

Draco toxopei De Jong, 1926a: 88

- Syntype: ZMA 10934 (1), male, Wa'katin, Buru, Indonesia, coll. L. J. Toxopeus, 1921; ZMA 10935 (1), male, Rana, Buru, Indonesia, coll. L. J. Toxopeus, 1921.
- Draco volans boschmai* Hennig, 1936: 180
 Holotype: ZMA 11025 (1), male, Maumeri, East-Flores, Indonesia, coll. M. Weber, 1888.
- Paratypes: ZMA 11026 (1), female, from type locality. ZMA 11027 (1) male, ZMA 11028 (1) female, Larantuka, Northcoast-Flores, Indonesia, coll. G. A. J. van de Sande, IV.1908; ZMA 11029 (1) female, Ende, Flores, Indonesia, coll. M. Weber, 1888; ZMA 11030 (1), male, ZMA 11031 (1), female, Sikka, Flores, Indonesia, coll. M. Weber, 1888. Other paratypes in RMNH.
- Eremias inornata* Roux, 1907a: 427
 Lectotype, designated by Daan & Hillenius (1966: 129), ZMA 11049, Oranje River, Little Namaqualand, South Africa, coll. M. Weber, IX.1894.
- Paralectotypes: ZMA 11050-055 (6), 3 adult and 3 juvenile specimens, from type locality. Other paralectotype in NMBA.
- Gehyra interstitialis* Oudemans, 1894: 134
 Holotype: ZMA unregistered, New Guinea, coll. R. Semon, 1893, could not be found in the collection.
- Gekko smaragdinus* Taylor, 1922b: 187
 Holotype: CAS 62336, Pollillo, Philippines, coll. E. H. Taylor, 12.VII.1920.
- Paratype: ZMA 11012 (1), from type locality. Other paratypes in EHT, FMNH, and MCZ.
- Gonatodes atricucullaris* Noble, 1921a: 135
 Holotype: MCZ 14042, Bellavista, Peru, coll. G. K. Noble, IX.1916.
- Paratypes: ZMA 11001-004 (4), Bellavista, Peru, coll. H. C. Raven. Other paratypes in MCZ, UINMH, UMMZ, and UNSM.
- Gymnodactylus darmandvillei* Weber, 1890: 163
 Syntypes: ZMA 10943 (1), ZMA 10944 (1), Sikka, East Flores, Indonesia, coll. M. Weber, 1888. Originally spelled as *Gymnodactylus d'armandvillei*.
- Gymnodactylus malayanus* De Rooy, 1915: 20
 Lectotype, designated by Daan & Hillenius (1966: 130), ZMA 11308, male, Indonesia, exact locality and collector unknown.
- Paralectotype: ZMA 11309 (1), female, from type locality.
- Gymnodactylus papuensis* Brongersma, 1934: 173
 Holotype: ZMA 10937, male, S. W. New Guinea, coll. Exploration Detachment, 1910-1911.
- Paratype: ZMA 10938 (1) female, from type locality. Other paratypes in MNHN.
- Gymnodactylus sermowaiensis* De Rooy, 1915: 9
 Holotype: ZMA 10910, male, upper Sermowai River, West New Guinea, coll. K. Gjellerup, V.1911.
- Gymnodactylus vankampeni* Brongersma, 1933b: 252
 Holotype: ZMA 10981, male, Modderlust, Northwest New Guinea, coll. P. N. van Kampen, 17.V.1910.
- Paratype: ZMA 10982 (1), female, from type locality.
- Hemidactylus vandermeermohri* Brongersma, 1928a: 1
 Holotype: ZMA 10939, male, Pulu Berhala, Deli, Sumatra, Indonesia, coll. J. C. van der Meer Mohr, VIII.1927.
- Paratype: ZMA 10940 (1), male, from type locality.
- Hemiphyllodactylus margarethae* Brongersma, 1931a: 11
 Holotype: ZMA 11095, Fort de Kock, Sumatra, Indonesia, coll. E. R. Jacobson, 1913.
- Paratypes: ZMA 11096 (1), from type locality. Other paratypes in IRSNB.
- Herpetosaura mira* Roux, 1907a: 435
 Lectotype, designated by Daan & Hillenius (1966: 131), ZMA 11036, Transvaal, South Africa, coll. Breyer, 1896.
- Paralectotypes: ZMA 11037-045 (9), from type locality. Other paralectotype in NMBA.
- Japalura nasuta* De Jong, 1930: 115
 Holotype: MZB, N. Borneo, Indonesia, coll. Mohari, 1912.
- Paratypes: ZMA 10989-990 (2), from type locality. Other paratypes in MZB.
- Lepidoblepharus barbouri* Noble, 1921a: 133
 Holotype: MCZ 14044, male, Perico N. of Bellavista, Peru, coll. G. K. Noble, 1916.
- Paratypes: ZMA 10996-999 (4), from type locality. Other paratypes in FMNH, MCZ, NMW, UINMH, UMMZ, and USNM.
- Lepidodactylus divergens* Taylor, 1918: 242
 Holotype: PNM 2026, females, Great Govenen Island, Philippines, coll. E. H. Taylor, 1917.
- Paratypes: ZMA 11014-015 (2), from type locality. Other paratypes in EHT, PNM, FMNH, and MCZ.
- Lygodactylus ocellatus* Roux, 1907a: 406
 Lectotype, designated by Daan & Hillenius (1966: 131), ZMA 11347, Pretoria District, South Africa, coll. Breyer, 1886.
- Paralectotypes: ZMA 11348-350 (3), from type locality. Other paralectotypes in MCZ and NMBA.
- Lygosoma acutirostre* Oudemans, 1894: 141
 Holotype: ZMA 11401, Saleyer, Indonesia, coll. M. Weber, 1889.
- Lygosoma beauforti* De Jong, 1927: 316
 Holotype: ZMA 12078, Parana Valley, N. W. New Guinea, coll. P. Wirz, 1922.
- Lygosoma derooyae* De Jong, 1927: 313
 Holotype: ZMA 11310, Doorman Peak, altitude 1450 m, W. New Guinea, coll. P. Wirz, 1922.
- Lygosoma florense* Weber, 1890: 173
 Syntypes: ZMA 11056-058 (3), Sikka, Flores, Indonesia, coll. M. Weber, 1888; ZMA 11059-064 (6), Maumeri, Flores, Indonesia, coll. M. Weber, 1888; ZMA unregistered, 2 speci-

- mens, Maumeri, Flores, Indonesia, could not found in the collection.
- Lygosoma kuekenthali* Boettger, 1895: 117
Lectotype, designated by Mertens (1922:175), SMF 15202, Soa Konorra, Halmaheira, Indonesia, coll. W. Kükenenthal, 1894.
Paralectotype: ZMA 11007 (1), from type locality. Other paralectotypes in MCZ and SMF.
- Lygosoma kuekenthali notomoluccense* Brongersma, 1931b: 335
Holotype: ZMA 11090, Ambon, Indonesia.
Paratypes: ZMA 11091-092 (2), Kelang, Indonesia, coll. F. Kopstein, VII.1923; ZMA 11094 (1), Sapa-rua, Indonesia, coll. F. Kopstein, 28.V.1923. Other paratypes in BMNH and ZSM (1).
- Lygosoma laeve* Oudemans, 1894: 144
Holotype: ZMA 10994, Cooktown, Queensland, Australia, coll. R. Semon, 1893.
- Lygosoma longicaudatum* De Rooy, 1915: 220
Holotype: ZMA 11033, Alkmaar, Lorentz River, W. New Guinea, coll. H. A. Lorentz, XI.1909.
- Lygosoma minutum obtusirostrum* De Jong, 1927: 312
Syntypes: ZMA 11442, Upper Sermowai River, W. New Guinea, coll. K. Gjellerup, 29.III.1911; ZMA 11443 (1), ZMA 12090 (1), Prauwenbivak, Idenburg River, W. New Guinea, coll. W. C. van Heurn, IX.1920.
- Lygosoma minutum rotundirostrum* De Jong, 1927: 313
Syntypes: ZMA 11406 (1), between Bronbeek and Modderlust, W. New Guinea, coll. P. N. van Kampen, 17.V.1910; ZMA 11407 (1), Prauwenbivak, Idenburg River, W. New Guinea, coll. W. C. van Heurn, IX.1920; ZMA unregistered (1), Parana Valley, New Guinea, coll. P. Wirz, 1922, could not be found in the collection.
- Lygosoma mivarti obscurum* De Jong, 1927: 317
Syntypes: ZMA 11445 (1), Prauwenbivak, Idenburg River, W. New Guinea, coll. W. C. van Heurn, IX 1920; ZMA 11446 (1), from type locality; ZMA unregistered (6), 2 from Prauwenbivak, Idenburg river, 3 from Pioniersbivak, Mam-beramo river, coll. W. C. van Heurn, 1920, 1 from northern New Guinea, coll. K. Gjellerup, 1911, could not be found in the collection.
- Lygosoma nigriventre* De Rooy, 1915: 214
Lectotype, designated by Daan & Hillenius (1966: 133), ZMA 11019, Hellwig Mountains, W. New Guinea, altitude 1700 m, coll. H. A. Lorentz, 7.IX.1907.
Paralectotypes: ZMA 11020-021 (2), Hellwig Mountains, West New Guinea, coll. H. A. Lorentz, 16.X 1909.
- Lygosoma semoni* Oudemans, 1894: 142
Holotype: ZMA 10993, New Guinea, coll. R. Semon, 1893.
- Lygosoma smaragdinum celebense* De Rooy, 1915: 201
Lectotype, designated by Daan & Hillenius (1966: 134), ZMA 11402, Posso, Celebes, Indonesia, coll. A. C. Kruyt, 1898.
Paralectotype: ZMA 11403, from type locality. Other paralectotype in RNMH.
- Lygosoma smaragdinum pisangense* Brongersma, 1931a: 24
Holotype: IRSNB, Tartaruga, Pisang Islands, Indonesia coll. S. M. Léopold de Belgique, 17.III.1929.
Paratype: ZMA 11307 (1), from type locality.
- Lygosoma striolatum* Weber, 1890: 172
Holotype: ZMA 10945, Reo, West Flores, Indonesia, coll. M. Weber, 1888.
- Lygosoma unilineatum* De Rooy, 1915: 208
Lectotype, designated by Daan & Hillenius (1966: 134), ZMA 10912, Tanah Merah Bay, W. New Guinea, coll. K. Gjellerup, 27.VIII.1911.
Paralectotype: ZMA 10913 (1), from type locality.
- Lygosoma verreauxii biunguiculata* Oudemans, 1894: 145
Syntypes: ZMA 11398-399 (2), Burnett River, Queensland, Australia, coll. R. Semon, 1893; ZMA unregistered (3), could not be found in the collection.
- Mabuya englei* Taylor, 1925: 101
Holotype: MCZ 26289, Saub, Cotabato Province, Mindanao, Philippines.
Paratypes: ZMA 11244-245 (2), from type locality. Other paratypes in EHT, MCZ, and UINMH.
- Neusticurus dejongi* Brongersma, 1927: 543
Holotype: ZMA 10241, male, Surinam, coll. Bruegel, 1890.
- Pachydactylus weberi* Roux, 1907a: 408
Lectotype, designated by Daan & Hillenius (1966: 135), ZMA 11046, Klipfontein, Little Namaqualand, South Africa, coll. M. Weber, IX.1894.
Paralectotypes: ZMA 11047-048 (2), from type locality. Other paralectotype in NMBA.
- Phyllodactylus magister* Noble, 1924: 110
Holotype: MCZ 17974, Perico, Chinchipe River, Peru.
Paratypes: ZMA 11216-219 (4), Bellavista, Peru, coll. G. K. Noble, IX.1916. Other paratypes in FMNH, MCZ, UINMH, UMMZ, and UNSM.
- Phyllodactylus muelleri* Parker, 1935: 483
Holotype: BMNH 1934.7.5.1., male, Patos island, Venezuela, coll. D. Vesey-Fitzgerald. Originally spelled *Phyllodactylus mülleri*.
Paratypes: ZMA 11005 male (1), ZMA 11006 juv (1), Maratay, Venezuela, coll. C. Vogl.
- Phyllodactylus rutteni* Wagenaar Hummelinck, 1940: 77
Holotype: RMNH 21973, Norte de Valuchu, Isla Blanquilla, Venezuela, coll. P. Wagenaar Hummelinck, 21.VII.1936.
Paratypes: ZMA 10924 (1), El Jaque, Isla Blanquilla, Venezuela, coll. P. Wagenaar Hummelinck, 22.VII.1936. ZMA 10925, Tortuga, Gran Roque, Los Roques, Venezuela, coll. P. Wagenaar Hummelinck, 25.VII.1936. Other paratypes in RMNH.

Scolecosaurus alleni Barbour, 1914: 315

Holotype: MCZ 7793, St. George, Grenada, British West Indies, coll. G. M. Allen, 1910.

Paratype: ZMA 11405 (1), from type locality. Other paratypes in MCZ.

Sepsina weberi Roux, 1907a: 437

Holotype: ZMA 10975, Steinkopf, Little Namaqualand, South Africa, coll. M. Weber, 1894.

Sphenomorphus leptofasciatus Greer & Parker, 1974: 50

Holotype: MCZ 108573, Chimbu district, Kundiawa, 5200 ft, Papua New Guinea, coll. F. Parker, 6.III.1968.

Paratypes: ZMA 15152 (10), from type locality, coll. F. Parker, 10.IV.1967. Other paratypes in AMNH, CAS and MCZ.

Sphenomorphus tanneri Greer & Parker, 1967: 1

Holotype: MCZ 76551, Kunua, Bougainville, coll. F. Parker, 8.VI.1963.

Paratypes: ZMA 15151 (10) from type locality, coll. F. Parker, 17.VII.1963. Other paratypes in AMNH, and MCZ.

Tetradactylus breyeri Roux, 1907a: 430

Holotype: ZMA 10974, Transvaal, South Africa, coll. Breyer, 1896.

Tiliqua gigas keiensis Oudemans, 1894: 138

Lectotype, designated by Daan & Hillenius (1966: 136), ZMA 10991, Kei Islands, Indonesia, coll. A.C. Wertheim, 1888.

Paralectotype: ZMA 10992, from type locality.

Tropidophorus stejneri Taylor, 1922b: 285

Holotype: CAS 60230, Abung Abung, Basilan Island, Philippines, coll. E. H. Taylor, 22.X.1920.

Paratypes: ZMA 11448-449 (2), ZMA 18578 (2), from type locality. Other paratypes in FMNH and MCZ.

REPTILIA, OPHIDIA

Boiga dendrophila occidentalis Brongersma, 1934: 211

Holotype: ZMA 10986, female, Nias, Indonesia, coll. J. P. Kleiweg de Zwaan, 1910.

Paratypes: ZMA 10987 (1), from type locality; ZMA 10988 (1) male, Airbangis, Padang lowlands, Sumatra, Indonesia, coll. E. Jacobson, 1913. Other paratypes in RMNH and UNSM.

Calamaria bogorensis Van Lidth de Jeude, 1890: 182

Syntypes: ZMA 10102 (2), Buitenzorg, Java, Indonesia, coll. M. Weber, 1888. Other syntypes in FMNH, MCZ, and RMNH.

Calamaria ceramensis De Rooy, 1913: 26

Syntypes: ZMA 10083, 2 specimens, Honitétu, W. Ceram, Indonesia, coll. L. F. de Beaufort, II.1910.

Calamaria doerianense Brongersma, 1928b: 255

Holotype: ZMA 11010, Doerian Island, Riouw Archipelago, Indonesia, coll. K. W. Dammerman, 11.XI.1923.

Calamaria elegans De Rooy, 1917: 158

Holotype: ZMA 11017, Ajer Dingin, Simalur, Indonesia, coll. E. R. Jacobson, VII.1913.

Calamaria forcarti Inger & Marx, 1965: 184

Holotype: NMW 16710, male, Deli, Sumatra, Indonesia, don. Steindachner, 1900.

Paratype: ZMA 10073 (1), Lelewoea, Nias, Indonesia, coll. J.P. Kleiweg de Zwaan, 1910-1911. Other paratypes in NMBA and NMW, cf. Tiedemann & Häupl (1980: 51).

Calamaria hilleniusi Inger & Marx, 1965: 96

Holotype: ZMA 10078, male, Samarinda, Borneo, Indonesia, coll. H. A. Lorentz, 24.V.1909. Paratypes in FMNH and SMK.

Calamaria javanica lineata Brongersma, 1928b: 256

Holotype: ZMA 11011, Long Petah, Central East Borneo, Indonesia, coll. H. C. Siebers, 15.IX.1925.

Calamaria lautensis De Rooy, 1917: 163

Lectotype designated by Inger & Marx (1965: 70), RMNH 4716a, Pulu si Laut, Cocos islands, Indonesia, coll. E. R. Jacobson, VI.1913.

Paralectotypes: ZMA 10111 (1), from type locality, coll. E. R. Jacobson, 21.VIII.1913, ZMA 10112 (1), Abail, Simalur, Indonesia, coll. E. R. Jacobson, IV.1913. Other paralectotype in RMNH 4716b, from type locality, abusively designated as lectotype by Daan & Hillenius (1966: 137).

Calamaria simalurensis De Rooy, 1917: 159

Holotype: ZMA 11013, Sibigo, Simalur, Indonesia, coll. E. R. Jacobson, VIII.1913.

Calamaria sumatrana Edeling, 1870: 379

Neotype: designated by Inger & Marx (1965: 181), ZMA 10237, Deli, Medan, Sumatra, Indonesia, coll. L. Ph. de Bussy, 1905-1920.

Cantoria annulata De Jong, 1926b: 304

Holotype: ZMA 11065, Prins Frederik Hendrik island, West New Guinea, coll. Dutch New Guinea Expedition, II.1910.

Chrysopelea paradisi celebensis Mertens, 1968: 206

Holotype: NMBA 5074, female, Kema, Minahassa, N.E. Celebes, Indonesia, coll. P. & F. Sarasin, 1894.

Paratypes: ZMA 13227 (2) male, female, Posso, Celebes, Indonesia, coll. A. C. Kruyt, 1898; ZMA 13228 (1) male, Celebes, Indonesia, coll. S. C. J. W. Musschenbroek, 1884. Other paratypes in NMBA and ZMB.

Crotalus horridus unicolor Van Lidth de Jeude, 1887: 133

Holotype: RMNH 613, Aruba, Dutch Leeward islands, coll. J. R. H. Neervoort van de Poll, 1885. Died in Zoo, Natura Artis Magistra, Amsterdam.

Paratypes: ZMA 11974 (1), ZMA 11169 (1), from type locality. Other paratype in RMNH.

Diadophis bipunctatus Van Lidth de Jeude, 1890: 184

Holotype: ZMA 11032, Kaju tanam, Sumatra, Indone-

- sia, coll. M. Weber, 1888.
- Elaps thepassi* Bleeker, 1859: 201
 Syntype: ZMA 11370 (1), Sintang, Indonesia, collection P. Bleeker, auction lot number 141, see Hubrecht (1879: 59).
- Enygrus asper schmidti* Stull, 1932: 26
 Holotype: MCZ 29778, Kaiserin Sepik, Augusta River, New Guinea, coll. K. Gjellerup, 1910.
 Paratype: ZMA 16934 (1), from type locality. Other paratypes in FMNH and UMMZ.
- Heurnia ventromaculata* De Jong, 1926b: 303
 Holotype: ZMA 11066, Pioniersbivak, Mamberamo river, N. W. New Guinea, coll. W. C. van Heurn, 12.VII.1920.
- Hydrophis caeruleus thai* Smith, 1920: 18
 Holotype: BMNH 1921.2.11.83, Hua Hin, Gulf of Siam, coll. M. A. Smith, VI.1917.
 Paratypes: ZMA 12227 (1), Meklong, Gulf of Siam, coll. M. A. Smith, VI.1917; NMS, Gulf of Siam and Cap St. Jacques, coll. M. A. Smith, VI.1917.
- Hydrophis torquatus aagaardi* Smith, 1920: 14
 Holotype: BMNH, 1921.2.11.44, Coast of Bangnara, Patani, Gulf of Siam, coll. C. J. Aagaard, VII.1917.
 Paratype: ZMA 12226 (1), from type locality. Other paratype in NMS.
- Leimadophis poecilogyrus reticulatus* Parker, 1931: 285
 Syntype: ZMA 11400, Province Sara, E. Bolivia, altitude 600 m. Other syntypes in BMNH.
- Lycodon florensis* De Rooy, 1917: 110
 Holotype: ZMA 11080, near Kotting along upper Maumeri River, Flores, Indonesia, altitude 220 m, coll. G. A. J. van der Sande, 14.VII.1908.
- Naja naja philippinensis* Taylor, 1922a: 265
 Holotype: PNM, destroyed in World War II, Brown & Alcalá (1980: 29).
 Paratype: ZMA 11269 (1), juv. hatched in Taylor's laboratory, Manila, Philippines. Other paratypes in CAS and CM.
- Natrix mairii multiscutellata* Brongersma, 1948: 372
 Holotype: RMNH 8669, male, Alkmaar, Lorentz river, West New Guinea, coll. H. A. Lorentz, 1909.
 Paratypes: ZMA 11081 (1) male, ZMA 11082 (1) female, Prauwenbivak, Idenburg River, New Guinea, coll. W. C. van Heurn, 14.IX.1920; ZMA 11083 (1), female, same locality as ZMA 11081, coll. Lt. Drost, X.1920; ZMA 11084 (1), female, Pioniersbivak, Mamberamo River, New Guinea, coll. W. C. van Heurn, 17.VIII.1920; ZMA 11085 (1), male, Betewi River, New Guinea, coll. P. N. van Kampen, 20.VI.1910; ZMA 11086 (1), female, Begowre River below zoutbron, New Guinea, coll. P. N. van Kampen, 3.VII.1910; ZMA 11087 (1), female, Zoutbron, New Guinea, coll. P. N. van Kampen, VII.1910; ZMA 11088 (1), female, between Modderlust on the Tami River and Kasawari on the coast, New Guinea, coll. P. N. van Kampen, 18.V.1910; ZMA 11089 (1), female, Hollandia, New Guinea, coll. Sain, 1910; ZMA 11097 (1), male, Hollandia, New Guinea, coll. P. N. van Kampen, 1910. Other paratypes in BMNH, MCZ, MZB, NMW, and RMNH.
- Psammophis longimentalis* Roux, 1907b: 736
 Holotype: ZMA 11098, Buffels river, South Africa, coll. M. Weber, 1894.
- Trigonocephalus blomhoffii* Boie, 1826.
 Lectotype, selected by Hoogmoed (1978: 99), ZMA 15179, environments of Nagasaki, Japan.
- Tropidonophis statisticus* Malnate & Underwood, 1988: 155
 Holotype: MCZ 142422, Orumba, Eastern Highlands Province, Papua New Guinea.
 Paratypes: ZMA 11432 (1), Cycloop Mountain, Irian Jaya, 21.VIII.1911. Other paratypes in AM, AMNH, BMNH, CAS, MCZ, NMW, RMNH, and UNSM.
- Typhlops florensis brongersmai* Mertens, 1929a: 294
 Holotype: ZMA 11179, male, Mao Marru, E. Sumba, Indonesia, altitude 450 m, coll. K. W. Dammersman, 5.V.1925. Paratype in SMF.
- Typhlops koekkoeki* Brongersma, 1934: 186
 Holotype: ZMA 11077, Bunju Island, near N. Borneo, Indonesia, coll. U. Driebergen, 1924.
- Typhlops soensis* De Jong, 1930: 117
 Holotype: MZB, Soë, S. Timor, Indonesia, altitude 880 m, coll. Mrs. Walsh, II.1929.
 Paratypes: ZMA 11078-79, from type locality. Other paratypes in MZB.
- Vipera latastei montana* Saint Girons, 1953a: 24
 Syntype: ZMA 11154 (1), Massif du Toubkal, Haut Atlas, Morocco, altitude 2800 m, coll. H. Saint Girons, 1952.
 Other syntypes in MHNG and MNHN.

INDEX

- aagaardi, *Hydrophus torquatus*134
Ablepharus boutonii furcata130
Ablepharus boutonii intermedius130
Ablepharus boutonii keiensis130
Ablepharus burnetti128
 acanthi, *Rana*131
 acutirostre, *Lygosoma*130
Agama stellio cypriaca130
 aguilina, *Ameiva*129
 albopunctata, *Sphenophryne*133
 alleni, *Scolecosaurus*126
 almogavarii, *Alytes obstetricans*126
Alytes dickhilleni126
Alytes obstetricans almogavarii126
Ameiva aguilina130
 anguinus parkelj, *Proteus*129
 angustopalmata, *Rana tigrina*129
 annulata, *Cantoria*133
 anodon, *Rhacophorus*129
Anolis bonairensis blanquillanus130
Anolis conspersus130
 asper schmidti, *Enygrus*134
Asterophrys minima126
 atricucularis, *Gonatodes*131
Austrochaperina robusta126
Bachia intermedia130
 barbouri, *Dicrodon*130
 barbouri, *Lepidoblepharus*131
 basipalmata, *Chaperina*127
 beauforti, *Cornufer*127
 beauforti, *Lygosoma*131
 bidens, *Xenorhina*129
 bifaciatus, *Rhacophorus*129
 billitonensis, *Ichthyophis*129
 bipunctatus, *Diadophis*133
 biunguiculata, *Lygosoma vereauxi*132
 blanquillanus, *Anolis bonairensis*130
 blomhoffii, *Trigonocephalus*134
 bogorensis, *Calamaria*133
Boiga dendrophila occidentalis133
 bonairensis blanquillanus, *Anolis*130
Boophis hillenii126
Boophis mandraka126
Boophis reticulatus126
Boophis viridis127
 boschmai, *Draco volans*131
 boutonii furcata, *Ablepharus*130
 boutonii intermedius, *Ablepharus*130
 boutonii keiensis *Ablepharus*130
Brachymeles pathfinderi130
Brachymeles vermis130
 brevicrus, *Oxydactyla*128
 breyeri, *Tetradactylus*133
 brongersmai, *Typhlops florensis*134
Bufo mcgregori127
 burnetti, *Ablepharus*130
 buruensis, *Draco*130
 caeruleus thai, *Hydrophus*134
Calamaria bogorensis133
Calamaria ceramensis133
Calamaria doerianense133
Calamaria elegans133
Calamaria forcarti133
Calamaria hilleniusi133
Calamaria javanica lineata133
Calamaria lautensis133
Calamaria simalurensis133
Calamaria sumatrana133
Cantoria annulata133
 celebense, *Lygosoma smaragdinum*132
 celebensis, *Chrysopeles paradisi*133
 ceramensis, *Calamaria*133
 ceratophthalmus, *Chaperina*127
Chamaeleon incomutus130
Chamaeleon laterispinnis130
Chaperina basipalmata127
Chaperina ceratophthalmus127
Chaperina punctata127
Choerophryne proboscidea127
Chrysopeles paradisi celebensis133
 conspersus, *Anolis*130
Cophixalus crucifer127
Cornufer beauforti133
Crotalus horridus unicolor127
 crucifer, *Cophixalus*130
 cypriaca, *Agama stellio*131
 darmandvillei, *Gymnodactylus*131
 d'armandvillei, *Gymnodactylus*131
 debussyi, *Rana*128
 dejongi, *Neusticurus*132
dendrophila occidentalis, *Boiga*133
 deroooyae, *Lygosoma*131
Diadophis bipunctatus133
 dickhilleni, *Alytes*126
Dicrodon barbouri130
 diminutiva, *Micrixalus*128
 divergens, *Lepidodactylus*131
 doerianense, *Calamaria*133
Draco buruensis130
Draco fimbriatus hennigi130
Draco quinquefasciatus longibarba130
Draco toxopei130
Draco volans boschmai131
Dyscophina volzi127
Elaps thepassi134
 elegans, *Calamaria*133
Eleutherodactylus zeuctotylus127
 englei, *Mabuya*132

<i>Enygrus asper schmidti</i>	134	<i>laterispinnis, Chamaeleon</i>	130
<i>Eremias inornata</i>	131	<i>lautensis, Calamaria</i>	133
<i>fallax, Rana</i>	128	<i>Lechroides platyceps</i>	127
<i>fimbriatus hennigi, Draco</i>	130	<i>Leimadophis poecilogyrus reticulatus</i>	134
<i>flava, Oreophryne</i>	128	<i>Lepidoblepharus barbouri</i>	131
<i>flavobrunneus, Mantidactylus</i>	128	<i>Lepidodactylus divergens</i>	131
<i>florense, Lygosoma</i>	131	<i>leptofasciatus, Sphenomorphus</i>	133
<i>florensis brongersmai, Typhlops</i>	134	<i>lineata, Calamaria javanica</i>	133
<i>florensis, Lycodon</i>	134	<i>linki, Polypedates</i>	128
<i>forcarti, Calamaria</i>	133	<i>longibarba, Draco quinquefasciatus</i>	130
<i>furcata, Ablepharus boutonii</i>	130	<i>longicaudatum, Lygosoma</i>	132
<i>Gehyra interstitialis</i>	131	<i>longimentalis, Psammophis</i>	134
<i>Gekko smaragdinus</i>	131	<i>Lycodon florensis</i>	134
<i>gigantea, Xenorhina</i>	129	<i>Lygodactylus ocellatus</i>	131
<i>gigas keiensis, Tiliqua</i>	133	<i>Lygosoma acutirostre</i>	131
<i>Gonatodes atricucullaris</i>	131	<i>Lygosoma beauforti</i>	131
<i>grisea, Rana</i>	128	<i>Lygosoma derooyae</i>	131
<i>guerreroi, Rana</i>	128	<i>Lygosoma florense</i>	131
<i>Gymnodactylus d'armandvillei</i>	131	<i>Lygosoma kuekenthali</i>	132
<i>Gymnodactylus malayanus</i>	131	<i>Lygosoma kuekenthali notomoluccense</i>	132
<i>Gymnodactylus papuensis</i>	131	<i>Lygosoma laeve</i>	132
<i>Gymnodactylus sermowaiensis</i>	131	<i>Lygosoma longicaudatum</i>	132
<i>Gymnodactylus vankampeni</i>	131	<i>Lygosoma minutum obtusirostrum</i>	132
<i>Hemidactylus vandermeermohri</i>	131	<i>Lygosoma minutum rotundirostrum</i>	132
<i>Hemiphyllodactylus margarethae</i>	131	<i>Lygosoma mivarti obscurum</i>	132
<i>hennigi, Draco fimbriatus</i>	130	<i>Lygosoma nigriventre</i>	132
<i>Herpetosaura mira</i>	131	<i>Lygosoma semoni</i>	132
<i>Heurnia ventromaculata</i>	134	<i>Lygosoma smaragdinum celebense</i>	132
<i>hillenii, Boophis</i>	126	<i>Lygosoma smaragdinum pisangense</i>	132
<i>hilleniusi, Calamaria</i>	133	<i>Lygosoma striolatum</i>	132
<i>horridus unicolor, Crotalus</i>	133	<i>Lygosoma unilineatum</i>	132
<i>Hydrophus caeruleus thai</i>	134	<i>Lygosoma vereauxii biunguiculata</i>	132
<i>Hydrophus torquatus aagaardi</i>	134	<i>Mabuya englei</i>	132
<i>Hyla papua</i>	127	<i>macra, Metopostira</i>	128
<i>Hyla proboscidea</i>	127	<i>macrops, Xenorhina</i>	129
<i>Hyla rhacophorus</i>	127	<i>magister, Phyllodactylus</i>	132
<i>Hyla sanguinolente</i>	127	<i>mairii multiscutellata, Natrix</i>	134
<i>Hylophorbus variegatus</i>	127	<i>malayanus, Gymnodactylus</i>	131
<i>Ichthyophis billitonensis</i>	129	<i>mandraka, Boophis</i>	126
<i>igorata, Rana</i>	128	<i>Mantella pulchra</i>	127
<i>incomutus, Chamaeleon</i>	130	<i>Mantidactylus flavobrunneus</i>	128
<i>inornata, Eremias</i>	131	<i>Mantidactylus punctatus</i>	128
<i>intermedia, Bachia</i>	130	<i>margarethae, Hemiphyllodactylus</i>	131
<i>intermedius, Ablepharus boutonii</i>	130	<i>mcgregori, Bufo</i>	127
<i>interstitialis Gehyra</i>	131	<i>Metopostira macra</i>	128
<i>Ixalus jacobsoni</i>	127	<i>Micrixalus diminutiva</i>	128
<i>jacobsoni Ixalus</i>	127	<i>Microhyla niasensis</i>	128
<i>Japalura nasuta</i>	131	<i>Microhyla superciliaris</i>	128
<i>javanica lineata, Calamaria</i>	133	<i>microtypanum, Rana</i>	128
<i>Kaloula rigida</i>	127	<i>minima, Asterophrys</i>	126
<i>keiensis Ablepharus boutonii</i>	130	<i>minutum obtusirostrum, Lygosoma</i>	132
<i>keiensis, Tiliqua gigas</i>	133	<i>minutum rotundirostrum, Lygosoma</i>	132
<i>koekkoeki, Typhlops</i>	134	<i>mira, Herpetosaura</i>	131
<i>kuekenthali, Lygosoma</i>	132	<i>mivarti obscurum, Lygosoma</i>	132
<i>kuekenthali notomoluccense, Lygosoma</i>	132	<i>Molge pyrrhogaster</i>	129
<i>laeve, Lygosoma</i>	132	<i>montana, Vipera latastei</i>	134
<i>latastei montana, Vipera</i>	134	<i>muelleri, Phyllodactylus</i>	132

mülleri, <i>Phyllodactylus</i>	132	<i>Rana microtypanum</i>	128
multiscutellata, <i>Natrix mairii</i>	134	<i>Rana novaeguineae</i>	129
<i>Naja naja philippinensis</i>	134	<i>Rana pantherina</i>	129
<i>naja philippinensis</i> , <i>Naja</i>	134	<i>Rana tigrina angustopalmata</i>	129
<i>nasuta</i> , <i>Japalura</i>	131	<i>Rana waigeensis</i>	129
<i>Natrix mairii multiscutellata</i>	134	<i>Rana yakani</i>	129
<i>Nectophryne sumatrana</i>	128	<i>reticulatus</i> , <i>Boophis</i>	126
<i>Neusticurus dejongi</i>	132	<i>reticulatus</i> , <i>Leimadophis poecilogyrus</i>	134
<i>niasensis</i> , <i>Microhyla</i>	128	<i>Rhacophorus anodon</i>	129
<i>nigriventre</i> , <i>Lygosoma</i>	132	<i>Rhacophorus bifaciatus</i>	129
<i>notomoluccense</i> , <i>Lygosoma kuekenthali</i>	132	<i>rhacophorus</i> , <i>Hyla</i>	127
<i>novaeguineae</i> , <i>Phanerotis</i>	128	<i>rigida</i> , <i>Kaloula</i>	127
<i>novaeguineae</i> , <i>Rana</i>	129	<i>robusta</i> , <i>Austrochaperina</i>	126
<i>obscurum</i> , <i>Lygosoma mivarti</i>	132	<i>rotundirostrum</i> , <i>Lygosoma minutum</i>	132
<i>obstetricans almogavarii</i> , <i>Alytes</i>	126	<i>rutteni</i> , <i>Phyllodactylus</i>	132
<i>obtusirostrum</i> , <i>Lygosoma minutum</i>	132	<i>sanguinolente</i> , <i>Hyla</i>	127
<i>occidentalis</i> , <i>Boiga dendrophila</i>	133	<i>Scolecosaurus alleni</i>	133
<i>ocellata</i> , <i>Xenorhina</i>	129	<i>schmidti</i> , <i>Enygrus asper</i>	134
<i>ocellatus</i> , <i>Lygodactylus</i>	131	<i>semoni</i> , <i>Lygosoma</i>	132
<i>Oedipina pseudouniformis</i>	129	<i>Sepsina weberi</i>	133
<i>Oreophryne flava</i>	128	<i>sermowaiensis</i> , <i>Gymnodactylus</i>	131
<i>Oxydactyla brevicrus</i>	128	<i>simalurensis</i> , <i>Calamaria</i>	133
<i>Pachydactylus weberi</i>	132	<i>smaragdinum celebense</i> , <i>Lygosoma</i>	132
<i>pantherina</i> , <i>Rana</i>	129	<i>smaragdinum pisangense</i> , <i>Lygosoma</i>	132
<i>papua</i> , <i>Hyla</i>	127	<i>smaragdinus</i> , <i>Gekko</i>	131
<i>papuensis</i> <i>Gymnodactylus</i>	131	<i>soensis</i> , <i>Typhlops</i>	134
<i>paradisi celebensis</i> , <i>Chrysopeles</i>	133	<i>Sphenomorphus leptofasciatus</i>	133
<i>parkelj</i> , <i>Proteus anguinus</i>	129	<i>Sphenomorphus tanneri</i>	133
<i>pathfinderi</i> , <i>Brachymeles</i>	130	<i>Sphenophryne albopunctata</i>	129
<i>Phanerotis novaeguineae</i>	128	<i>statisticus</i> <i>Tropidonophis</i>	134
<i>Philautus pollilensis</i>	128	<i>stejnegeri</i> , <i>Tropidophorus</i>	133
<i>Philautus williamsi</i>	128	<i>stellio cypriaca</i> , <i>Agama</i>	130
<i>philippinensis</i> , <i>Naja naja</i>	134	<i>striolatum</i> , <i>Lygosoma</i>	132
<i>Phyllodactylus magister</i>	132	<i>sumatrana</i> , <i>Calamaria</i>	133
<i>Phyllodactylus mülleri</i>	132	<i>sumatrana</i> , <i>Nectophryne</i>	128
<i>Phyllodactylus rutteni</i>	132	<i>superciliaris</i> , <i>Microhyla</i>	128
<i>pisangense</i> , <i>Lygosoma smaragdinum</i>	132	<i>tanneri</i> , <i>Sphenomorphus</i>	133
<i>platyceps</i> , <i>Lechroides</i>	127	<i>Tetradactylus breyeri</i>	133
<i>poecilogyrus reticulatus</i> , <i>Leimadophis</i>	134	<i>thai</i> , <i>Hydrophus caeruleus</i>	134
<i>pollilensis</i> , <i>Philautus</i>	128	<i>thepassii</i> , <i>Elaps</i>	134
<i>Polypedates linki</i>	128	<i>tigrina angustopalmata</i> , <i>Rana</i>	129
<i>proboscidea</i> , <i>Choerophryne</i>	127	<i>Tiliqua gigas keiensis</i>	133
<i>proboscidea</i> , <i>Hyla</i>	127	<i>torquatus aagaardi</i> , <i>Hydrophus</i>	134
<i>Proteus anguinus parkelj</i>	129	<i>toxopei</i> , <i>Draco</i>	130
<i>Psammophis longimentalis</i>	134	<i>Trigonocephalus blomhoffii</i>	134
<i>pseudouniformis</i> , <i>Oedipina</i>	129	<i>Tropidonophis statisticus</i>	134
<i>pulchra</i> , <i>Mantella</i>	127	<i>Tropidophorus stejnegeri</i>	133
<i>punctata</i> , <i>Chaperina</i>	127	<i>Typhlops florensis brongersmai</i>	134
<i>punctatus</i> , <i>Mantidactylus</i>	128	<i>Typhlops koekkoeki</i>	134
<i>pyrrhogaster</i> , <i>Molge</i>	129	<i>Typhlops soensis</i>	134
<i>quinquefasciatus longibarba</i> , <i>Draco</i>	130	<i>unicolor</i> , <i>Crotalus horridus</i>	133
<i>Rana acanthi</i>	128	<i>unilineatum</i> , <i>Lygosoma</i>	132
<i>Rana debussyi</i>	128	<i>vandermeermohri</i> , <i>Hemidactylus</i>	131
<i>Rana fallax</i>	128	<i>vankampeni</i> , <i>Gymnodactylus</i>	131
<i>Rana grisea</i>	128	<i>variegatus</i> , <i>Hylophorbus</i>	127
<i>Rana guerreroi</i>	128	<i>ventromaculata</i> , <i>Heurnia</i>	134
<i>Rana igorata</i>	128	<i>vereauxii biunguiculata</i> , <i>Lygosoma</i>	130

vermis, <i>Brachymeles</i>	130
<i>Vipera latastei montana</i>	134
viridis, <i>Boophis</i>	127
volans boschmai, <i>Draco</i>	131
volzi, <i>Dyscophina</i>	127
waigeensis, <i>Rana</i>	129
weberi, <i>Pachydactylus</i>	132
weberi, <i>Sepsina</i>	133

williamsi, <i>Philautus</i>	128
<i>Xenorhina bidens</i>	129
<i>Xenorhina gigantea</i>	129
<i>Xenorhina macrops</i>	129
<i>Xenorhina ocellata</i>	129
yakani, <i>Rana</i>	129
zeuctotylus, <i>Eleutherodactylus</i>	127

REFERENCES

- Arntzen, J.W. & M. Garcia-Paris, 1994. Morphological and allozyme studies of Midwife Toads (Genus *Alytes*), Including the description of two new Taxa from Spain. *Contributions to Zoology*, 65 (1): 5-34.
- Barbour, T., 1914. A contribution to the zoogeography of the West Indies, with special reference to amphibians and reptiles. *Mem. Mus. Comp. Zool. Harv.* 44 (2): 209-3.
- Barbour, T. & G.K. Noble, 1915. A revision of the lizard of the genus *Ameiva* *Bull. Mus. Comp. Zool. Harv.* 59 (6): 417-479.
- Barbour, T. & A. Loveridge, 1929. Typical reptiles and amphibians. *Bull. Mus. Comp. Zool. Harv.* 69: 205-360.
- Barbour, T. & A. Loveridge, 1946. First supplement to Typical reptiles and amphibians. *Bull. Mus. Comp. Zool. Harv.* 96 (2): 58-214.
- Bleeker, P., 1859. Reptilien van Sintang. *Natuurk. Tijdschr. Ned. Ind., [Wetlevreden]*, 20: 200-201.
- Blommers-Schlösser, R. M. A., 1979a. Biosystematics of the Malagasy frogs. I. Mantellinae (Ranidae). *Beaufortia*, 29 (352): 1-77.
- Blommers-Schlösser, R. M. A., 1979b. Biosystematics of the Malagasy frogs. II. The Genus *Boophis* (Rhacophoridae). *Bijdr. Dierk.*, 49 (2): 261-312, pls. 1-4.
- Boettger, O., 1895. Liste der Reptilien und Batrachier der Insel Halmaheira nach den Sammlungen Prof. Dr. W. Kükenthals. I. Reptilien *Zool. Anz. Leipzig*, 18: 116-121.
- Boie, H., 1826. Merkmale einiger Japanischen Lurche. *Isis*,: 203-216.
- Brame, A. H., 1968. Systematics and evolution of the mesoamerican salamander genus *Oedipina*. *J. Herpet.*, 2 (1/2): 1-64.
- Brongersma, L. D., 1927. *Neusticurus dejongi*, sp. n., a new lizard from Surinam. *Ann. Mag. nat. Hist., London*, (9) 20: 543-545.
- Brongersma, L. D., 1928a.. Lizards from Pulu Berhala. *Misc. zool. Sumatra., Medan*, 26: 1-3.
- Brongersma, L. D., 1928b. Neue Reptilien aus dem Zoologischen Museum Amsterdam. *Zool. Anz. Leipzig*, 75 (11/12): 251-257.
- Brongersma, L. D., 1931a.. Reptilia. In: *Résultats scientifiques du voyage aux Indes Orientales Néerlandaises de LL.AA.RR. le Prince et la Princesse Léopold de Belgique*. *Mém. Mus. roy. Hist. nat. Belg., Bruxelles (hors série)*, 5 (2): 3-39.
- Brongersma, L. D., 1931b. Eine neue Rasse von *Lygosoma kuekenthali* *Bttgr. Zool. Anz. Leipzig*, 96 (11/12): 335-336.
- Brongersma, L. D., 1933a . *Herpetological notes I-X. Zool. Meded., Leiden*, 16: 1-29.
- Brongersma, L. D., 1933b. A new gecko of the genus *Gymnodactylus* from New Guinea. *Ann. Mag. nat. Hist., London*, (10) 11: 252-253.
- Brongersma, L. D., 1933c. Ein neuer Laubfrosch aus Surinam. *Zool. Anz., Leipzig*, 103 (9/10): 267-270.
- Brongersma, L. D., 1934 *Contributions to Indo-Australian herpetology (Thesis Univ. Leiden)*. *Zool. Meded., Leiden*, 17: 161-251, pls 1-2.
- Brongersma, L. D., 1946. Some notes on species of the genera *Bachia* and *Scolecosaurus*. *Zool. Meded., Leiden*, 26: 237-246.
- Brongersma, L. D., 1948. A new subspecies of *Natrix mairii* (Gray) from Dutch New Guinea. *Proc. K. Akad. Wet., Amsterdam*, 51 (3): 372-381.
- Brown, W. C. & A. C. Alcala, 1980. Philippine lizards of the family Scincidae. *Suliman University Natural Science Monograph Series No. 2.* :1-264.
- Cochran, D.M., 1961. Type specimens of Reptiles and amphibians in the U.S. National Museum *Bulletin of the United States National Museum, Washington* (220): 1-291.
- Cogger, H. G., 1979. Type specimens of Reptiles and Amphibians in the Australian Museum. *Rec. Aust. Mus., Vol. 32* (4): 163-210.
- Crumly, C. R. 1990 Type catalogues of herpetological collections: An annotated list of lists. *Society for Study of Amphibians and Reptiles. Herpetological Circular No.* 18: 1-50.
- Daan S. & D. Hillenius, 1966. Catalogue of the type specimens of amhibians and reptiles in the Zoological Museum, Amsterdam. *Beaufortia*, 13: 117-144.
- Daan, S., 1967. Variation and taxonomy of the Hardun, *Agama stellio* (Linnaeus, 1758) (Reptilia, Agamidae). *Beaufortia*, 14 (172): 109-134. .
- Duellman W.E. & B. Berg, 196.2 Type specimens of Amphibians and Reptiles in the Museum of Natural History, The University of Kansas. *Univ. Kansas Publ.*

- Mus. Nat. Hist., vol. 15, No. 4, :185-204.
- Edeling, A. C. J., 1870. Recherches sur la fauna erpétologique de Sumatra. Nat. Tijdschr. Ned. Indië, 31: 376-388.
- Forcart, L., 1946. Katalog der Typusexemplare in der Amphibiensammlung des Naturhistorischen Museums zu Basel. Verh. naturf. Ges. Basel, Band 57: 118-142.
- Fry, D. B., 1912. Description of *Austrochaperina*, a new species of Engostomatidae from North Australia. Rec. Aust. Mus., Sydney, 9: 87-106, pls. 8-9.
- Garman, S., 1887. On the batrachians of Grand Cayman. Proc. Amer. Philos. Soc. 24: 273-285.
- Garman, S., 1888. On West-Indian reptiles and batracians in the Museum of Comparative Zoology at Cambridge, Mass. Bull. Essex Inst., 19: 1-53.
- Goldman, J., L. Hill & P.J. Stanbury, 1969. Type specimens in the Macleay Museum, University of Sydney, III. Amphibians and reptiles. Proc. Linnean Soc. New South Wales, Sydney, 93 (3): 427-438.
- Greer, A. E. & F. Parker, 1967. A new scincid lizard from the northern Solomon Islands. Breviora, 275: 1-20.
- Greer, A. E. & F. Parker, 1974. The *Fasciatus* species group of *Sphenomorphus* (Lacertilia: Scincidae); notes on eight previously described species and description of new species. Papua New Guin. Sci. Soc. Proc., 1973, 25: 31-61.
- Haas, C. P. J., 1950. Checklist of the snakes of the Indo-Australian Archipelago (Reptiles, Ophidia). Treubia 20 (3): 511-625.
- Hennig, W., 1936. Revision der Gattung *Draco* (Agamidae). Temminckia, 1: 153-220.
- Hoogmoed, M. S., 1978. An annotated review of the salamander types described in the Fauna Japonica. Zool. Meded., Leiden, 53 (9): 91-105.
- Hubrecht, A. A. W., 1879. Catalogue des collections formées et laissées par M. P. Bleeker, ... : 1-71 (De Breuk & Smits, Leiden).
- Inger, R. F. & H. Marx 1965 The systematics and evolution of the oriental colubrid snakes of the genus *Calamaria*. Fieldiana, Zool. Vol., 49: 1-304.
- Jong, J. K. de, 1926a. Fauna buruana: Reptiles. Treubia, 7 (2): 85-96.
- Jong, J. K. de, 1926b. *Heurnia ventromaculata* n. g. n. sp. und *Cantoria annulata* n. sp., zwei neue Schlangen von Neu-Guinea. Zool. Anz., Leipzig, 67: 302-304.
- Jong, J. K. de, 1927 Reptiles from Dutch New Guinea. Nova Guinea, 15 (Zool.) 3: 296-318.
- Jong, J. K. de, 1930. Notes on some reptiles from the Dutch East Indies. Treubia, 12 (1): 115-119.
- Kampen P. N. van, 1905. Amphibien von Palembang (Sumatra). Zool. Jahrb. (Syst.), Jena, 22 : 701-715, pl. 26.
- Kampen P. N. van, 1906. Amphibien in : Résultats de l'expédition scientifique Néerlandaise à la Nouvelle-Guinée en 1903, sous les auspices de Arthur Wichmann, chef de l'expédition. Nova Guinea, 5 (Zool.) 1: 163-180, pl. 6.
- Kampen P. N. van, 1907. Amphibien des Indischen Archipels. In : M. Weber, Zoologische Ergebnisse einer Reise in Niederländisch Ost-Indien, 4: 383-418, pl. 16. (E. J. Brill, Leiden).
- Kampen P. N. van, 1909a. Die Amphibienfauna von Neu-Guinea. Nova Guinea, 9 (Zool.) 1: 31-49, pl. II.
- Kampen P. N. van, 1909b. Eine neue Nectophryne-Art und andere Amphibien von Deli (Sumatra). Natuurk. Tijdschr. Ned. Ind. Weltevreden, 69: 18-24, 1 pl.
- Kampen P. N. van, 1912 Javanische Amphibien, gesammelt von Edw. Jacobson. Notes Leyden Mus., Leiden, 34: 75-79.
- Kampen P. N. van, 1913a. Amphibien von Waigeu und den Molukken. Bijdr. Dierk. 19: 89-92.
- Kampen P. N. van, 1913b. Amphibien, gesammelt von der Niederländischen Süd- Neu-Guinea Expedition von 1909-1910. Nova Guinea, 9 (Zool.) 3: 453-465, pl. 11.
- Kampen P. N. van, 1914. Zur Fauna von Nord Neu Guinea, Amphibien. Zool. Jahrb. (Syst.), Jena, 37: 365-378.
- Kampen P. N. van, 1915a. Amphibien von Nias. In: J.P. Kleiweg de Zwaan, Die Insel Nias bei Sumatra: 277-281 (Nijhoff, Den Haag) .
- Kampen P. N. van, 1915b. Amphibien von Neu Guinea. Nova Guinea, 13 (Zool.) 1: 39-41.
- Kampen P. N. van, 1919 Die Amphibienfauna von Neu-Guinea. Bijdr. Dierk., 21: 51-56.
- Kampen P. N. van, 1923. The Amphibia of the Indo-Australian Archipelago: 1-304, (E. J. Brill, Leiden).
- Kluge, A. G., 1983. Type-specimens of amphibians in the University of Michigan Museum of Zoology, Misc. Pubs. Mus. Zool. Univ. Mich. 166: 1-68.
- Kluge, A. G., 1984. Type-specimens of reptiles in the University of Michigan Museum of Zoology. Misc. Pubs. Mus. Zool. Univ. Mich. 167: 1-85.
- Kluge, A. G., 1993. Checklist, Gekkonoid Lizard Taxonomy. International Gecko Society San Diego, California.
- Kramer, E., 1978. Typenkatalog der schlangen im Naturhistorischen Museum Basel (BM) Stand 1977 Revue suisse Zool. 85 (3): 159-166.
- Kramer, E., 1979. Typenkatalog der Echsen im Naturhistorischen Museum Basel (BM) Stand 1978 Revue suisse Zool. 86 (1): 657-665.
- Lidth de Jeude, J. T. van, 1887. On a collection of reptiles and fishes from the West Indies. Notes Leyden Mus., 9: 129-139, pl. 2.
- Lidth de Jeude, J. T. van, 1890. Reptilia from the Malay Archipelago, II. Ophidia. In: M. Weber, Zoologische Ergebnisse einer Reise in Niederländisch Ost-Indien, 1: 178-192, pls. 15-16. (E. J. Brill, Leiden).
- Loveridge, A., 1932. New reptiles and amphibians from Tanganyika Territory and Kenya Colony. Bull. Mus. comp. Zool. Harv., 72 (10): 373-387.
- Loveridge, A., 1934. Australian reptiles in the museum of Comparative Zoology Bull. Mus. comp. Zool. Harv., 77 (6): 243-383.

- Loveridge, A., 1948. New Guinean Reptiles and Amphibians in the Museum of Comparative Zoology and United States National Museum. Bull. Mus. comp. Zool. Harv., 101 (2): 305-430.
- Lynch, J. D. & M. S. Hoogmoed, 1977. Two new species of *Eleutherodactylus* (amphibia: Leptodactylidae) from northeastern South America. Proc. Biol. Soc. Wash., 90 (2): 424-439.
- Mahnert, V., 1976. Catalogue des types de poissons, amphibiens et reptiles du Muséum d'Histoire naturelle de Genève. Revue Suisse de Zoologie, Genève, 83 (2): 471-496.
- Malnate, E. V., 1971. A catalog of the primary types in the herpetological collections of the Academy of Natural Sciences, Philadelphia (ANSP). Proc. Acad. nat. Sciences, Philadelphia, 123 (9): 345-375.
- Malnate, E. V. & G. Underwood, 1988. Australasian Natricine Snakes of the Genus *Tropidonophis*. (ANSP). Proc. Acad. nat. Sciences, Philadelphia, 140 (1): 59-201.
- Marx, H., 1958. Catalogue of reptiles and amphibians in Chicago Natural History Museum. Fieldiana Zool., 36 (4): 409-496.
- Marx, H., 1976. Supplementary catalogue of type specimens of reptiles and amphibians in Field Museum of Natural History. Fieldiana Zool., 69 (2): 33-94.
- McCoy, C. J. & N. D. Richmond, 1966. Herpetological type specimens in Carnegie Museum. Ann. Carneg. Mus., Pittsburgh, 38 (10): 233-264.
- Menzies, J. I., 1987. A taxonomic revision of Papuan *Rana* (Amphibia: Ranidae). Aust. J. Zool., 35: 373-418.
- Mertens R., 1922. Verzeichnis der typen in der herpetologischen Sammlung des Senckenbergischen Museums. Senckenbergiana, 4: 162-183.
- Mertens R., 1929. Bemerkungen über *Typhlops florensis*, seine Rassen und seine Verbreitung. Treubia, 11: 293-296.
- Mertens R., 1931. *Ablepharus boutonii* (Desjardin) und seine geographische Variation. Zool. Jahrb. (Syst.), 61 (1/2): 63-210.
- Mertens R., 1967. Die herpetologische Sektion des Natur-Museums und Forschungs-Institutes Senckenberg in Frankfurt a. M. nebst einen Verzeichnis ihrer Typen. Senckenbergiana, 48: 1-106.
- Mertens R., 1968. Die Arten und Unterarten der Schmuckbaumschlangen (*Chrysopelea*). Senckenbergiana, 49: 191-217.
- Musters, C. J. M., 1983. Taxonomy of the genus *Draco* L. Zool. Verh. Leiden, 199: 1-120, pls. 1-4.
- Noble, G. K., 1921a. Some new lizards from northwestern Peru. Ann. New York Acad. Sci., 29: 133-139.
- Noble, G. K., 1921b. Two new lizards from northwestern Peru. Ann. New York Acad. Sci., 29: 141-143.
- Noble, G. K., 1924. New lizards from northwestern Peru. Occ. Pap. Boston Soc. nat. Hist., 5: 107-113.
- Oudemans, J. T., 1894. Eidechsen und Schildkröten. In: R. Semon, Zoologische Forschungsreisen in Australien und dem malayischen Archipel, 5 Denkschr. Med. naturw. Ges. Jena, 8: 129-146.
- Parker, H. W., 1925. New and rare reptiles and batrachians from Madagascar. Ann. Mag. nat. Hist., London, (9) 16: 390-394.
- Parker, H. W., 1928. The *Brevicipitid* frogs of the genus *Microhyla*. Ann. Mag. nat. Hist. London, (10) 2: 473-499.
- Parker, H. W., 1931. Reports of an expedition to Brazil and Paraguay in 1926-1927, supported by the Trustees of the Percy Sladen Memorial Fund and the Executive Committee of the Carnegie Trust for Scotland, Amphibia and Reptilia. Proc. Linn. Soc. London, (Zool.) 37 (252): 285-289, pl. 16.
- Parker, H. W., 1934. A monograph of the frogs of the family *Microhylidae*: 1-28 (British Museum, London).
- Parker, H. W., 1935. Some lizards from Venezuela and the Dutch Leeward islands. Ann. Mag. nat. Hist., (10) 15: 480-484.
- Parker, H. W., 1940. The Australian frogs of the family *Leptodactylidae*. Novit. zool., London, 42 (1): 1-106.
- Peters, J. A., 1952. Catalogue of type specimens in the Herpetological Collections of the University of Michigan Museum of Zoology. Occ. Pap. Mus. Zool., Univ. Mich., 539: 1-55.
- Rooij. N. de, 1909. Reptilien (Eidechsen, Schildkröten and Krokodile). Nova Guinea, 5 (Zool.) (3): 375-383, pls. 14-18.
- Rooij. N. de, 1913. *Praeda itineris a L.F. de Beaufort in Archipelago indico facti ann. 1909-1910, III. Reptilien*. Bijdr. Dierk., 19: 15-29.
- Rooij. N. de, 1915. The reptiles of the Indo-Australian Archipelago, I. *Lacertilia, Chelonia, Emydosauria*: i-xiv, 1-384. (E. J. Brill, Leiden).
- Rooij. N. de, 1917. The reptiles of the Indo-Australian Archipelago, II. *Ophidia*: i-xiv, 1-334. (E. J. Brill, Leiden).
- Roux, J., 1907a. Beiträge zur Kenntnis der Fauna von Süd-Afrika. VII. *Lacertilia*. Zool. J. (syst.), Jena, 25: 403-444. pls. 14-15.
- Roux, J., 1907b. Beiträge zur Kenntnis der Fauna von Süd-Afrika. VIII. *Ophidia*. Zool. J. (syst.), Jena, 25: 732-741. pls. 27.
- Roux, J., 1910. Reptilien und Amphibien der Aru und Kei-Inseln. Abh. Senck. Ges. naturf. Fr., 33: 209-247, 2 pls.
- Saint Girons, H., 1953. Un vipère naine : *Vipera latastei montana*. Bull. Soc. zool. France, 78: 24-28.
- Schwartz A. & R. Thomas 1975. A check-list of West Indian amphibians and reptiles. Carnegie Museum of Natural History, Pittsburgh, special publication No.1: 1-216.
- Sket, B. & J. W. Arntzen, 1994. A black, non-trogomorphic amphibian from the karst of Slovenia: *Proteus anguinus parkelj* n. ssp. (Urodela: Proteidae) Bijdr. Dierk., 64 (1): 33-53.
- Slevin, J. R. & A. E. Leviton, 1956. Holotype specimens of

- reptiles and amphibians in the collection of the California Academy of Sciences. Proc. California Acad. Sciences, San Francisco, 4th ser., 28 (14): 529-560.
- Smith, M. A., 1920. On Sea snakes from the coast of the Malay Peninsula, Siam and Cochin-China. J. Fed. Malay St. Mus., X: 1-64.
- Smith H. M. & D. A. Langebartel & K. L. Williams, 1964. Herpetological type-specimens in the University of Illinois Museum of Natural History. Illinois biol., Monogr., 32: 1-80.
- Stull, O. G., 1932. Five new subspecies of the family Boidae. Occ. Pap. Boston Soc. Nat. Hist., 8: 25-30, pls. 1-2.
- Taylor, E. H., 1918. Reptiles of Sulu Archipelago. Phil. J. Sci., Manila, (D) 13 (5): 233-267, pls. 1-3.
- Taylor, E. H., 1920. Philippine Amphibia. Phil. J. Sci., Manila, 16 (3): 213-359, 10 pls.
- Taylor, E. H., 1922a. The snakes of the Philippine islands. Mon. Bur. Sci., Manila, publ. No. 16: 1-312, pls. 1-37.
- Taylor, E. H., 1922b. Additions to the herpetological fauna of the Philippine islands. I. Phil. J. Sci., Manila, 21 (2): 161-206, pls. 1-7.
- Taylor, E. H., 1922c. Additions to the herpetological fauna of the Philippine islands. II. Phil. J. Sci., Manila, 21 (3): 257-303, pls. 1-4.
- Taylor, E. H., 1923. Additions to the herpetological fauna of the Philippine islands. III. Phil. J. Sci., Manila, 22 (5): 515-557, pls. 1-3.
- Taylor, E. H., 1925. Additions to the herpetological fauna of the Philippine islands. IV. Phil. J. Sci., Manila, 26 (1): 97-111.
- Taylor, E. H., 1944. Present location of certain herpetological and other type specimens. The Univ. of Kansas Science Bulletin Vol., 30 (11): 117-187.
- Taylor, E. H., 1965. New Asiatic and African Caecilians with redescription of certain other species. The Univ. of Kansas Science Bulletin Vol., 46 (6): 253-302.
- Taylor, E. H., 1966. Herpetology of the Philippine Islands I (reprint).
- Taylor, E. H., 1966. Herpetology of the Philippine Islands II (reprint).
- Tiedemann F. & M. Häupl 1980. Typenkatalog der Herpetologischen Sammlung Teil II : Reptilia. Kataloge der wissenschaftlichen Sammlungen des Naturhistorischen Museum in Wien, Vertebrata, Heft 2: 1-79.
- Tyler, M. J., 1964. An Investigation of the systematic position and synonymy of *Hyla montana* Peters & Doria, (Anura, Hylidae). Zool. Abh. Dresden Bd. 27 (10): 265-270.
- Tyler, M. J., 1965. Taxonomic studies of some hylid frogs of Australia and New Guinea. Proc. Zool. Soc. London Vol., 145 (1): 91-106.
- Tyler, M. J., 1968. Papuan hylid frogs of the genus *Hyla*. Zool. Verh., Leiden, 96: 1-223.
- Wagenaar Hummelinck, P., 1940. Studies on the fauna of Curacao, Aruba, Bonaire and the Venezuelan islands: 1-130, pls. 1-16 (Thesis Univ. Utrecht).
- Weber, M., 1890. Reptilia from the Malay Archipelago, I. In: M. Weber, Zoologische Ergebnisse einer Reise in Niederländisch Ost-Indien, Leiden, I: 159-177, pls. 14.

L. van Tuijl
Instituut for Systematics
and Population Biology
(Zoölogisch Museum)
University of Amsterdam
P.O. Box 94766
1090 GT Amsterdam
The Netherlands

Received: April 25, 1995.
Distributed: 8 September 1995