

NOTE XII.

PELARGODERUS SIJTHOFFII, N. SP.

BY

C. RITSEMA Cz.

Agreeing with *P. semitigrinus* Rits. (Notes Leyd. Mus. Vol. VII, p. 43, pl. 3, fig. 3) but easily recognizable by two very obvious differences.

In the new species the elytra are very distinctly speckled with white, about as in *P. alcanor* Newm.; in *semitigrinus*, however, they are indistinctly speckled with greyish.

In the new species a narrow dirty white stripe is present along the middle of the apical fourth of each elytron, extending down to the pointed outer angle; in *semitigrinus* this apical stripe is entirely wanting.

Two males and one female from Padang Sidempoean (West-Sumatra) belonging to Mr. P. F. Sijthoff's collection. One of these males is now in the collection of Dr. H. J. Veth, the female in that of the Leyden Museum.

Obs. *Pelargoderus nigromaculatus*, described by me in December 1897 and published in Notes Leyd. Mus. Vol. XX, p. 34, has been concisely described in the same year by Dr. L. von Heyden under the name of *Pelargoderus trigonalis* in Abh. Senckenb. Ges. Bd. XXIII, p. 573.

Pelargoderus (Rhamses) vitticollis Thoms. (Ann. Soc. Ent. France, 1878, Bull. p. 18) of which I have examined the type in Mr. R. Oberthür's collection = *Epepeotes polorator* Newm. and consequently not from Borneo but from the Philippine islands.

Leyden Museum, 30 November 1901.