Serpocaulon × *manizalense*: a new hybrid between simple- and pinnate-leaved species of *Serpocaulon* (*Polypodiaceae*) from Colombia

D. Sanín¹, V. Torrez²

Key words

endangered fern
hybridization
multivariate analyses
Serpocaulon adnatum
Serpocaulon levigatum

Abstract During a revision of *Serpocaulon* from Colombia, a new hybrid was found between *S. adnatum* and *S. levigatum* near to Manizales city, which is described and illustrated herein. Qualitative and quantitative spore and macro-morphological characters were evaluated using principal component analyses to distinguish the new taxon. Our results suggest that the perispore with leasura, lamina width, rhizome diameter, blade dissection and number of pinnae are important characters to distinguish *S. × manizalense*. This is the first record of a hybrid between a simple and pinnate-leaved species in *Serpocaulon*, which is considered to be Critically Endangered (CR).

Published on 30 October 2014

INTRODUCTION

Serpocaulon A.R.Sm. is a Neotropical genus of *Polypodiaceae* with 42 species and three known hybrids (Smith et al. 2006, Labiak & Prado 2008, Rojas-Alvarado & Chaves-Fallas 2013, Schwartzburd & Smith 2013, Sanín 2014). The genus is characterized by a suite of morphological characters that includes long-creeping rhizomes, clathrate scales, veins regularly anastomosing (goniophleboid), and areoles chevron-shaped with a single, free, included veinlet (Smith et al. 2006).

The two known hybrid taxa are *Serpocaulon* × *pubescens* (Rosenst.) Schwartzb. & A.R.Sm. (Schwartzburd & Smith 2013), *S. × sessilipinnum* A.Rojas & J.M.Chaves (Rojas-Alvarado & Chaves-Fallas 2013), and one other taxon is often suggested as being of hybrid origin: *S. semipinnatifidum* (Fée) A.R.Sm. (Tryon & Stolze 1993, Moran 1995). However, there is the possibility to find more hybrids mainly between *S. levigatum* and others Andean species (Tryon & Stolze 1993, Moran 1995, Sanín 2011).

During a review of *Serpocaulon* from Colombia, a new hybrid was found co-occurring with *S. adnatum* (Kunze ex Klotzsch) A.R.Sm., and *S. levigatum* (Cav.) A.R.Sm. Considering that hybrids occur in areas that were subject to extreme changes where the parents used to be or are present (Rieseberg 1997), and that they have intermediate characters to their parental taxa (Moran & Watkins 2004), we presume that *S. adnatum* and *S. levigatum* are the putative parental taxa of the new hybrid, which also slightly resembles *S. semipinnatifidum*, of which no records are known from the area. Our study aims to describe the new hybrid and distinguish it from its putative parents and *S. semipinnatifidum*.

MATERIALS AND METHODS

Morphological sampling and characters

To discriminate the new hybrid, 411 herbarium specimens of *Serpocaulon adnatum* (133), *S. levigatum* (222), *S. semipinnatifidum* (54), and the new hybrid (2) were selected from the herbaria CAUP, CHOCO, COL, CUVC, FAUC, FMB, HUA, HUQ, JAUM, MEDEL, MO, NY, PSO, and TOLI. Each studied specimen was sampled from a different individual.

Serpocaulon adnatum and *S. levigatum* were included in the analyses as they were found co-existing with the new hybrid. *Serpocaulon semipinnatifidum* was not found to co-exist with them, but it was included in the analyses as it can be confused with the new hybrid. The putative parents suggested for *S. semipinnatifidum*, in combination with *S. levigatum*, *S. funckii* (Mett.) A.R.Sm. (Moran 1995) and *S. lasiopus* (Klotzsch) A.R.Sm. (Tryon & Stolze 1993), were excluded from the analysis as they were not found to co-exist with the new hybrid. Furthermore, they are not recorded in the Caldas department (Fraume et al. 1990, Sanín & Duque-Castrillón 2006, Sanín et al. 2006, 2008, Álvarez-Mejía et al. 2007, Sanín 2011).

For morphometric analyses, 38 specimens were used as operational taxonomic units (OTU), representing the entire geographical range and the morphological variability within each taxon. A total of 59 morphological characters were measured, from which 35 were qualitative and 24 were quantitative. From these quantitative characters, eight were spore characters and 16 were macro-morphological characters.

The analysed spore characters are equatorial axis, length of aperture, polar axis, endospore, exospore, central verrucae, verrucae height and width. The spore description follows Ramírez-Valencia et al. (2013). The macro-morphological characters are rhizome diameter, rhizome scale length and width, phyllopodia distance, petiole length, lamina length and width, number of pinnae, lamina scale length and width. Additionally, we measured for the medial pinna length and width, the number of areolae and sori between the principal vein and the costa, and along the principal vein and the costa. For *S. levigatum*, which has a simple lamina, these characters were measured

¹ Botanic Garden Meise, Bouchout Domain, Nieuwelaan 38, 1860 Meise, Belgium; corresponding author e-mail: dav.sanin@gmail.com.

² Plant Conservation and Population Biology, KU Leuven University, Kasteelpark Arenberg 31, B-3001, Heverlee, Belgium; torflorvania@gmail.com.

for the entire lamina. The macro-morphology description follows Lellinger (2002).

Numerical and statistical analyses

Quantitative characters were analysed by range and median values. Principal component analyses (PCA) were based on quantitative values of spore and macro-morphological characters. PCA results were used to support distinctions among the taxa. These analyses were performed using R (R Development Core Team 2013).

Conservation status

The conservation status of the new hybrid was assessed by applying the IUCN Red List Categories and Criteria (IUCN 2001).

RESULTS

Principal component analyses

Spore characters (Table 1)

The first three components accounted for 80 % of the total variance. The first principal component had high contributing loading values from verrucae height, verrucae width, central verruca, and polar axis. The second component had high contributing loadings from the endospore, exospore, central verrucae, and verrucae height. Finally, the third component

had high contributing loadings from polar axis, equatorial axis, exospore and endospore.

In the scatterplot against the first two components (Fig. 1), the OTUs are arranged in loose and slightly overlapping groups, corresponding to *S. levigatum*, *S. semipinnatifidum* and *S. × manizalense*. Despite the overlap, it is possible to recognize the new hybrid (*S. × manizalense*) between *S. levigatum* and *S. adnatum*. In addition, the spores of the other hybrid (*S. semipinnatifidum*) are in a cluster separate from the other taxa.

Macro-morphological characters (Table 2)

The first three components accounted for 68.7 % of the total variance observed. The first principal component had high contributing loading values from petiole length, lamina width, rhizome diameter and number of pinnae, among the four most important. The second component had high contributing loadings from medial pinna width and lamina length. Finally, the third component had high contributing loadings from rhizome scale length, rhizome scale width, rhizome diameter, and lamina length.

In the scatterplot of the first two components, the OTUs are arranged in four different clusters, each representing a different taxon (Fig. 2). This result allowed us to easily characterize the four taxa and support the description of the new hybrid (*S. × manizalense*). Nevertheless, *S. levigatum* and *S. semipinnatifidum* are slightly overlapping.

Table 1 Summary of the principal component weights for the spore characters. In **bold**, morphological characters showing the highest values.

Character	Axes 1	Axes 2	Axes 3
Equatorial axis	0.607	-0.538	0.418
Polar axis	0.715	-0.285	0.515
Endospore	0.362	0.791	0.110
Exospore	0.365	0.717	0.191
Central verrucae	0.906	0.103	-0.184
Verrucae width	0.910	-0.021	-0.154
Verrucae height	0.914	0.072	-0.109
Aperture longitude	0.547	-0.289	-0.594

Table 2 Summary of the principal component weights for the macro-morphological characters. In **bold**, morphological characters showing the highest values.

Character	Axes 1	Axes 2	Axes 3
Rhizome diameter	0.899	-0.229	0.135
Rhizome scale length	0.766	-0.100	0.460
Rhizome scale width	0.706	-0.072	0.404
Phyllopodia distance	0.142	-0.029	0.012
Petiole length	0.925	-0.163	0.089
Number of pinnae	0.895	-0.265	0.023
Lamina length	0.856	0.268	0.122
Lamina width	0.917	-0.271	0.024
Pinna media length	0.803	0.473	-0.027
Pinna media width	0.752	0.374	-0.067

Fig. 1 Plot of the first two components of the PCA from the spore morphology. *Serpocaulon × manizalense* (●); *S. semipinnatifidum* (○); *S. adnatum* (×); *S. levigatum* (Δ).

Fig. 2 Plot of the first two components of the PCA from the macro-morphology. *Serpocaulon × manizalense* (●); *S. semipinnatifidum* (○); *S. adnatum* (×); *S. levigatum* (Δ).

DISCUSSION

Spore characteristics of the hybrid

The slight overlap of *S. levigatum*, *S. semipinnatifidum* and *S.* × *manizalense* in the PCA of spore morphology, could be explained by the fact that *S. levigatum* is a parent of *S. semipinnatifidum* (Tryon & Stolze 1993, Moran 1995, Sanín 2011) and with the present results we propose that it is also one of the parental species of the new hybrid. In addition, the new hybrid is arranged between *S. levigatum* and *S. adnatum* (Fig. 1), both proposed as putative parental species of *S.* × *manizalense*.

According to the PCA results, the most important spore characters to differentiate the new hybrid are the shape and orna-

mentation of the spores, in particular the verrucae (Table 1). Similarly, Ramírez-Valencia et al. (2013) reported these spore characters as useful to distinguish 21 species of *Serpocaulon* from Colombia.

Interestingly, the described hybrid has well-formed spores (see description of the hybrid), an unusual characteristic in fern hybrids. Usually, fern hybrids have malformed spores (Haufler 2008, Sharpe et al. 2010). However, *S.* × *manizalense* is not the first described hybrid with well-formed spores. Such a characteristic has been described in other fern hybrid taxa such as *Polystichum* Roth. (Mullenniex et al. 1998), *Asplenium* L. (Morzenti 1967), *Polypodium* L. (Haufler et al. 1995), and also in *Serpocaulon* (Rojas-Alvarado & Chaves-Fallas 2013).

Fig. 3 *Serpocaulon* × *manizalense* D.Sanín & Torrez. a. Habit; b. segment of the blade; c. detail of the veins; d. rhizome scale; e. lamina scale (all: Sanín et al. 2646, FAUC, HUA, COL). — Scale bars: a = 5 cm; b, c = 1 cm; d = 1 mm; e = 0.7 mm. — Drawn by: a–c. L.F. Coca; d, e. D. Sanín.

Macro-morphology of the hybrid

Some of the most frequently used characters to distinguish *Serpocaulon* species are: rhizome diameter, laminar length and width, petiole length, rhizome scale length and width (Lellinger 1989, Tryon & Stolze 1993, Moran 1995, Labiak & Prado 2008, Sanín 2014). However, in the present study other less frequently used characters such as number of pinnae or segments, medial pinnae length and width appeared to be important to distinguish the new hybrid.

The PCA analysis of the macro-morphology resulted in better-resolved groups than that of the spore characters, and only *S. levigatum* and *S. semipinnatifidum* are overlapping. This is not surprising, since it was reported that *S. levigatum* is the parental taxon of *S. semipinnatifidum* (Tryon & Stolze 1993, Moran 1995, Sanín 2011), while the other parental taxon remains uncertain. Moran (1995) proposed that the other putative parent may be *S. funckii* for the north of the Andes hybrids (Colombia and Venezuela). However, Tryon & Stolze (1993) suggested that *S. lasiopus* may be the other putative parental taxon of *S. semipinnatifidum* for the hybrids located in Peru.

Hybrid habitat

Despite the extensive botanical exploration of the Chinchiná basin river (Fraume et al. 1990, Orrego et al. 2004, Sanín & Duque-Castrillón 2006, Sanín et al. 2006, 2008, Álvarez-Mejía et al. 2007), there is only one wild population of *S. × manizalense* known, and another single plant propagated by rhizome, which has been maintained for conservational purposes in the JBOUC.

Frequently, the formation of a hybrid is triggered by an extreme change in the habitat of the parental taxa (Rieseberg 1997, Rieseberg et al. 2006, Kentner & Mesler 2000).

Thus, most of the hybrids appear in highly disturbed areas such as road cuts (Barrington 1985). Probably, the generation of *S. × manizalense* was promoted by the conversion of the natural area to a landfill, which has been opened since 1991.

TAXONOMIC TREATMENT

Serpocaulon × manizalense D.Sanín & Torrez, *hybrid nov.*
— Fig. 3, 4

Hybrida inculta e *Serpocaulon adnatum* et *S. levigatum* genita, epiphytica et terrestrium; a *S. semipinnatifidum* differ rhizoma longe 5–5.5 mm (versus 1.7(–2.9–)4.7 mm), lamina longe 25(–30–)33 per 8(–28–)30 cm (versus 15(–24.7–)32 per 1.2(–4–)9) cm, areolis magis numerosis secus pinna basal 1–5 series (versus 1–3 series). — Type: Sanín et al. 2646 (holo FAUC; iso COL, HUA), Colombia, Caldas, Manizales, flanco occidental de la Cordillera

Central, vía a Neira, cerca del Relleno Sanitario La Esmeralda, zona a la derecha que se pronostica conservar en la ampliación del Relleno [N05° 04'49" W75°30'27"], 2200 m, 3 Nov. 2008.

Etymology. The new hybrid is named after the city of Manizales and its people.

Plants terrestrial and epiphytic. *Rhizomes* 5–5.5 mm diam, long-creeping, dark brown to reddish, faintly farinose. *Phyllopodia* 7–7.3 cm apart. *Scales* 1.3(–2–)2.5 by 0.8(–1–)2.0 mm, appressed, with rounded bases and rounded to acute apices, ovate to ovate-lanceolate, dispersed along the rhizome, dark brown at centre, clathrate, hyaline at the margin. *Petioles* 10–12.2 cm, articulate, subterete, 1/2 length of blade, stramineous. *Laminae* 25(–30–)33 by 8(–28–)30 cm, simple with crenate margins to base and lanceolate shape or basally pinnate and ovate-lanceolate truncate shape, glabrous, rachis distally stramineous. *Pinnae* absent or up to 4 pairs, of irregular size, firm to coriaceous, proximally adnate, with 7 basal lobules that are attenuate, apex long-caudate to acute in the distal pinnae. Proximal pinnae 15.8–16.3 by 3.5–3.8 cm. *Veins* goniophleboid, closed chevron-shape areoles 28–30 by 2–5 rows between the costae and pinna margins, with one included fertile veinlet or open, irregularly shaped with two included fertile veinlets. Laminar indument consisting of scales, 0.8–1 by 0.5–0.6 mm, acicular, bicolorous, to 9 cells wide. Sori in 1–5 rows between costae and margin. Spores 54.5–56.5 by 33.5–35.5 µm, well-formed, monolete, aperture 21.8–24.8 µm long, laesurae straight, ambit ellipsoid, plane-convex to concave-convex, ornamentation densely verrucate, verrucae 3–4 by 4.5–5.5 µm, regular in size and rounded, evenly distributed, exospore 1–1.5 µm, perispore 1.3–1.5 µm, sulcate.

Distribution & Ecology — *Serpocaulon × manizalense* occurs on the western slope of the Central Andean Cordillera of Colombia, near Manizales on the road to Neira, in the locality of La Esmeralda landfill (Fig. 5a, Map 1). It was found epiphytic on *Rhus striata* Ruiz & Pav. (*Anacardiaceae*) and terrestrial in secondary forests (Fig. 5b). In addition, the only *Serpocaulon* species found in the area were *S. adnatum* and *S. levigatum*. *Serpocaulon × manizalense* was collected fertile in November 2008, and also it was seen fertile in September 2013 from cultivated plants in the Caldas University Botanical Garden (JBOUC) (Fig. 5c–e).

Notes on related taxa — The characters of *S. × manizalense* that are crucial to distinguish it from *S. semipinnatifidum* are present in the rhizome, lamina, and spores (Table 3). The rhizome in *S. × manizalense* is wider than in *S. semipinnatifidum*. The blade dissection in *S. × manizalense* is lobate to proximally pinnate (Fig. 3, 6), whereas *S. semipinnatifidum* is lobate to proximally pinnatisect (Fig. 6). Frequently, the lamina length of

Fig. 4 Scanning electron micrographs of spores of *S. × manizalense* D.Sanín & Torrez, 1200×, 15 kV. a. Lateral view; b. clustered spores with visible laesura in the perispore (all: Sanín et al. 2646, FAUC). — Scale bars = 10 µm.

Fig. 5 Ecosystem, location, and propagated plant of the type collection. a. Ecosystem of *S.* × *manizalense* D.Sanín & Torrez; b. location of the type collection; c–e. asexual cultivated plants of *S.* × *manizalense*.

Table 3 Contrasting characters between *S. × manizalense*, *S. semipinnatifidum*, *S. adnatum*, and *S. levigatum*. Minimum, median, and maximum values are given for quantitative characters.

Character	<i>S. × manizalense</i>	<i>S. semipinnatifidum</i>	<i>S. adnatum</i>	<i>S. levigatum</i>
Rhizome diameter (mm)	5–5.5	1.7(–2.9–)4.7	4(–7–)10	1.6(–3.3–)4.7
Scale length (mm)	1.3(–2–)2.5	0.8(–1–)1.8	1(–2–)3	1(–1.2–)1.7
Blade dissection	Lobated to proximally pinnate	Lobated to proximally pinnatisect	Pinnate	Simple
Lamina shape	Lanceolate (when it is simple) or ovate-lanceolate (when it is pinnate)	Lanceolate	Narrow to amply ovate	Ovate-lanceolate
Lamina length (cm)	25(–30–)33	15(–24.7–)32.5	20(–52–)88	5.7(–14–)26
Number of areoles between the costae and pinna margin	1–5	1–3	4(–6–)7	3(–4–)6
Perine with laesura	Present	Absent	Absent	Absent

Fig. 6 Silhouettes. a. *Serpocaulon × manizalense* (Sanín et al. 2646, FAUC, HUA, COL); b. *S. adnatum* (Coca et al. 199, FAUC); c. *S. levigatum* (Sanín et al. 5124, NY); d. *S. semipinnatifidum* (Rodríguez et al. 6103, HUA, NY). All plants collected in Colombia. — Scale bar = 10 cm.

Map 1 Distribution of *Serpocaulon × manizalense* D.Sanín & Torrez.

S. × manizalense is larger than of *S. semipinnatifidum* (Fig. 6). In addition, the species differ in the number of areoles between the costae and pinna margin, *S. × manizalense* has 1–5 areoles and *S. semipinnatifidum* has 1–3 areoles. The spores of *S. × manizalense* show a laesura in the perispore (Fig. 4), while the spores of *S. semipinnatifidum* do not have a laesura in the perispore (Ramírez-Valencia et al. 2013).

The most conspicuous intermediate characters between the putative parents and the new hybrid are the blade dissection and the number of pinnae in the lamina: pinnate with 4(–7–)10 pairs in *S. adnatum*, lobate to pinnate with up to 4 pinnae pairs in the hybrid, and a simple lamina in *S. levigatum* (Fig. 6). The rhizome diameter of *S. × manizalense* is intermediate between *S. adnatum* and *S. levigatum*. Less conspicuous, but also intermediate were the colour and rhizome scales length. The rhizome scales in *S. adnatum* are dark brown to nearly black with 1–3 mm long (Fig. 7), in *S. levigatum* are pale orange with 1–1.7 mm long (Fig. 7), whereas in *S. × manizalense* are pale orange to brown with 1.3–2.5 mm long (Fig. 7).

Conservation — IUCN Red List Category: Critically Endangered [CR B2a + D]. The EOO cannot be estimated for *S. × manizalense* because it is known from only one location. The AOO is 9 km², and it falls completely outside any protected and pristine area under the Colombian System of Protected Areas.

Acknowledgements The first author thanks L.M. Álvarez, A. Pardo, and F. Cardona for their support, as well as the staff of the Caldas University, especially the research student group (SByRFG). We thank colleagues at BR, CAUP, CHOCO, COL, CUVC, FAUC, FMB, HUA, HUQ, JAUM, LPB, MBM, MEDEL, MO, NY, PSO, QCNE, TOLI, UFP, and UPCB for allowing access to specimens. Thanks to L.F. Coca for the drawing of the habit of the plant, V. Ramírez-Valencia for her help in the spore analyses, M. López

Fig. 7 Rhizome scales. a. *Serpocaulon adnatum*; b. *S. × manizalense*; c. *S. levigatum* (a. Sanín et al. 3080, FAUC; b. Sanín et al. 2646, FAUC, HUA, COL; c. Sanín et al. 3315, FAUC). — Scale bars: a = 2.5 mm; b = 1 mm; c = 1.5 mm.

for the SEM photographs, B. Bassuner for the IUCN threat categorization, and A. Pérez for digital design assistance. Special thanks to M. Sundue, P. Hovenkamp, and an anonymous reviewer for their kind comments that improved substantially the manuscript. This research was supported by the Caldas University (FAUC Herbarium and IIES Institute) and the Colombian Association of Herbaria (ACH).

REFERENCES

- Álvarez-Mejía LM, Sanín D, Álzate-Q NF, et al. 2007. Plantas de la región Centro-Sur de Caldas. Cuadernos de Investigación No. 28. Editorial Universidad de Caldas, Manizales.
- Barrington DS. 1985. Ecological and historical factors in fern biogeography. *Journal of Biogeography* 20: 275–279.
- Fraume M, Álvarez-Mejía LM, Gallego JH. 1990. Monteleón, relicto de selva andina de Manizales, Colombia. *Revista Agronomía* 4: 24–35.
- Haufler CH. 2008. Species and speciation. In: Ranker TA, Haufler CH (eds), *Biology and evolution of ferns and lycophytes*: 303–331. Cambridge University Press, New York.
- Haufler CH, Windham MD, Rabe EW. 1995. Reticulate evolution in the *Polypodium vulgare* complex. *Systematic Botany* 20: 89–109.
- IUCN. 2001. IUCN Red List Categories and Criteria: Version 3.1. Gland, Switzerland and Cambridge, UK. Available from: <http://www.iucnredlist.org/technical-documents/categories-and-criteria/2001-categories-criteria> [accessed 24 Nov. 2013].
- Kentner EK, Mesler MR. 2000. Evidence for natural selection in a fern hybrid zone. *American Journal of Botany* 87: 1168–1174.
- Labiak P, Prado J. 2008. New combinations in *Serpocaulon* and a provisional key for the Atlantic rain forest species. *American Fern Journal* 98: 139–159.
- Lellinger DB. 1989. The ferns and ferns-allied of Costa Rica, Panamá, and the Chocó (Part 1: Psilotaceae through Dicksoniaceae). *Pteridologia* 2A.
- Lellinger DB. 2002. A modern multilingual glossary for taxonomic pteridology. *Pteridologia* 3. American Fern Society.
- Moran RC. 1995. *Polypodium* L. In: Moran RC, Riba R (eds), *Psilotaceae a Salviniaceae*. Flora Mesoamericana. Vol. 1: 349–365. Mexico City, Universidad Nacional Autónoma de México, Saint Louis, Missouri Botanical Garden & London, The Natural History Museum. Distrito Federal, México D.F.
- Moran RC, Watkins JE. 2004. *Lomariopsis × farrarii*: a new hybrid fern between *L. japurensis* and *L. vestita* (Lomariopsidaceae) from Costa Rica. *Brittonia* 56: 205–209.
- Morzenzi VM. 1967. *Asplenium plenum*: A fern which suggest an unusual method of species formation. *American Journal of Botany* 54, 9: 1061–1068.
- Mullenniex A, Hardig TM, Mesler MR. 1998. Molecular confirmation of hybrid swarms in the fern genus *Polystichum* (Dryopteridaceae). *Systematic Botany* 23: 421–426.
- Orrego O, Botero JE, Verhelst JC, et al. 2004. Plantas vasculares del municipio de Manizales, Caldas. *Boletín Científico Museo Historia Natural Universidad de Caldas* 8: 61–106.
- R Development Core Team. 2013. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. Available at <http://www.R-project.org>. Last accessed 24 December 2013.
- Ramírez-Valencia V, Sanín D, Pardo-Trujillo A. 2013. Análisis morfológico de las esporas de *Serpocaulon* (Polypodiaceae) de la Cordillera Central de Colombia. *Caldasia* 35: 177–197.
- Rieseberg LH. 1997. Hybrid origin of plant species. *Annual Review of Ecology and Systematics* 28: 359–389.
- Rieseberg LH, Wood TE, Baack E. 2006. The nature of plant species. *Nature* 440: 524–527.
- Rojas-Alvarado AF, Chaves-Fallas JM. 2013. A new hybrid of *Serpocaulon* (Polypodiaceae) from Costa Rica. *American Fern Journal* 103: 175–181.
- Sanín D. 2011. Revisión taxonómica de *Serpocaulon* (Polypodiaceae), para la Cordillera Central y el Macizo Colombiano. MSc Thesis. Vegetal Biology, Caldas University, Colombia.
- Sanín D. 2014. *Serpocaulon obscurinervium* (Polypodiaceae), a new fern species from Colombia and Ecuador. *Plant Ecology and Evolution* 147: 154–161.
- Sanín D, Álvarez-Mejía LM, Mancera-Santa JC, et al. 2008. Monilofitos y Licofitos de la Cuenca del Río Chinchiná (Caldas, Colombia). Clave para géneros y catálogo de las especies. *Revista Academia Colombiana de Ciencias* 32: 331–352.
- Sanín D, Duque-Castrillón CA. 2006. Estructura y composición florística de dos transectos localizados en la Reserva Forestal Protectora de Río Blanco (Manizales, Caldas, Colombia). *Boletín Científico Museo Historia Natural Universidad de Caldas* 10: 19–44.
- Sanín D, Mancera-Santa JC, Castaño-R N, et al. 2006. Catálogo preliminar de las plantas vasculares de la Reserva Forestal Protectora “Río Blanco” (Manizales, Caldas, Colombia). *Boletín Científico Museo Historia Natural Universidad de Caldas* 10: 44–60.
- Schwartsburd P, Smith AR. 2013. Novelties in *Serpocaulon* (Polypodiaceae). *Journal of the Botanical Research Institute of Texas* 7: 85–93.
- Sharpe JM, Mehltreter K, Walker LR. 2010. Ecological importance of ferns. In: Mehltreter K, Walker LR, Sharpe JM, *Fern Ecology*: 1–21. Cambridge University Press, New York.
- Smith AR, Kreier HP, Haufler CH, et al. 2006. *Serpocaulon*, a new genus segregated from *Polypodium*. *Taxon* 55: 919–930.
- Tryon RM, Stolze RG. 1993. *Pteridophyta of Perú*. Part V. 18. *Aspleniaceae* – 21. *Polypodiaceae*. *Feldiana Botany n.s.* 32: 1–190.

INDEX OF COLLECTIONS

The type collection is in **bold**. ad = *S. adnatum*; le = *S. levigatum*; ma = *S. × manizalense*; se = *S. semipinnatifidum*

Acevedo et al. 1405: le – Acosta-Arteaga et al. 104: le; 449: le – Agudelo et al. 302: ad; 307: ad; 1586: le; 3289: le; 3460: le – Aguilar 339: ad; 391: ad; 723: le – Albert de Escobar & Brand 2098: le; 2118: le – Albert de Escobar & Giraldo 5172a: le; 6712: le – Albert de Escobar et al. 3695: se; 3957: se; 3977: ad; 4179: se; 4517: le; 4531: le; 5241: le; 5870: le; 6005: le; 6982: le; 8809: se – Álvarez-Mejía 1336: ad – Álvarez-Mejía & Ospina 2171:

le – Alvear & Sánchez 628: le – Álzate & Cardona 492: le – Arbeláez et al. 54: le; 108: se; 111: le; 112: se; 115: le; 245: le; 272: ad; 293: le; 321: le; 513: le – Arbeláez-G. et al. 93: le; 352: ad; 1057: le; 1256: le; 1855: le; 2081: le – Atehortúa 117: se; 337: le; 351: le; 403: ad; 406: ad; 407: ad. Barclay 4673: le – Becking 1542: le – Benavides & Jiménez 4078: le – Betancur et al. 3172: ad – Bischler 1896: le; 1943: le; 2246: ad – Bittner & Guzmán 2700: le – Bolaños et al. 7: le – Bristol 404: le; 427: le.

- Cabrera et al. 5081: le – Calle & Muriel 99: le – Callejas & Echeverri 11465: se; 11512: le; 11513: se; 11567: le – Callejas & Escobar 7408: se – Callejas & Roldán 10493: le; 10506: le; 10671: le – Callejas et al. 368: ad; 823: le; 3741: le; 3792: ad; 6375: le – Cardona 544: le – Cardona-Naranjo 565: ad – Casañas-Suarez 563: le – Castaño-Rubiano 581: ad – Ceballos & Lasso 223: le – Churchill & Rengifo 17317: ad – Coca & Duque-Buitrago 66: le – Coca et al. 199: ad – Correa 461: le – Correa & Yarce 1666: se – Croat 56406: le; 71389: ad – Croat & Galeano 70776: le – Cuatrecasas 8617: ad; 9335: le; 9607: ad; 11654: le.
- David & Giraldo 727: ad – David & Idárraga 484: se; 486: le – De Benavides 2101: le; 2879: ad; 9186: ad; 9612: ad; 9702: ad – De Fraume et al. 598: ad – Devia 742: ad – Díaz S.L. & Alcázar 51: le – Díaz-Ibarra 282: le – Díaz-P. 3145: le; 3708: le – Díaz-P. et al. 517: le; 906: le – Duque-Jaramillo 3339: le – Duque-Quintero 14: ad; 18: ad.
- Echeverri 426: se – Echeverri et al. 457: le – Escobar & Zuluaga 76: le – Espinal & Ramos 3580: ad.
- Feuillet et al. 79: ad – Fonnegra et al. 2231: le; 2233: le; 2234: se; 5274: le; 5287: ad; 5317: ad; 8211: le – Forero 2256: le – Foster & Foster 1943: le – Franco-Roselli et al. 1654: ad; 1729: ad.
- Galindo et al. 450: le – García-Barriga 17344: ad; 20472: le – Gil et al. 59: le – Gómez et al. 612: ad – González 1027: le – Grant 9932: le; 10658: le; 10833: se – Grubb & Guymier 91: ad – Guarín 345: ad – Gutiérrez & Barkley 17c004: ad; 18c277: se – Gutiérrez-V. 14: se – Guzmán 5: ad.
- Hagemann & Leist 1207: le; 1503: le; 1567: ad; 1872: le – Haught 4097: le – Henao 21: le – Hernández 307: le; 362: le – Hernández et al. 532: le – Hernández-Schmidt 1309: se – Hno. Daniel 42: se; 678: ad – Hodge 6648: se – Hoyos et al. 29: le – Huertas & Camargo 5053: le.
- Idrobo & Schultes 1135: le – Idrobo et al. 2909: ad; 9593: le; 10452: le; 10521: ad.
- Jaramillo-Mejía 7287: le; 7324: le – Jaramillo-Mejía & Van der Hammen 4149: le; 4199: le; 4248: le – Jaramillo-Mejía et al. 3454: ad; 5284: le – Johnson & Barkley 18c792: se – Juncosa & Misas 1094: ad.
- Killip 38464: le – Killip & García 33919: ad – Killip & Smith 15147: se – Kirkbride 2036: se – Kirkbride & Forero 1902: se.
- Leist & Mohle 2150: ad – Lellinger & De la Sota 813: ad – Little 9353: se – Llanos et al. 3369: le – Llanos & Botache 2461: ad; 2500: ad – Llanos & Camacho 1797: ad – Llanos & Gerardino 2891: le – Londoño et al. 139: le – López-Jurado & Riascos 565: le; 575: le; 582: le; 604: le – Lozano et al. 4236: le; 4514: le – Luteyn & Escobar 13266: le.
- MacDougal & Roldán 3589: le; 3627: se – Medina 1116: le – Mejía 112a: le; 121: se – Mendoza 1938: le – Mendoza & Betancur 4643: ad; 4956: ad – Mora 4456: ad; 4461: le; 4564: le – Morales & Giraldo 3953: se – Munar 711: ad; 1336: ad – Muñoz s.n. 699: le – Muñoz et al. 1739: le; 1917: ad – Murillo 1500: se – Murillo & Jaramillo-Mejía 1127: le; 1324: se – Murillo et al. 736: ad; 1517: ad – Murillo-A. 3648: ad.
- Nee & Mori 3833: se.
- Osorio 78: ad; 156: le – Ótalora 239: le; 262: le; 278: se.
- Pérez 6: ad; 59: le – Posada et al. 315: le – Prado & Pinilla 67: le.
- Ramírez-P. 3464: le; 4662: le; 6219: le; 7131: se – Ramírez-P. & Cuayal 3910: le – Ramírez-P. & Díaz 12219: ad – Ramírez-P. & Drosero 12013: se – Ramos 1163: ad – Restrepo de Fraume 102: ad; 110: ad; 343: ad – Ríos 16: ad – Rivera et al. 83: le – Rivera-Díaz et al. 970: le – Rodríguez 3100: le; 3122: se; 3125: ad; 3199: se; 3560: ad; 4835: ad; 6103: se – Rodríguez & Arboleda 3452: le – Rodríguez et al. 3251: se; 3387: le; 3651: le; 3667: se; 3693: se; 4835: ad; 4875: se; 4878: le; 4907: ad; 5011: le; 5475: le; 5556: le; 5881: le; 5987: se; 6006: se – Roldán et al. 34: se; 41: le; 64: le; 68: ad – Romero-Castañeda 7399: le; 7545: le – Ruiz & Correa 934: le – Ruiz & Rengifo 1082: le.
- Sandoval et al. 159: le – Sanín 653: ad; 768: le; 1643: ad; 1714: ad; 1736: ad; 2157: ad; 2805: le; 2806: ad – Sanín & Mancera 949: le – Sanín & Sanín 2704: le; 3208: se; 3218: le; 3262: le; 3282: se – Sanín et al. 1596: le; 2442: le; 2446: ad; 2495: le; **2646**: ma; 2774: ad; 2794: ad; 3061: ad; 3065: ad; 3076: ad; 3080: ad; 3089: ad; 3103: le; 3177: le; 3315: le; 4240: le; 5700: le – Sarria 740: le – Silva et al. 129: ad; 237: ad; 557: ad – Silverstone-Sopkin & Giraldo-G. 6414: le – Skolnik et al. 353: ad – Soejarto 3040: se.
- Torres et al. 2147a: ad – Triana-Moreno 60: ad.
- Vargas 7419: ad – Vélez 3425: le; 3670: le; 5028: le; 5826: le.
- Wilde 2542: le.
- Zarucchi et al. 4357: le; 4633: le; 5592: le; 6290: se; 6816: le; 6846: le; 7011: le.