



Hoya wongii (Apocynaceae, Asclepiadoideae): a new campanulate flowered species from Brunei (Borneo)

M. Rodda¹, N. Simonsson², L. Wanntorp³

Key words

Apocynaceae
Asclepiadoideae
Borneo
Brunei
campanulate corolla
corona
Hoya wongii

Abstract A new species from kerangas heath forest of Brunei, Borneo, *Hoya wongii*, is here described and illustrated. This species differs from the previously described species of *Hoya* in having a corona with inflated outer processes and laterally compressed and erect inner processes, ending in two bidentate membranaceous appendages. The corona is inserted above a pale yellow campanulate corolla. We discuss morphological affinities between the new species and other campanulate flowered species of *Hoya*.

Published on 12 August 2011

INTRODUCTION

Hoya R.Br is a genus of tropical and subtropical plants, generally epiphytic, lactiferous climbers with fleshy, sometimes succulent leaves. *Hoya* has a wide Indomalayan-Australian-Western Pacific distribution with the highest species diversity in the Philippines and New Guinea (Tsiang & Li 1977, Li et al. 1995, Forster & Liddle 1996, Forster 1996, Wanntorp et al. 2006a, b).

Presently, no comprehensive taxonomic revision is available for the genus (Meve 2002). Due to the lack of taxonomic studies on the whole genus, the number of species in *Hoya* is difficult to estimate. There are more than 500 species names listed in the International Plant Names Index (September 2010). This list, however, includes many synonyms and a number of c. 300 species for *Hoya* is probably more close to reality (Forster & Liddle 1996). In the past ten years, more than ten *Hoya* species have been described from Borneo. Among these, only one species, the newly described *H. danumensis* Rodda & Nyhuus (2009) has a strictly campanulate corolla. In Borneo two species with semi-campanulate corollas have been recorded, *H. phyllura* O.Schwartz and *H. nyhuusiae* Kloppenb. (Kloppenb. 2003); *H. vacciniiflora* O.Schwartz has a campanulate-urceolate corolla. Species with campanulate or semi-campanulate corollas are also found in other areas of the genus distribution (Rodda & Nyhuus 2009).

We studied herbarium specimens of *Hoya* at A, B, BM, BRUN, E, FI, HBG, K, L, P, SAN, SING, SNP and TO to provide evidence for a new campanulate flowered species from Brunei, which we describe below.

DESCRIPTION

Hoya wongii Rodda, Simonsson & L.Wanntorp, *sp. nov.* —
Fig. 1, 2

Ad Hoyam campanulatam Blume similis ob corollam campanulatam, sed differt coronae lobis inferius inflatis et superne erectis, compressis cum binis appendicibus membranaceis bidentatis. — Typus: *K.M. Wong* (KMW) 1566 (holo BRUN; iso K, L, SAN, SING), Borneo, Brunei, Tutong district, Pasir Puteh, 5 Oct. 1989.

Etymology. *Hoya wongii* is named after its first collector Dr. Wong Khoon Meng, who has greatly contributed to the knowledge of the SE Asian flora.

All measurements from rehydrated material.

Semi-woody, slender glabrous vine with white latex. *Leafy stems* cylindrical 2.5–3 mm diam, older stems leafless, bark waxy, peeling at least on dry material. *Internodes* 15–150 mm, adventitious roots absent. *Leaves* opposite, petiolate; petiole 3–7 by 0.7–1.2 mm; lamina coriaceous, flexible, not succulent, elliptic, 50–90 by 20–40 mm, widest in the central portion, apex apiculate-cuspidate, base shortly attenuate, margin entire, penninerved, main vein depressed on adaxial surface, evident on abaxial surface, secondary veins 5–7 pairs evident when dry, anastomosing near leaf margins. *Inflorescences* pseudo-umbelliform, 1-flowered; peduncle extra-axillary, perennial, about 10 mm long, pedicels 15–25 by 0.4–1 mm, glabrous. *Calyx* c. 5 mm diam, sepals oblong, about 2 by 1.6 mm, apex rounded, margins ciliate. *Corolla* campanulate, pale yellow, 25–30 mm diam; corolla lobe fused for c. 18 by 16 mm, free lobes c. 5 mm long with acute tips. *Corona* staminal pale yellow, 5–6 mm high, 6 mm diam, outer processes broadened into a rounded free process with revolute lateral margins, inner processes laterally compressed, ending in two bidentate membranaceous appendages, about 2 mm higher than outer processes and 0.3 mm higher than inner processes; anther skirt revolute, free from the filament tube. *Guide rail* forming a ridge, laterally compressed, prominent at the base of the anther skirt and extending 0.3–0.5 mm laterally. *Pollinarium* c. 1 mm long, pollinia oblanceolate, 830 by 300 µm, apex rounded to obtuse, retinaculum 280 by 180 µm, caudicles 120 µm long. *Ovary* rounded, 2.5 mm long. *Fruits and seeds* not seen.

Distribution — Brunei, possibly other areas of Borneo.

¹ Department of Plant Biology, Viale P. A. Mattioli 25, 10125, Torino, Italy; current address: Singapore Botanic Gardens Herbarium, 1 Cluny Road, Singapore 259569; corresponding author e-mail: rodda.michele@gmail.com.

² Hästhagevägen 1, 461 58 Trollhättan, Sweden.

³ Department of Phanerogamic Botany, Swedish Museum of Natural History Stockholm, P.O. Box 50007, SE-104 05 Stockholm, Sweden.




Fig. 1 *Hoya wongii* Rodda, Simonsson & L.Wanntorp. a. Flowering branch; b. pollinarium; c. corolla lobes; d. corona, underside; e. corona, side view (M. Rodda delineavit) (prepared from the holotype). — Scale bars: a = 1 cm; b = 200 µm; c = 5 mm; d, e = 1 mm.

Habitat & Ecology — A climbing epiphyte only found in the type locality. The type locality is in a coastal kerangas heath forest with white sand, 15–30 m above sea level.

Additional specimens seen. *Hoya campanulata* Blume *sin. coll.*, L0004389 (type L), Java, Ex Hort Bogor. *Hoya collettii* (Collett & Hemsl.) Schltr., *Collett* 833 (type K), Myanmar, Shan Hills, 6000 ft, June 1888. *Hoya manipurensis* Deb, *Micholitz s.n.* (holotype of *Micholitzia obcordata* N.E.Br., K), 'India & Burma', cult. Royal Bot. Gard. Glasnevin, Dublin, 1909. *Hoya danumen-*

sis Rodda & Nyhuus, *M. Rodda 2008.1h* (holotype FI) ex hort., 3 Sept. 2008. *Hoya phyllura* O.Schwartz, *Winkler* 339 (type HBG), Lebang Hara, West Borneo, 24 Nov. 1924. *Hoya telosmoides* Omlor, *Clemens* 29828 (holotype BM), Malaysia, Sabah, Mt Kinabalu, Tenompok, 1500 m, 7 June 1932. *Hoya vacciniiflora* O.Schwartz, *Winkler* 518 (type HBG), Bukit Mulu, West Borneo, 2 Dec. 1924.

Conservation status — *Hoya wongii* should be considered Data Deficient (DD) according to IUCN Red List criteria (IUCN




Fig 2 *Hoya wongii* Rodda, Simonsson & L.Wanntorp. The peculiar single-flowered inflorescences are evident on this specimen (holotype in BRUN herbarium). Photo by Dr. Jacqueline Henrot, BRUN.

2001) because it is known from only one collection and thus remains in need of further investigation with respect to future conservation efforts. This species was found in kerangas heath forest, a nutrient poor habitat rarely used for cultivation. However, felling and burning activities may degrade kerangas into an open savanna of shrubs (Whitmore 1984) making it inhospitable for many epiphytic plant species including *Hoya*.

Notes — Due to its broadly campanulate corolla, specimens of *H. wongii* can be easily confused with specimens of *H. campanulata*, *H. danumensis*, *H. nyhuusiae*, *H. phyllura* or *H. vacciniiflora*. However, none of these species have coronas showing the same morphological characters as in *H. wongii*. Here, the corona has broadly inflated outer processes and laterally compressed inner processes with two bidentate membranaceous appendages. All other Bornean species of *Hoya*

with a campanulate or semi-campanulate corolla have spreading elongated corona lobes and a well-developed anther skirt forming a disk at the base of the corona. Among species with a campanulate to semi-campanulate corolla, *H. collettii*, resembles *H. wongii* in the morphology of the corona. This species, which is endemic to Myanmar, is so far only known from the type specimen. It differs from *H. wongii* in being a small shrub with very thick, narrowly lanceolate leaves and not a vine with thin, elliptic leaves. Further, *H. collettii* has rounded outer corona processes much less developed than those of *H. wongii*. The outer corona processes are very elongated but not laterally compressed and lack the two bidentate appendages of *H. wongii*.

Two additional species of *Hoya*, one again endemic to Borneo and the other found in India, Thailand and China, show a slightly similar corona to that of the new species. The first, *H. telosmoides* (Omlor 1996), from Borneo, has only been collected from Mt Kinabalu, to which it is possibly restricted. The corolla in *H. telosmoides* is not campanulate as that of *H. wongii* but urceolate and hairy on the inside. These two species have similar coronas, with well-developed erect outer processes. The inner coronal processes of *H. telosmoides* are, however, much less developed than those of *H. wongii* and lack the bidentate appendages. The second species, *H. manipurensis* (Deb 1955), is a species growing at low to middle altitudes in cool sub-tropical zones of India, China and Thailand. It also has an urceolate corona similar to that of *H. telosmoides*. In addition, *H. manipurensis* also differs in habit, being an epiphytic shrub and not a vine, and in having c. 10 mm long greenish red tubular flowers instead of pale yellow campanulate flowers.

Acknowledgements This study is part of an ongoing research project on the systematics of *Hoya*. Financial support has been received from the SYNTHESYS programme, grants no GB-TAF-5657, NL-TAF-676 and DE-TAF-675 to MR. We would like to thank the curators of the following herbaria: A, B, BM, BRUN, E, FI, HBG, K, L, P, SAN, SING, SNP and TO for help with material and for providing high quality images of herbarium specimens; G. Pandolfo for help with the latin diagnosis; Anita Walsmit Sachs who much improved the line drawing and two anonymous reviewers for their valuable comments on the manuscript.

REFERENCES

- Deb DB. 1955. *Hoya manipurensis*. Journal of the Indian Botanical Society 34: 50.
- Forster PI. 1996. A checklist of the Asclepiadaceae of Papuasia. Science in New Guinea 22, 1: 15–22.
- Forster PI, Liddle DJ. 1996. Flora of Australia 28: 231–237. CSIRO, Canberra.
- International Plant Name Index. 2010. <http://www.ipni.org> [accessed September 2010].
- IUCN. 2001. IUCN Red List categories and criteria, version 3.1. Prepared by the IUCN Species Survival Commission. Gland & Cambridge.
- Kloppenborg D. 2003. Our cover story, *Hoya nyhuusiae* Kloppenborg. Fraterna 16, 3: 1–6.
- Li PT, Gilbert MG, Stevens WD. 1995. Asclepiadaceae. Flora of China 16: 228–236. Science Press, Beijing & Missouri Botanical Garden Press, St. Louis.
- Meve U. 2002. Species numbers and progress in asclepiad taxonomy. Kew Bulletin 57, 2: 459–464.
- Omlor R. 1996. Notes on Marsdenieae (Asclepiadaceae): A new, unusual species of *Hoya* from Northern Borneo. Novon 6, 3: 288–294.
- Rodda M, Nyhuus T. 2009. *Hoya danumensis*, a new species of *Hoya* (Apocynaceae, Asclepiadoideae) from Borneo. Webbia 64, 2: 163–167.
- Tsiang Y, Li PT. 1977. Asclepiadaceae. Flora Reipublicae Popularis Sinicae 63: 475–492. Science Press Beijing. [In Chinese].
- Wanntorp L, Kocyan A, Renner SS. 2006a. Wax plants disentangled: A phylogeny of *Hoya* (Marsdenieae, Apocynaceae) inferred from nuclear and chloroplast DNA sequences. Molecular Phylogenetics and Evolution 39: 722–733.
- Wanntorp L, Kocyan A, Van Donkelaar R, Renner SS. 2006b. Towards a monophyletic *Hoya* (Marsdenieae, Apocynaceae): Inferences from the chloroplast trnL region and the rbcL-atpB spacer. Systematic Botany 31, 3: 586–596.
- Whitmore TC. 1984. Tropical rainforests of the Far East: 352. 2nd edition. Oxford University Press, Oxford.