



Novel *Curvularia* species from clinical specimens

H. Madrid¹, K.C. da Cunha², J. Gené², J. Dijksterhuis¹, J. Cano², D.A. Sutton³,
J. Guarro², P.W. Crous^{1,4}

Key words

Bipolaris
Curvularia
gpd
ITS
LSU
phylogeny
RPB2
systematics

Abstract The fungal genus *Curvularia* includes numerous plant pathogens and some emerging opportunistic pathogens of humans. In a previous study we used morphology and sequences of the nuclear ribosomal internal transcribed spacer region (ITS) and the glyceraldehyde-3-phosphate dehydrogenase (*gpd*) gene to identify species within a set of 99 clinical *Curvularia* isolates from the USA. Seventy-two isolates could be identified while the remaining 27 isolates belonged in three unclassified clades that were tentatively labelled *Curvularia* sp. I, II and III. In the present study, we further assess the taxonomic placement of these isolates using sequences of ITS, *gpd*, the large subunit rDNA, and the second largest subunit of RNA polymerase II. DNA sequence comparisons with a set of 87 isolates representing 33 *Curvularia* spp. and members of the closely-related genera *Bipolaris* and *Exserohilum* revealed that *Curvularia* sp. I, II and III represent novel lineages in *Curvularia*. These lineages are morphologically different from the currently accepted species. In the phylogenetic tree, *Curvularia* sp. I and sp. III were each split into two distinct lineages. Morphology and phylogeny supported the proposal of five new species, to be named *C. americana*, *C. chlamydospora*, *C. hominis*, *C. muehlenbeckiae* and *C. pseudolunata*. The concatenated 4-locus phylogeny revealed the existence of six clades in *Curvularia*, which are associated with particular morphological features. They were named after representative species, namely *americana*, *eragrostidis*, *hominis*, *lunata*, *spicifera* and *trifolii*.

Article info Received: 2 February 2014; Accepted: 5 March 2014; Published: 29 July 2014.

INTRODUCTION

Curvularia, typified by *C. lunata*, is a species-rich genus, which includes numerous grass pathogens and saprobes occurring on plant material, dung and soil (Faurel & Schotter 1965, Sivanesan 1987, Jiang & Zhang 2007). At least eight species of this genus have been reported from opportunistic diseases in humans ranging from mild skin and nail infections to severe invasive disease, depending on route of infection and immune status of the host (Kamalam et al. 1992, Ismail et al. 1993, Lopes & Jobim 1998, Ebright et al. 1999, de Hoog et al. 2000). Morphologically, *Curvularia* is characterised by the production of sympodial conidiophores with tretic, terminal and intercalary conidiogenous cells and elongate, transversely septate conidia with a dark basal scar. Conidia are often curved at an asymmetrically swollen intermediate cell, but species with straight conidia also have been described (Sivanesan 1987). Authors such as Ellis (1971, 1976), de Hoog et al. (2000) and Revankar & Sutton (2010) have described the conidia as truly septate or 'euseptate', i.e. composed of a single wall with septa that are formed as inward extensions of that wall (Luttrell 1963). A similar genus is *Bipolaris*, type species *B. maydis*, which traditionally has been distinguished from *Curvularia* by producing conidia which lack an asymmetrically swollen intermediate cell and are 'distoseptate' (Domsch et al. 2007, Revankar & Sutton 2010), i.e. they have a common outer wall enclosing more or less spherical cells, each of which is surrounded by an individual

wall (Luttrell 1963). The separation of the two genera has been a matter of controversy and many authors have stated that *Curvularia* species also have distoseptate conidia (Alcorn 1983a, Sivanesan 1987, Seifert et al. 2011).

Sexual stages of *Bipolaris* and *Curvularia* were traditionally placed in *Cochliobolus*. Typically, they feature thick-walled, ostiolate ascomata with pseudoparaphyses, and bitunicate asci that give rise to filiform, multiseptate ascospores (Sivanesan 1987, Zhang et al. 2012). The ascospores often appear more or less helically coiled within the ascus. A similar genus, *Pseudocochliobolus*, was segregated from *Cochliobolus* to accommodate species producing ascomata on columnar stromata, with ascospores appearing linearly parallel or loosely coiled within the asci. The asexual stages of *Pseudocochliobolus* species were *Curvularia* and *Bipolaris* species with short, rather straight conidia (Tsuda et al. 1977, Tsuda & Ueyama 1981). Most authors have not accepted *Pseudocochliobolus* as a separate genus because the degree of coiling of the ascospores can vary greatly within a species. Also, the addition of a second genus with *Curvularia* and *Bipolaris* asexual stages would introduce unnecessary complexity into the taxonomy of this group of fungi instead of clarifying it (Alcorn 1983a, 1988, Sivanesan 1987).

Cochliobolus, *Pseudocochliobolus* and their *Bipolaris* and *Curvularia* asexual morphs were previously considered to be related either to the *Dothideales* (Eriksson 1981) or to the *Pleosporales* (Barr 1979, Sivanesan 1984). Molecular data confirmed their placement in the latter order and more precisely in its largest family, *Pleosporaceae*, along with other important genera of plant pathogens and clinically-relevant fungi such as *Alternaria* and *Exserohilum* (Olivier et al. 2000, Zhang et al. 2009, 2012). Berbee et al. (1999) performed a phylogenetic study to assess the evolutionary relationships of *Cochliobolus*, *Pseudocochliobolus*, *Curvularia* and *Bipolaris*. Their phylogenetic trees, based on the internal transcribed spacer (ITS)

¹ CBS-KNAW Fungal Biodiversity Centre, Uppsalalaan 8, 3584 CT Utrecht, The Netherlands; corresponding author e-mail: h.madrid@cbs.knaw.nl.

² Mycology Unit, Medical School and IISPV, Universitat Rovira i Virgili, 43201 Reus, Spain.

³ Fungus Testing Laboratory, University of Texas Health Science Center, San Antonio, TX 78229-3900, USA.

⁴ Wageningen University and Research Centre (WUR), Laboratory of Phytopathology, Droevendaalsesteeg 1, 6708 PB Wageningen, The Netherlands.

region of the rDNA and the glyceraldehyde-3-phosphate dehydrogenase (*gpd*) gene, revealed that isolates were distributed mainly in two clades which were named 'Cochliobolus groups 1 and 2'. Group 1 exclusively encompassed species with *Bipolaris* asexual morphs, including the type species, *B. maydis*, agent of southern corn leaf blight, as well as other economically-relevant phytopathogenic species. The sexual morph of *B. maydis* is *Cochliobolus heterostrophus*, type species of *Cochliobolus* (Sivanesan 1987). Group 2 included mostly plant pathogens and saprobes with *Bipolaris* and *Curvularia* asexual morphs, including the type species of the latter genus, *C. lunata* and all species of *Pseudocochliobolus*.

Manamgoda et al. (2012), with a wider sampling of species and based on the analysis of ITS, large subunit (LSU) rDNA, *gpd* and elongation factor 1- α (EF1- α) genes, applied the one fungus = one name concept (Hawksworth et al. 2011) to the *Bipolaris-Curvularia*, *Cochliobolus-Pseudocochliobolus* complex. Their phylogenies confirmed the existence of the same two main groups reported by Berbee et al. (1999). Based on those results, *Cochliobolus* and *Pseudocochliobolus* were synonymized with the more commonly used generic names *Bipolaris* and *Curvularia*, respectively, and the generic concept of the latter genus was expanded to accommodate some species with rather straight conidia formerly placed in *Bipolaris* but grouping in the *Curvularia* clade (Manamgoda et al. 2012). These included important agents of opportunistic infections in vertebrates, such as *B. australiensis*, *B. hawaiiensis* and *B. spicifera* (de Hoog et al. 2000). The last of these had been previously considered a *Curvularia* species by Boedijn (1933).

Curvularia spp. have been identified mostly based on morphology, but the names applied often do not correlate with DNA sequence-based identifications. Furthermore, the species most commonly reported from humans, *C. lunata*, appeared to be a species complex (Berbee et al. 1999, Yanagihara et al. 2010). Da Cunha et al. (2013) recently characterised a set of 99 clinical *Curvularia* strains from the USA using sequences of the ITS region and the *gpd* gene. They could identify 73.2 % of the isolates, including *C. aerea*, which was the most common species. The remaining isolates were distributed over three different lineages which did not correlate with any known species. In this study we used DNA sequence data of four nuclear loci to further assess the taxonomic position of these isolates.

MATERIALS AND METHODS

Fungal isolates

Twenty-seven clinical *Curvularia* isolates from the USA were studied (Table 1). These isolates were obtained from the Fungus Testing Laboratory, University of Texas Health Science Center, San Antonio, and represent the clades named *Curvularia* sp. I, II and III in the study by da Cunha et al. (2013). These isolates were compared with ex-type or reference strains of different *Curvularia* spp. from the CBS-KNAW Fungal Biodiversity Centre, Utrecht, The Netherlands.

Phenotypic study

Colony morphology and growth rates were studied on potato carrot agar (PCA; 20 g of potatoes, 20 g of carrots, 20 g of agar, 1 L of distilled water) and oatmeal agar (OA; 30 g of filtered oat flakes, 20 g of agar, 1 L of distilled water) after 7 d of incubation at 25 °C in the dark. Microscopic features were studied in lactic acid from colonies on the same media after 10–21 d of incubation. Size ranges in the species descriptions are derived from at least 30 measurements.

Cryo-Scanning Electron microscopy

Relevant areas of fungal cultures were carefully selected by means of a stereo microscope (Nikon SMZ1500, Nikon, Amsterdam, The Netherlands). Small (c. 3 × 5 mm) agar blocks were carefully cut out with a surgical blade (no. 11, Swan-Morton, Sheffield, UK), while disturbing of fungal structures was kept to a minimum during cutting and transferring of the samples to a copper cup (diam 10 mm, height 8 mm). Agar blocks were glued to the copper cup with frozen tissue medium (KP-Cryoblock, Klinipath, Duiven, The Netherlands). The copper cup was placed on an agar surface inside a closed Petri dish to prevent drying of the sample. The sample was quickly frozen in nitrogen slush and immediately transferred to a JEOL 5600LV scanning electron microscope (JEOL, Tokyo, Japan) equipped with an Oxford CT1500 cryostation. The sample was viewed at 2.5 kV and ice was removed by sublimation after heating of the SEM-stage to -85 °C. Then the sample was sputter-coated in the cryostation by means of a gold target for three times 90 s holding the sample at different angles for an optimal coating. Electron micrographs were acquired with the F3 or F4 scan at 5 kV and contrast levels digitally enhanced in Adobe® Photoshop® Creative Suite v. 6.

Molecular study

DNA extraction of *Curvularia* spp. I–III was performed with the PrepMan Ultra sample preparation reagent (Applied Biosystems, Foster City, CA, USA) as described by da Cunha et al. (2013). DNA extraction of isolates of the other species studied was carried out from colonies growing on malt extract agar (Oxoid, Basingstoke, England) with the UltraClean® Microbial DNA Isolation Kit (Mo Bio Laboratories, Inc., Solana Beach, CA, USA). Amplification and sequencing of the ITS and RNA polymerase II second largest subunit (RPB2) were performed with primers ITS5 + ITS4 (White et al. 1990) and 5F2 + 7cR (O'Donnell et al. 2007) following the protocols of Amaradasa et al. (2014). Amplification of the *gpd* and LSU genes were performed with primers *gpd*1 + *gpd*2 (Berbee et al. 1999) and LR0R + LR5 (Vilgalys & Hester 1990) as described in Manamgoda et al. (2012). The ITS PCR products were purified and sequenced at Macrogen Europe (Amsterdam) using a 3739 XL DNA analyser (Applied Biosystems). The *gpd*, LSU and RPB2 loci were sequenced at the CBS-KNAW Fungal Biodiversity Centre (Utrecht, The Netherlands), using the BigDye terminator sequencing kit v. 3.1 (Applied Biosystems) and an ABI Prism™ 3100 DNA sequencer (Applied Biosystems). The program SeqMan Pro (Lasergene, Madison, WI, USA) was used to obtain consensus sequences from the complementary sequences of each isolate. Sequences of the clinical isolates were aligned with those of a set of 60 isolates representing 33 species of *Curvularia*, and two phylogenetically related genera of *Pleosporaceae*, i.e. *Bipolaris* (nine spp.) and *Exserohilum* (one sp., used as outgroup) using ClustalX v. 1.81 (Thompson et al. 1997), followed by manual adjustments with a text editor. Individual alignments of ITS, LSU, *gpd* and RPB2 and a concatenated 4-locus dataset were analysed with maximum likelihood (ML) using MEGA5 (Tamura et al. 2011) with partial deletion of gaps, substitution models proposed by this program and 1 000 bootstrap replicates. Bootstrap support values (bs) ≥ 70 % were considered significant. Incongruence among datasets was tested by a visual inspection of all groups with ≥ 70 bs in the partial trees to search for potentially conflicting groups. A Markov Chain Monte Carlo (MCMC) algorithm was used to generate phylogenetic trees with Bayesian probabilities using MrBayes v. 3.1.1 (Ronquist & Huelsenbeck 2003). The best models of nucleotide substitution for each locus for the Bayesian analysis were determined using MrModeltest v. 2.3 (Nylander 2004). Two analyses of four MCMC chains were run from

Table 1 Isolates included in the phylogenetic study, their origins, and GenBank accession no.

Taxon	Isolate no. ¹	Source	GenBank accession no. ²			
			ITS	LSU	<i>gpd</i>	RPB2
<i>Bipolaris chloridis</i>	CBS 242.77B	<i>Chloris gayana</i> , Australia	HF934928	HF934869	HG779083	HF934830
<i>B. cynodontis</i>	CBS 285.51	<i>Cynodon transvaalensis</i> , Kenya	HF934929	HF934874	HG779081	HF934831
	CBS 305.64	<i>Cynodon dactylon</i> , USA	HF934930	HF934883	HG779082	HF934832
<i>B. maydis</i>	CBS 130.26	Unknown	HF934923	HF934873	HG779084	HF934825
	CBS 136.29	<i>Zea mays</i> , Japan	HF934926	HF934879	HG779086	HF934828
	CBS 307.64	<i>Zea mays</i> , USA	HF934925	HF934875	HG779085	HF93482
<i>B. microlaenae</i>	CBS 280.91 ^T	<i>Microlaena stipoides</i> leaf, Australia	HF934933	HF934877	HG779092	HF934835
<i>B. oryzae</i>	CBS 157.50	<i>Oryza sativa</i> grain, Indonesia	HF934931	HF934870	HG779090	HF934833
	CBS 199.54	<i>Oryza sativa</i> grain, New Guinea	HF934932	HF934884	HG779091	HF934834
<i>B. sorghicola</i>	CBS 249.49	<i>Sorghum vulgare</i> var. <i>sudanense</i> , Locality unknown	HF934927	HF934868	HG779087	HF934829
<i>B. sorokiniana</i>	CBS 140.31	Substrate unknown, Japan	HF934935	HF934876	HG779088	HF934837
	CBS 145.32	<i>Triticum durum</i> , Locality unknown	HF934934	HF934885	HG779089	HF934836
<i>B. zeae</i>	CBS 1277.16	Unknown	HG778980	HG779027	HG779095	HG779158
<i>B. zeicola</i>	CBS 316.64	<i>Zea mays</i> , USA	HF934938	HF934871	HG779093	HF934840
	CBS 317.64	<i>Zea mays</i> , USA	HF934939	HF934878	HG779094	HF934841
<i>Curvularia aeria</i>	CBS 294.61 ^T	Air, Brazil	HF934910	HF934902	HF565450	HF934812
<i>C. affinis</i>	CBS 154.34 ^T	<i>Manihot utilissima</i> , Java	HG778981	HG779028	HG779126	HG779159
	CBS 185.49	<i>Manihot utilissima</i> , Java	HG778982	HG779029	HG779127	HG779160
<i>C. akaii</i>	CBS 318.86	Substrate unknown, Japan	HF934921	HF934897	HG779118	HF934823
	CBS 127728	Substrate unknown, Japan	HF934920	HF934898	HG779119	HF934822
	CBS 127730	Substrate unknown, Japan	HF934922	HF934899	HG779120	HF934824
<i>C. australiensis</i>	CBS 172.57	<i>Oryza sativa</i> seed, Vietnam	HF934912	HF934901	HG779139	HF934814
<i>C. brachyspora</i>	CBS 186.50	Soil, Java	HG778983	HG779030	HG779150	HG779161
<i>C. carica-papayae</i>	CBS 135941 ^T	<i>Carica papaya</i> leaf, India	HG778984	HG779031	HG779146	HG779162
<i>C. coicis</i>	CBS 192.29 ^T	<i>Coix lacrima-jobi</i> var. <i>typica</i> , Japan	HF934917	HF934895	HG779130	HF934819
<i>C. cymbopogonis</i>	CBS 419.78	<i>Yucca</i> sp. leaf, Netherlands	HG778985	HG779032	HG779129	HG779163
<i>C. ellisii</i>	CBS 193.62	Air, Pakistan	HF934913	HF934896	HG779143	HF934815
<i>C. eragrostidis</i>	CBS 189.48	Sorghum seed, Java	HG778986	HG779033	HG779154	HG779164
<i>C. gladioli</i>	CBS 210.79	<i>Gladiolus</i> sp. leaf, Romania	HG778987	HG779034	HG779123	HG779165
<i>C. graminicola</i>	BRIP 23186	<i>Aristida ingrata</i> , Australia	JN192376	JN600986	JN600964	–
<i>C. hawaiiensis</i>	CBS 173.57 ^T	<i>Oryza sativa</i> , Hawaii	HG778988	HG779035	HG779140	HG779166
	CBS 448.72	Salt-marsh soil, Kuwait	HG778989	HG779036	HG779142	HG779167
	CBS 727.96	Substrate unknown, USA	HG778990	HG779037	HG779141	HG779168
<i>C. heteropogonis</i>	CBS 284.91 ^T	<i>Heteropogon contortus</i> leaf, Australia	HF934919	HF934893	HF934919	HF934821
	CBS 511.91	<i>Heteropogon contortus</i> leaf, Australia	HF934918	HF934894	HF934918	HF934820
<i>C. intermedia</i>	CBS 334.64	<i>Avena vesicolor</i> , USA	HG778991	HG779038	HG779155	HG779169
<i>C. ischaemi</i>	CBS 630.82 ^T	<i>Ischaemum indicum</i> leaf, Solomon Islands	HG778992	HG779039	HG779131	HG779170
<i>C. lunata</i>	CBS 730.96 ^{MT}	Lung biopsy, USA	HF934911	HF934900	JX256429	HF934813
<i>C. oryzae</i>	CBS 169.53 ^T	<i>Oryza sativa</i> seed, Vietnam	HF934906	HF934867	HF934808	HF934808
<i>C. ovariicola</i>	CBS 285.91	<i>Eragrostis parviflora</i> , Australia	HG778993	HG779040	HG779144	HG779171
	CBS 286.91	<i>Eragrostis parviflora</i> , Australia	HG778994	HG779041	HG779145	HG779172
<i>C. perotidis</i>	CBS 350.90 ^T	<i>Perotis rara</i> , Australia	HG778995	HG779042	HG779138	HG779173
<i>C. prasadii</i>	CBS 143.64 ^T	<i>Jasminum sambac</i> , India	HG778996	HG779043	HG779147	HG779174
	CBS 144.64	Substrate unknown, England	HG778997	HG779044	HG779149	HG779175
<i>C. protuberata</i>	CBS 376.65 ^T	<i>Deschampsia flexuosa</i> leaf, Scotland	HG778998	HG779045	HG779135	HG779176
<i>C. ravenelii</i>	CBS 127709	Unknown	HG778999	HG779046	HG779109	HG779177
<i>C. robusta</i>	CBS 624.68 ^T	<i>Dichanthium annulatum</i> leaf, USA	HG779000	HG779047	HG779125	HG779178
<i>C. senegalensis</i>	CBS 149.71	Substrate unknown, Nigeria	HG779001	HG779048	HG779128	HG779179
<i>C. spicifera</i>	CBS 198.31	<i>Capsicum anuum</i> , Cyprus	HF934916	HF934905	HG779136	HF934818
	CBS 199.31	<i>Cucurbita maxima</i> , Cyprus	HF934915	HF934903	HG779137	HF934817
<i>C. trifolii</i>	CBS 173.55	<i>Trifolium repens</i> , USA	HG779023	HG779077	HG779124	HG779208
<i>C. tripogonis</i>	BRIP 12375 ^T	<i>Tripogon jacquemonti</i> , India	JN192388	JN601002	JN600980	–
<i>C. tuberculata</i>	CBS 146.63 ^T	<i>Zea mays</i> leaf, India	HF934907	HF934866	HG779157	HF934809
<i>C. uncinata</i>	CBS 221.52 ^T	<i>Oryza sativa</i> leaf, Vietnam	HG779024	HG779078	HG779134	HG779209
	CBS 531.70	<i>Oryza sativa</i> seeds, Denmark	HG779025	HG779079	HG779132	HG779210
<i>C. verruciformis</i>	CBS 537.75	<i>Lobibyx</i> sp. feather, New Zealand	HG779026	HG779080	HG779133	HG779211
<i>C. verruculosa</i>	CBS 149.63	<i>Elaeis guineensis</i> , Nigeria	HF934909	HF934891	HG779110	HF934811
	CBS 150.63	<i>Punica granatum</i> leaf, India	HF934908	HF934892	HG779111	HF934810
<i>Curvularia</i> sp. I Lineage A (= <i>C. muehlenbeckiae</i> sp. nov.)	CBS 144.63 ^T	<i>Muehlenbeckia</i> sp. leaf, India	HG779002	HG779049	HG779108	HG779180
	UTHSC 08-2905	Chest, USA	HE861836	HG779050	HF565484	HG779189
<i>Curvularia</i> sp. I Lineage B (= <i>C. hominis</i> sp. nov.)	UTHSC 07-2791	Cornea, USA	HG779003	HG779057	HG779105	HG779181
	UTHSC 07-3105	Nasal sinus, USA	HG779004	HG779058	HG779104	HG779182
	UTHSC 07-3184	Nasal sinus, USA	HG779005	HG779059	HG779099	HG779183
	UTHSC 07-3581	Nail, USA	HG779006	HG779060	HG779102	HG779184
	UTHSC 08-849	Eye, USA	HE861837	HG779051	HF565483	HG779185
	UTHSC 08-1296	Nail, USA	HG779007	HG779061	HG779103	HG779186
	UTHSC 08-2418	Bronchial wash, USA	HG779008	HG779062	HG779096	HG779187
	UTHSC 09-464 ^T	Cornea, USA	HG779011	HG779065	HG779106	HG779191
	UTHSC 08-2517	Foot, USA	HG779009	HG779063	HG779107	HG779188
	UTHSC 08-3737	Bronchial wash, USA	HG779010	HG779064	HG779101	HG779190
	UTHSC 09-1692	Nasal sinus, USA	HG779012	HG779066	HG779097	HG779192
	UTHSC 09-2197	Nasal sinus, USA	HE861835	HG779052	HF565485	HG779193
	UTHSC 09-2532	Nasopharynx, USA	HG779013	HG779067	HG779100	HG779194
	UTHSC 09-3403	Unknown tissue, USA	HG779014	HG779068	HG779098	HG779195

Table 1 (cont.)

Taxon	Isolate no. ¹	Source	GenBank accession no. ²			
			ITS	LSU	<i>gpd</i>	RPB2
<i>Curvularia</i> sp. II (= <i>C. americana</i> sp. nov.)	UTHSC 08-3414 [†]	Ankle, USA	HE861833	HG779056	HF565488	HG779200
	UTHSC 07-2649	Toe tissue, USA	HE861834	HG779054	HF565486	HG779196
	UTHSC 08-84	Nasal sinus, USA	HG779015	HG779069	HG779115	HG779197
	UTHSC 08-278	Peritoneal dialysis fluid, USA	HE861832	HG779055	HF565487	HG779198
	UTHSC 08-2697	Leg, USA	HG779016	HG779070	HG779117	HG779199
	UTHSC 09-2907	Toe nail, USA	HG779017	HG779071	HG779114	HG779201
	UTHSC 09-2806	Bone marrow, USA	HG779018	HG779072	HG779112	HG779202
	UTHSC 09-2863	Bronchial wash, USA	HG779019	HG779073	HG779113	HG779203
	UTHSC 10-1276	Maxillary sinus, USA	HG779020	HG779074	HG779116	HG779204
<i>Curvularia</i> sp. III Lineage A (= <i>C. chlamydo-<i>spora</i></i> sp. nov.)	UTHSC 07-2764 [†]	Toe nail, USA	HG779021	HG779075	HG779151	HG779205
	UTHSC 08-1283	Nasal sinus, USA	HG779022	HG779076	HG779152	HG779206
<i>Curvularia</i> sp. III Lineage B (= <i>C. pseudolunata</i> sp. nov.)	UTHSC 09-2092 [†]	Nasal sinus, USA	HE861842	HG779053	HF565459	HG779207
<i>Exserohilum turcicum</i>	CBS 330.64	<i>Zea mays</i> , USA	HF934950	HF934887	HG779153	HF934852

¹ BRIP: Queensland Plant Pathology Herbarium, Queensland, Australia; CBS: CBS-KNAW Fungal Biodiversity Centre, Utrecht, The Netherlands; UTHSC: Fungus Testing Laboratory, University of Texas Health Science Center, San Antonio, Texas; [†] ex-type strain; ^{NT} ex-neotype strain (Manamgoda et al. 2012).

² Sequences generated during this study appear in **bold**; other sequences originate from Manamgoda et al (2012), da Cunha et al. (2013) and Amaradasa et al. (2014).

random trees for 4 598 100 generations and sampled every 100 generations, resulting in 45 981 trees, of which 25 % were discarded as the burn-in phase. Posterior probabilities (pp) were determined from the remaining trees. The sequences generated during this study and the alignments used in the phylogenetic analyses were deposited in GenBank (Table 1) and TreeBASE (submission ID <http://purl.org/phylo/treebase/phyloids/study/TB2:S14881>), respectively.

RESULTS

Phylogenetic study

After removing ambiguously aligned regions, we obtained ITS, LSU, *gpd* and RPB2 alignments of 533, 830, 434, and 793 positions of which 64 (12 %), 39 (4.69 %), 111 (25.57 %) and 259 (32.66 %) were variable, respectively. MEGA5 proposed a K2 + G + I model for the ITS and RPB2 loci, K2 + I for LSU, T92 + G for *gpd* and GTR + G + I for the concatenated 4-locus dataset. These models were used in the ML analyses. Partial trees (not shown) were congruent except for the following clades: *Curvularia gladioli* CBS 210.79 grouped with *C. ischaemi* CBS 630.82 (93 % bs) in the ITS tree, but in the RPB2 tree the former isolate grouped with *Curvularia trifolii* CBS 173.55 (77 % bs), while the CBS 630.82 grouped with *Curvularia coicis* CBS 192.29 (100 % bs). These incongruencies affected species that are not closely related to *Curvularia* sp. I–III of da Cunha et al. (2013) and therefore the four loci were combined. Partial trees revealed that RPB2 was the most informative locus with 35 clades with significant bs, followed by *gpd* with 23. ITS and LSU both showed only 10 clades with significant bs. The ITS and LSU ML trees provided good support for a clade representing the genus *Bipolaris*, but *Curvularia* species appeared in several clades, some of which had low bootstrap support. The *gpd* ML tree separated *Bipolaris* and *Curvularia* as two clades with 93 % and 70 % bs, respectively, whereas these clades showed 99 % and 95 % bs in the RPB2 tree. In the concatenated 4-locus ML tree (not shown) the *Bipolaris* and *Curvularia* clades had 100 % and 97 % bs, respectively. For Bayesian analysis, MrModeltest proposed a SYM + I + G model for the ITS locus and GTR + I + G for LSU, *gpd* and RPB2. These models were incorporated in the analysis. The consensus tree obtained from the Bayesian analysis (Fig. 1) agreed with the topology of the ML tree (not shown) for the 4-locus dataset.

The 4-locus tree (Fig. 1) revealed that *C. carica-papayae*, listed as a synonym of *C. aeri* by Sivanesan (1987), is a phylogenetically distinct species. The concatenated tree also corroborated

that isolates in *Curvularia* spp. I–III of da Cunha et al. (2013) are different from accepted species of this genus represented in the CBS collection (Fig. 1). However, *Curvularia* spp. I and III were each split into two lineages that are sufficiently distant from each other to represent different species. These lineages were named here Ia, Ib and IIIa, IIIb, accordingly. One of them, Ib, shows considerable genetic variation, but is treated here as a single taxon because its complex topology does not seem to suggest a clear separation of species within it. The ITS and LSU ML trees did not provide enough resolution to separate lineages within *Curvularia* spp. I and III, but showed 87 % and 99 % bs for *Curvularia* sp. II. The *gpd* ML tree gave 80 % bs to *Curvularia* sp. II and separated lineages Ia (98 % bs) and Ib (52 % bs) of *Curvularia* sp. I, but did not separate the two lineages of *Curvularia* sp. III. The RPB2 ML tree gave 100 % bs to *Curvularia* sp. II and provided enough resolution to separate lineages Ia, Ib, IIIa and IIIb with bs \geq 75 %. *Curvularia* sp. Ia, Ib, II, IIIa and IIIb are morphologically and phylogenetically different from other members of the genus and therefore are proposed here as new taxa. These species were respectively named *C. muehlenbeckiae*, *C. hominis*, *C. americana*, *C. chlamydo-*spora** and *C. pseudolunata* and described in alphabetical order in the Taxonomy section.

Within the *Curvularia* clade, six well-supported lineages were associated with certain combinations of morphological features. These lineages were named the *americana*-, *eragrostidis*-, *hominis*-, *lunata*-, *spicifera*- and *trifolii*-clades (Fig. 1). The *eragrostidis*-clade (82 % bs, 1 pp) was formed by *C. eragrostidis*, *C. graminicola* and *C. intermedia*, and is characterised by producing inconspicuously distoseptate (i.e. the two cell wall layers within the conidium are difficult to distinguish in mature conidia), straight to somewhat unequal-sided, 4-celled conidia. In mature conidia of *C. graminicola*, all septa are accentuated by dark transverse bands, whereas in *C. eragrostidis* and *C. intermedia*, only the median septum is accentuated (Sivanesan 1987, Alcorn 1998). The *americana*- (99 % bs, 1 pp), *hominis*- (99 % bs, 1 pp) and *lunata*- (93 % bs, 0.99 pp) clades included species with mostly 4-celled, inconspicuously distoseptate conidia with the central cells usually darker than the end cells. In these clades the conidia are often curved at the third cell from base, and this cell is usually larger than the others (the only exceptional case is *C. brachyspora*, in which both central cells are more or less the same size (Sivanesan 1987)). This morphology, however, is also observed in *C. ischaemi*, which falls outside these three clades. The *americana*-clade included *C. americana* and *C. verruculosa*. These species appeared




Fig. 1 Bayesian consensus tree obtained from the combined ITS, LSU, *gpd* and RPB2 alignment of *Curvularia* and related genera. The scale bar represents the average number of substitutions per site. Bootstrap values $\geq 70\%$ and posterior probabilities ≥ 0.95 (in *italics*) are given near the internodes. The new species proposed in this study are shown in the coloured boxes. Ex-type and ex-neotype isolates for each species are indicated with a 'T' or 'NT', respectively, after the isolate number.

in Fig. 1 as two distinct species separated by relatively long branches. The *hominis*-clade included two new species, *C. hominis* and *C. muehlenbeckiae*. One isolate of the latter taxon, CBS 144.63, had been labelled '*C. lunata*' in the CBS collection, but in this study it proved to be phylogenetically quite distant from the ex-neotype strain of that species, CBS 730.96. The *lunata*-clade was formed by *C. aerea*, *C. brachyspora*, *C. carica-papayae*, *C. chlamydospora*, *C. lunata*, *C. prasadii* and *C. pseudolunata*. Accentuated septa can be observed in all members of this clade and elongate blackish stromata have been reported in *C. carica-papayae* and *C. aerea* (Mathur & Mathur 1959, Ellis 1966, 1971, Sivanesan 1987). This kind of stromata is also produced by old cultures of the ex-neotype strain of *C. lunata*, CBS 730.96 (unpubl. data). Isolates of *C. chlamydospora* and *C. pseudolunata* can produce aggregates of brown chlamydo-spores in culture (Fig. 3k and 6i). The *spicifera*-clade (98 % bs, 1 pp) was formed by *C. australiensis*, *C. ellisii*, *C. hawaiiensis*,

C. perotidis and *C. spicifera*. Members of this clade produce conspicuously distoseptate conidia that are straight in all species, except in *C. ellisii* which produces both straight and curved conidia (Sivanesan 1987). Three taxa of this clade are agents of opportunistic infections in humans, i.e. *C. australiensis*, *C. hawaiiensis* and *C. spicifera* (McGinnis et al. 1986, de Hoog et al. 2000). The *trifolii*-clade (95 % bs, 1 pp) included *C. akaii*, *C. heteropogonis*, *C. gladioli* and *C. trifolii*. These species produce 4-celled, usually curved, inconspicuously distoseptate conidia which, in contrast to those seen in the other clades discussed here, show a strongly protruding hilum (Sivanesan 1987, Boerema & Hamers 1989, Alcorn 1990). Two other species in our study produce conidia with a protruding hilum, i.e. *C. cymbopogonis* and *C. protuberata*. Their conidia, however, are 5-celled (Sivanesan 1987).

Not all *Curvularia* species were included in the six clades previously mentioned, and other well-supported lineages were




Fig. 2 *Curvularia americana* (a, b, d–j: CBS 136983; c, k: FMR 11674). a, b. Colonies on OA and PCA, respectively, at 25 °C after 7 d; c–i. conidiophores and conidia; j, k. microconidiation. — Scale bars: c–i = 10 µm.

observed. *Curvularia oryzae* and *C. tuberculata*, for example, appeared as sister taxa with 99 % bs and 1 pp. These species are morphologically very different, i.e. the conidia of *C. oryzae* are 3-distoseptate and smooth while those of *C. tuberculata* are 3–8-distoseptate and tuberculate at maturity (Sivanesan 1987). We preferred not to name morphologically heterogeneous lineages because future studies including more taxa might reveal more homogeneous groupings within such lineages.

Taxonomy

Curvularia americana Da Cunha, Madrid, Gené & Cano, sp. nov. — MycoBank MB806052; Fig. 2

Etymology. The name refers to the continent where this species was found.

Vegetative hyphae septate, branched, subhyaline to brown, smooth to asperulate, 1.5–4 µm wide, anastomosing. *Conidiophores* semi- to macronematous, mononematous, septate,

usually simple, slightly geniculate, subhyaline to dark brown, smooth to asperulate, with cell walls often thicker than those of the vegetative hyphae, 60–299 × 2–5 µm. *Conidiogenous cells* terminal or intercalary, polytretic, proliferating sympodially, subcylindrical to slightly swollen, 8–22 × 4–8 µm. *Conidia* 4(–5)-celled, straight to slightly curved, 13–28 × 7–15 µm, usually with the third cell unequally sided and larger than the others, second and third cells pale brown to brown, apical and basal cell subhyaline, apical cell smooth-walled, intermediate smooth (slightly verruculose under SEM), basal cell often verruculose; hilum non-protruding, flat, darkened and thickened, 1.5–3 µm wide. *Microconidiation* sometimes present, forming 1-celled, pale brown, globose conidia 5–6 µm wide. *Chlamydospores* not observed. *Sexual morph* not observed.

Culture characteristics — Colonies on OA and PCA attaining 62 and 69 mm diam, respectively, in 7 d at 25 °C, funiculose and greenish grey to dark green at the centre, effuse and greyish white towards the periphery, with a fimbriate margin; reverse olive to dark green.




Fig. 3 *Curvularia chlamydospora* (a–d, h–k: CBS 136984; e–g: FMR 11040). a, b. Colonies on OA and PCA, respectively, at 25 °C after 7 d; c–g, i, j. conidiophores and conidia; h. microconidiation; k. chlamydospore. — Scale bars: c, i, k = 20 µm; d–h = 10 µm; j = 5 µm.

Specimens examined. USA, Minnesota, culture from ankle (human), 2008, D.A. Sutton (holotype CBS H-21465, culture ex-type FMR 11551 = UTHSC 08-3414 = CBS 136983); California, culture from maxillary sinus (human), 2010, D.A. Sutton (FMR 11500 = UTHSC 10-1276); Ohio, culture from peritoneal dialysis fluid (human), 2008, D.A. Sutton (FMR 11691 = UTHSC 08-278); Oklahoma, culture from toe nail (human), 2009, D.A. Sutton (FMR 11005 = UTHSC 09-2907); Tennessee, culture from leg (human), 2008, D.A. Sutton (FMR 11674 = UTHSC 08-2697); Texas, culture from toe tissue (human), 2007, D.A. Sutton (FMR 11687 = UTHSC 07-2649); Texas, culture from bronchial wash (human), 2009, D.A. Sutton (FMR 11514 = UTHSC 09-2863); Utah, culture from nasal sinus (human), 2008, D.A. Sutton (FMR 11693 = UTHSC 08-84); Virginia, culture from bone marrow (human), 2009, D.A. Sutton (FMR 11515 = UTHSC 09-2806).

Notes — *Curvularia americana* is similar to *C. lunata* and *C. prasadii* in conidial morphology. However, the conidia of *C. lunata* are slightly narrower, up to 13 µm wide (Manamgoda et al. 2012) and, in contrast to *C. americana*, all septa in conidia of *C. prasadii* are accentuated and up to 2.4 µm wide (Mathur & Mathur 1959, Ellis 1966, 1971). The phylogenetic study placed *C. lunata* and *C. prasadii* in the *lunata*-clade, a lineage relatively distant from *C. americana*. The 4-locus tree indicated that *C. americana* is the sister taxon of *C. verruculosa*, but these species were separated by a considerable genetic distance (Fig. 1). The conidia of *C. verruculosa* are slightly larger (20–40 × 12–17 µm) than those of *C. americana* and show distinctly verruculose intermediate cells (Tandon & Bilgrami 1962, Ellis 1966, 1971, Sivanesan 1987).

Curvularia chlamydospora Madrid, Da Cunha, Gené & Guarro, *sp. nov.* — MycoBank MB806053; Fig. 3

Etymology. The name refers to the presence of chlamydospores.

Vegetative hyphae septate, branched, subhyaline to brown, smooth-walled, 1.5–4 µm wide, anastomosing. *Conidiophores* semi- to macronematous, mononematous, septate, usually simple, geniculate or bent at the apex, brown to dark brown, smooth to asperulate, 22–323 × 2–5 µm. *Conidiogenous cells* terminal or intercalary, polytretic, proliferating sympodially, subcylindrical to irregularly shaped, 7–18 × 5–10 µm. *Conidia* 4-celled, mostly slightly curved, 16–25 × 7–12 µm wide in the broadest part, smooth-walled (basal cell verruculose under SEM), usually with the central septum appearing slightly accentuated, the third cell from the base slightly larger and unequal sided, second and third cells darker than the others, brown to dark brown, end cells paler; hilum non-protruding, flat, darkened and thickened, 1.5–3 µm wide. *Chlamydospores* present, initially as intercalary chains but later forming clusters of swollen cells, 13–80 µm, smooth to verruculose and thick-walled. *Microconidiation* present, forming conidia 1–2-celled, pale brown, globose to subglobose, 4–6 µm diam. *Sexual morph* not observed.

Culture characteristics — Colonies on OA attaining 76 mm diam in 7 d at 25 °C, funiculose, greenish grey or dark green, margin fimbriate; reverse olive grey to dark green. Colonies on PCA attaining 68 mm diam at the same temperature and time of incubation, funiculose at the centre, effuse towards the periphery, dark green, with a fimbriate margin; reverse dark green.

Specimens examined. USA, Montana, culture from toe nail (human), 2007, D.A. Sutton (holotype CBS H-21466, culture ex-type FMR 11709 = UTHSC 07-2764 = CBS 136984); Nevada, culture from nasal sinus (human), 2008, D.A. Sutton (FMR 11040 = UTHSC 08-1283).

Notes — *Curvularia chlamydospora* is superficially similar to three species producing 4-celled conidia with an accentuated median septum, namely *C. brachyspora*, *C. eragrostidis* and *C. intermedia*. However, the third cell from base is usually larger and more pigmented than the second one in *C. chlamydo-*

spora, while in the three similar taxa both intermediate cells are rather equal in size and pigmentation. These species have not been reported to produce chlamydospores in culture and have wider conidia, i.e. 10–14 µm in *C. brachyspora*, 11–20 µm in *C. eragrostidis* and 13–20 µm in *C. intermedia* (Sivanesan 1987). *Curvularia eragrostidis* and *C. intermedia* reside in the *eragrostidis*-clade, while *C. chlamydospora* belongs to the *lunata*-clade. *Curvularia brachyspora* appeared as the sister taxon of *C. chlamydospora* but this relationship received poor statistical support (Fig. 1).

Curvularia hominis Da Cunha, Madrid, Gené & Cano, *sp. nov.* — MycoBank MB806054; Fig. 4

Etymology. The name refers to the origin of the isolates, all of which were isolated from clinical human specimens.

Vegetative hyphae septate, branched, subhyaline to brown, smooth to slightly asperulate 1.5–5 µm wide, anastomosing. *Conidiophores* semi- to macronematous, mononematous, septate, simple or branched, geniculate towards the apex, subhyaline to dark brown, smooth to asperulate, with cell walls often thicker than those of the vegetative hyphae, 55–325 × 2–5 µm wide. *Conidiogenous cells* terminal or intercalary, polytretic, proliferating sympodially, subcylindrical to irregularly shaped, 6–26 × 4–9 µm; conidiogenous loci usually somewhat thickened and darkened. *Conidia* 4–5-celled, slightly curved, 18–30 × 7–14 µm wide in the broadest part, with the third cell from the base often larger and unequal sided, intermediate cells usually verruculose and darker than the others, brown, end cells subhyaline to pale brown and smooth-walled; hilum non-protruding, flat, darkened and thickened, 1.5–3 µm wide. *Microconidiation* and *chlamydospores* were not observed. *Sexual morph* not observed.

Culture characteristics — Colonies on OA and PCA attaining 70–72 mm diam in 7 d at 25 °C, funiculose and dark green at the centre, floccose and olive to white towards the periphery, with a fimbriate margin; reverse olive to dark green.

Specimens examined. USA, Florida, culture from cornea (human), 2009, D.A. Sutton (holotype CBS H-21467, culture ex-type FMR 11539 = UTHSC 09-464 = CBS 136985); Arkansas, culture from nasal sinus (human), 2007, D.A. Sutton (FMR 11172 = UTHSC 07-3184); Louisiana, culture from eye (human), 2008, D.A. Sutton (FMR 11688 = UTHSC 08-849); Minnesota, culture from nail (human), 2007, D.A. Sutton (FMR 11698 = UTHSC 07-3581); Minnesota, culture from nasal sinus (human), 2009, D.A. Sutton (FMR 11527 = UTHSC 09-2197); Ohio, culture from nasal sinus (human), 2009, D.A. Sutton (FMR 11535 = UTHSC 09-1692); Texas, culture from nasal sinus (human), 2007, D.A. Sutton (FMR 11704 = UTHSC 07-3105); Texas, culture from nail (human), 2008, D.A. Sutton (FMR 11683 = UTHSC 08-1296); Texas, culture from bronchial wash (human), 2008, D.A. Sutton (FMR 11680 = UTHSC 08-2418); Texas, culture from bronchial wash (human), 2008, D.A. Sutton (FMR 11542 = UTHSC 08-3737); Texas, culture from foot (human), 2008, D.A. Sutton (FMR 11678 = UTHSC 08-2517); Texas, culture from nasopharynx (human), 2009, D.A. Sutton (FMR 11521 = UTHSC 09-2532); Texas, culture from tissue (human), 2009, D.A. Sutton (FMR 11509 = UTHSC 09-3403); Utah, culture from cornea (human), 2007, D.A. Sutton (FMR 11708 = UTHSC 07-2791).

Notes — Although all isolates of this fungus were obtained from humans, the species might also be common in the environment. *Curvularia hominis* resembles other species of the genus with 4-celled conidia and an asymmetrically swollen, dark third cell, such as *C. aerea*, *C. carica-papayae*, *C. lunata* and *C. prasadii*, but differs from them in producing conidia with verruculose intermediate cells (Fig. 4e–i). The latter four species are members of the *lunata*-clade, whereas *C. hominis* and *C. muehlenbeckiae* form a distinct lineage, the *hominis*-clade (Fig. 1).




Fig. 4 *Curvularia hominis* (CBS 136985). a, b. Colonies on OA and PCA, respectively, at 25 °C after 7 d; c, e–i. conidiophores and conidia; d. conidium showing two wall layers. — Scale bars: c = 20 µm; d–g, i = 10 µm; h = 5 µm.

Curvularia muehlenbeckiae Madrid, Da Cunha, Gené, Guarro & Crous, *sp. nov.* — MycoBank MB806055; Fig. 5

Etymology. The name refers to the substrate from which the ex-type strain was obtained, *Muehlenbeckia* sp.

Vegetative hyphae septate, branched, subhyaline to brown, smooth-walled, 1.5–5 µm wide, anastomosing. *Conidiophores* semi- to macronematous, mononematous, septate, simple to branched, straight or flexuous, geniculate towards the apex, subhyaline to dark brown, smooth to asperulate with cell walls often thicker than those of the vegetative hyphae, 21.5–398 × 2–5 µm with subnodulose and nodulose intercalary swellings up to 9.5 µm wide, swellings coinciding with conidiogenous loci. *Conidiogenous cells* integrated, terminal and intercalary, subcylindrical to irregularly shaped, mono- to polytretic, proliferating sympodially; intercalary conidiogenous cells 5–18 µm long, terminal conidiogenous cells 5–25 µm long. *Conidia* 4-celled, asymmetrical to more or less curved at the third cell from base, 17–26 × 8.5–12 µm, intermediate cells dark brown and usually verruculose, end cells paler and smooth-walled

or less ornamented than central cells. *Chlamydospores* and *microconidiation* not observed. *Sexual morph* not observed.

Culture characteristics — Colonies on OA attaining 76 mm diam in 7 d at 24 °C, cottony to funiculose, pale grey at the centre, dark olive towards the periphery, with a fimbriate margin; reverse olivaceous-black. Colonies on PCA attaining 40 mm diam at the same temperature and period of incubation, radiate, funiculose, dark olive with a slightly fimbriate margin; reverse concolorous with surface.

Specimens examined. INDIA, from *Muehlenbeckia* sp. leaf, 1962, K.S. Bilgrami (holotype CBS H-10451, culture ex-type CBS 144.63). — USA, Utah, culture from chest (human), 2008, D.A. Sutton (UTHSC 08-2905 = FMR 11671 = CBS 136986).

Notes — This species is the sister taxon of *C. hominis*, which has slightly larger conidia (18–30 × 7–14 µm) with a similar ornamentation consisting of small but conspicuous warts. Some *Curvularia* species outside the *hominis*-clade produce conidia ornamented with warts, e.g. *C. tuberculata*, *C. verruculosa* and *C. verruciformis*. The first two species produce larger conidia, i.e. 23–52 × 13–20 µm and 20–40 × 12–17 µm, respectively,




Fig. 5 *Curvularia muehlenbeckiae* (CBS 144.63). a, b. Colonies on OA after 7 d and on PCA after 5 d, respectively, at 25 °C; c–h. conidiophores and conidia. — Scale bars: c–g = 10 µm; h = 5 µm.

and the third one differs from members of the *hominis*-clade by having mostly 5-celled, more strongly ornamented conidia (Jain 1962, Agarwal & Sahni 1963, Ellis 1966, Sivanesan 1987). *Curvularia* species with warted conidia appear in different clades, suggesting that this kind of ornamentation evolved several times in *Curvularia*.

Curvularia pseudolunata Da Cunha, Madrid & Gené, *sp. nov.*
— MycoBank MB806056; Fig. 6

Etymology. The name refers to the morphological resemblance and phylogenetic closeness of this species to *Curvularia lunata*.

Vegetative hyphae septate, branched, subhyaline to brown, smooth-walled, 1.5–5 µm wide. *Conidiophores* macronema-

tous, mononematous, septate, unbranched, geniculate near the apex, brown, smooth-walled, 100–350 × 2–4.5 µm. *Conidiogenous cells* mostly terminal, polytretic, proliferating sympodially, subcylindrical, subglobose to irregularly shaped, 4.5–30 × 6–10 µm. *Conidia* 4-celled, mostly curved, 20–27 × 8–12 µm, with the third cell from base usually unequally sided, larger and darker than the others, brown, second and end cells subhyaline to pale brown, smooth-walled, basal cell often verruculose; hilum non-protruding, flat, darkened and thickened, 1.5–2.5 µm wide. *Chlamydospores* abundant, initially as intercalary chains, later forming clusters of swollen cells, up to 60 µm diam, smooth- and thick-walled. *Microconidiation* not observed. *Sexual morph* not observed.




Fig. 6 *Curvularia pseudolunata* (CBS 136987). a, b. Colonies on OA and PCA, respectively, at 25 °C after 7 d; c–h. conidiophores and conidia; i. chlamydo-spores. — Scale bars: c = 20 µm; d–i = 10 µm.

Culture characteristics — Colonies on OA attaining 71 mm diam in 7 d at 25 °C cottony to lanose, greenish grey, with a fimbriate margin; reverse dark green. Colonies on PCA attaining 78 mm diam at the same temperature and time of incubation, lanose at the centre, floccose towards the periphery, greyish green, with a fimbriate margin; reverse olive green.

Specimen examined. USA, California, culture from nasal sinus (human), 2009, *D.A. Sutton* (holotype CBS H-21468, cultures ex-type FMR 11529 = UTHSC 09-2092 = CBS 136987).

Notes — *Curvularia pseudolunata* is morphologically similar to *C. lunata* and these taxa grouped together in the 4-locus phylogeny (Fig. 1). However, the conidia of *C. lunata* are slightly larger (21–31 × 9–13 µm) and this species is separated from *C. pseudolunata* by a considerable genetic distance.

DISCUSSION

Traditionally, *Curvularia* and *Bipolaris* have been distinguished by conidial features, i.e. euseptate and typically curved at a swollen intermediate cell in *Curvularia*, but straight to slightly curved and distinctly distoseptate in *Bipolaris* (Kwon-Chung & Bennett 1992, de Hoog et al. 2000, Revankar & Sutton 2010). We agree with the view of authors like Alcorn (1983b), Sivanesan (1987) and Seifert et al. (2011) that both genera have distoseptate conidia. Phylogenetic studies (Berbee et al. 1999, Manamgoda et al. 2012) have demonstrated that species with conspicuously distoseptate conidia previously placed in *Bipolaris* actually belong in *Curvularia*, e.g. members of the *spicifera*-clade (Fig. 1). Furthermore, in the new species described herein, two wall layers were often evident in young conidia (Fig. 2f, 4d, 5f) and septa are already visible at this stage; however, in mature conidia the layers may appear so close to one another that the conidia may look euseptate under the light microscope (Fig. 2g, 4i, 5e). A recently described

pleosporalean genus, *Porocercospora*, the causal agent of bufalograss false-smut disease, also shows two-layered conidial cell walls, but mature conidia often seem to have both eu- and distosepta, depending on how closely together the two cell wall layers cohere near the septa. This genus is phylogenetically closely related to *Bipolaris* and *Curvularia* and is similar to them in having tretic conidiogenesis and darkly pigmented mycelium. However, *Porocercospora* has conidiophores without a geniculate rachis, conidiogenous cells with inconspicuous, non-darkened conidiogenous loci and long, obclavate to cylindro-obclavate conidia (Amaradasa et al. 2014).

Although *Bipolaris* and *Curvularia* cannot be distinguished based on the morphology of their conidial septa, other morphological features seem to be of diagnostic value. None of the species of *Bipolaris* s.str. included in this study (Fig. 1) and in previous works (Berbee et al. 1999, Manamgoda et al. 2012) has conidia curved at an intermediate swollen cell. As described by Berbee et al. (1999) for ‘*Cochliobolus* group 1’, the conidia of *Bipolaris* s.str. can show a gentle curve that continues along the whole length of the conidium. Conidia ornamented with small to coarse warts are produced by some *Curvularia* species, e.g., *C. tuberculata*, *C. verruciformis* and *C. verruculosa*, but this kind of ornamentation has not been reported in *Bipolaris* s.str. (Jain 1962, Tandon & Bilgrami 1962, Agarwal & Sahni 1963, Ellis 1966, Sivanesan 1987). Another helpful character is the morphology of the hilum. None of the species in *Bipolaris* s.str. has conidia with a strongly protruding hilum, but it is observed in several members of *Curvularia* s.str., such as *C. cymbopogonis* and *C. protuberata*, as well as all species in the *trifolii*-clade (Sivanesan 1987). A protruding hilum is also observed in a closely related genus, *Exserohilum*, which also includes clinically relevant and plant-pathogenic species (McGinnis et al. 1986, de Hoog et al. 2000). Members of this genus sometimes form curved conidia, but the hilum is different from those seen in *Curvularia* spp. In *Exserohilum*

the hilum appears as a protrusion of the cell wall that is not delimited by a septum and that often appears double-walled, with the outer wall forming an enveloping collar or 'hilum bubble' around it (Alcorn 1983b, 1988). In *Curvularia*, by contrast, when the hilum protrudes, it appears single-walled in light microscopy and is delimited by a septum (Nelson & Hodges 1965, Sivanesan 1987, Zhang et al. 2004). Conidial size might also be helpful to distinguish *Bipolaris* s.str. from *Curvularia* s.str. Among the species falling in the *Bipolaris* clade in Berbee et al. (1999) and Manamgoda et al. (2012), the longest conidia are those of *B. zeae*, up to 225 µm long (Sivanesan 1987). Conidia of *Curvularia* s.str. tend to be shorter. Among species in the *Curvularia* clade (Berbee et al. 1999, Manamgoda et al. 2012), the longest conidia are produced by *C. tripogonis* and are up to 130 µm long (Sivanesan 1987).

Boedijn (1933) divided *Curvularia* into three groups of species, i.e. groups Maculans, Lunata and Genuculata. The Maculans group was characterised by producing 4-celled, straight or somewhat asymmetrical conidia with the central cells darker and larger than the end cells. This group included *C. maculans* (currently considered a synonym of *C. eragrostidis*), *C. cesatii* (this species was transferred to the genus *Endophragmiella* as *E. cesatii* by Hughes in 1979), *C. intermedia* and *C. spicifera*, all of which were unable to produce stromata in culture. The Lunata group included species with 4-celled, more or less curved conidia in which one of the intermediate cells is enlarged and darker than the others. Some of its members were *C. lunata*, *C. ramosa* and *C. trifolii*. This group was reported to produce subcylindrical stromata in culture. The Genuculata group was proposed for species with 5-celled conidia which often produced stromata, such as *C. genuculata*, *C. affinis*, *C. fallax* and *C. falcata* (this species was synonymized with *C. senegalensis* by Sivanesan in 1987). In our phylogenetic study, three species of Boedijn's Maculans group were included, i.e. *C. eragrostidis*, *C. intermedia* and *C. spicifera*. The group is polyphyletic since only the former two species grouped together in the *eragrostidis*-clade, a lineage characterised by rather straight, inconspicuously distoseptate, 4-celled conidia. This lineage also included *C. graminicola* (Fig. 1). *Curvularia spicifera* clustered in a different clade with other species whose conidia show evident distosepta. No DNA sequences have as yet been analysed for *Endophragmiella cesatii*, which, as indicated previously, originally was considered a *Curvularia* species and a member of the Maculans group (Boedijn 1933). Its morphology clearly suggests a phylogenetically distant fungus (Hughes 1979, 1980). The genus *Endophragmiella* is considered a member of the *Lasiosphaeriaceae*, *Sordariales* by Seifert et al. (2011). Boedijn's Lunata group also proved to be polyphyletic since two of its members, *C. lunata* and *C. trifolii*, clustered in separate, relatively distant clades in Fig. 1. Two members of Boedijn's Genuculata group included in this study, *C. affinis* and *C. senegalensis*, formed a well-supported clade. CBS isolates of *C. genuculata* could not be clearly distinguished molecularly from isolates labelled *C. senegalensis* in a study by da Cunha et al. (2013); other authors also suggested that these taxa might be conspecific (Hosokawa et al. 2003, Sun et al. 2003). Unfortunately, no ex-type strains of these species are available and epitypification is necessary to clarify their taxonomy. Isolate CBS 155.34, labelled 'Syntype' of *C. fallax* has 4-celled conidia and clustered with morphologically similar isolates of *Curvularia* spp. in a preliminary phylogeny (not shown). CBS 155.34 is possibly mislabelled and therefore was excluded from the analysis.

The well-documented *Bipolaris* s.l. opportunists, i.e. *B. australiensis*, *B. hawaiiensis* and *B. spicifera* (McGinnis et al. 1986), were transferred to *Curvularia* (Manamgoda et al. 2012), suggesting that pathogenicity to vertebrates in this group of fungi

might be restricted to the latter genus (Manamgoda et al. 2012). Interestingly, however, a recent study by da Cunha et al. (2012) reported that two species of *Bipolaris* s.str., *B. cynodontis* and *B. setariae*, represented 8.6 % and 1 %, respectively, of a total of 104 clinical *Bipolaris* s.l. isolates from the USA. These species were isolated from various anatomical sites, including the eyes, legs, nasal sinuses, nails and lower respiratory tract. Their pathogenicity still needs to be demonstrated and no cases of human disease by these fungi have been published yet.

The ITS locus has been widely used in the identification of plant pathogenic and clinically relevant fungi and recently has been proposed as a universal barcode marker for these organisms (Iwen et al. 2002, Schoch et al. 2012). It has been used by some authors to identify isolates of *Curvularia* from clinical samples and plants (Fryen et al. 1999, Bagyalakshmi et al. 2008, Dyer et al. 2008, Chowdhary et al. 2011, Funnell-Harris et al. 2013). This marker, however, is not optimal for species identification since it provided little resolution for closely related *Curvularia* species in our study. Similar results were published by da Cunha et al. (2012), in which an ITS tree gave < 70 % bs for clades representing *C. spicifera* and *C. hawaiiensis*, two of the main clinically relevant members of the genus. Other authors have found limited species resolution in ITS phylogenies of other members of the *Pleosporales* (Pryor & Gilbertson 2000, de Hoog & Horr  2002, Pryor & Bigelow 2003, Park et al. 2008, Brun et al. 2013), indicating that additional genes need to be used for reliable species identification in this group of fungi. Protein-coding loci have been reported to be phylogenetically more informative than rDNA in *Ascomycota* (Schoch et al. 2009) and this is confirmed here in *Curvularia*. In our work, species discrimination improved with the *gpd* and *RPB2* loci, which revealed more than double the percentage of variable sites seen in ITS. These protein-coding loci are promising markers for future phylogenetic studies in *Curvularia* and related genera.

Acknowledgements We thank the technical staff, Arien van Iperen (cultures) and Janneke Bloem (DNA isolation, amplification and sequencing of some of the isolates studied) for their invaluable assistance. The study has been economically supported in part by the Spanish Ministry of 'Econom a y Competitividad', grant CGL 2011-27185.

REFERENCES

- Agarwal GP, Sahni VP. 1963. *Curvularia verruciformis* Agarwal & Sahni, a new fungus from Jabalpur (M.P.). *Current Science* 32: 276–277.
- Alcorn JL. 1983a. On the genera *Cochliobolus* and *Pseudocochliobolus*. *Mycotaxon* 16: 353–379.
- Alcorn JL. 1983b. Generic concepts in *Drechslera*, *Bipolaris* and *Exserohilum*. *Mycotaxon* 17: 1–86.
- Alcorn JL. 1988. The taxonomy of 'Helminthosporium' species. *Annual Review of Phytopathology* 26: 37–56.
- Alcorn JL. 1990. Additions to *Bipolaris*, *Cochliobolus* and *Curvularia*. *Mycotaxon* 39: 361–392.
- Alcorn JL. 1998. A new *Cochliobolus* species and its *Curvularia* anamorph. *Proceedings of the Royal Society of Queensland* 107: 1–4.
- Amaradasa BS, Madrid H, Groenewald JZ, Crous PW, Amundsen K. 2014. *Porocercospora seminalis* gen. et comb. nov. the causal organism of buf-falgrass false smut. *Mycologia* 106: 77–85.
- Bagyalakshmi R, Therese KL, Prasanna S, Madhavan HN. 2008. Newer emerging pathogens of ocular nonsporulating molds (NSM) identified by polymerase chain reaction (PCR)-based DNA sequence technique targeting internal transcribed spacer (ITS) region. *Current Eye Research* 33: 139–147.
- Barr ME. 1979. A classification of *Loculoascomycetes*. *Mycologia* 71: 935–957.
- Berbee ML, Pirseyedi M, Hubbard S. 1999. *Cochliobolus* phylogenetics and the origin of known, highly virulent pathogens inferred from ITS and glyceraldehyde-3-phosphate dehydrogenase gene sequences. *Mycologia* 91: 964–977.
- Boedijn KB. 1933.  ber einige phragmosporen Dematiaceen. *Bulletin du Jardin Botanique de Buitenzorg* 13: 120–134.

- Boerema GH, Hamers MEC. 1989. Check-list for scientific names of common parasitic fungi. Series 3b: Fungi on bulbs: Amaryllidaceae and Iridaceae. Netherlands Journal of Plant Pathology 95: 1–32 (Supplement 3).
- Brun S, Madrid H, Gerrits van den Ende B, Andersen B, Marinach-Patrice C, et al. 2013. Multilocus phylogeny and MALDI-TOF analysis of the plant pathogenic species *Alternaria dauci* and relatives. Fungal Biology 117: 32–40.
- Chowdhary A, Randhawa HS, Singh V, Khan ZU, Ahmad S, et al. 2011. *Bipolaris hawaiiensis* as etiologic agent of allergic bronchopulmonary mycosis: first case in a pediatric patient. Medical Mycology 49: 760–765.
- Cunha KC da, Sutton DA, Fothergill AW, Cano J, Gené J, et al. 2012. Diversity of *Bipolaris* species in clinical samples in the United States and their antifungal susceptibility profiles. Journal of Clinical Microbiology 50: 4061–4066.
- Cunha KC da, Sutton DA, Fothergill AW, Gené J, Cano J, et al. 2013. In vitro antifungal susceptibility and molecular identity of 99 clinical isolates of the opportunistic fungal genus *Curvularia*. Diagnostic Microbiology and Infectious Disease 76: 168–174.
- Domsch KH, Gams W, Anderson TH. 2007. Compendium of soil fungi. 2nd ed. IHW-Verlag, Germany.
- Dyer ZA, Wright RS, Rong IH, Jacobs A. 2008. Back pain associated with endobronchial mucus impaction due to *Bipolaris australiensis* colonization representing atypical allergic bronchopulmonary mycosis. Medical Mycology 46: 589–594.
- Ebright JR, Chandrasekar PH, Marks S, Fairfax MR, Aneziokoro A, McGinnis MR. 1999. Invasive sinusitis and cerebritis due to *Curvularia clavata* in an immunocompetent adult. Clinical Infectious Diseases 28: 687–689.
- Ellis MB. 1966. Dematiaceous Hyphomycetes VII: *Curvularia*, *Brachysporium*, etc. Mycological Papers 106: 1–57.
- Ellis MB. 1971. Dematiaceous Hyphomycetes. Commonwealth Mycological Institute, United Kingdom.
- Ellis MB. 1976. More dematiaceous Hyphomycetes. Commonwealth Mycological Institute, United Kingdom.
- Eriksson O. 1981. The families of bitunicate ascomycetes. Opera Botanica 60: 1–220.
- Faurel L, Schotter G. 1965. Champignons coprophiles du Tibesti. Revue Mycologique 30: 330–351.
- Fryen A, Mayser P, Glanz H, Füssle R, Breithaupt H, Hoog GS de. 1999. Allergic fungal sinusitis caused by *Bipolaris (Drechslera) hawaiiensis*. European Archives of Otorhinolaryngology 256: 330–334.
- Funnell-Harris DL, Prom LK, Pedersen JF. 2013. Isolation and characterization of the grain mold fungi *Cochliobolus* and *Alternaria* spp. from sorghum using semiselective media and DNA sequence analysis. Canadian Journal of Microbiology 59: 87–96.
- Hawksworth DL, Crous PW, Redhead SA, Reynolds DR, Samson RA, et al. 2011. The Amsterdam declaration on fungal nomenclature. IMA Fungus 2: 105–112.
- Hoog GS de, Guarro J, Gené J, Figueras MJ. 2000. Atlas of clinical fungi. 2nd ed. Centraalbureau voor Schimmelfcultures, Utrecht, The Netherlands.
- Hoog GS de, Horré R. 2002. Molecular taxonomy of the *Alternaria* and *Ulocladium* species from humans and their identification in the routine laboratory. Mycoses 45: 259–276.
- Hosokawa M, Tanaka C, Tsuda M. 2003. Conidium morphology of *Curvularia geniculata* and allied species. Mycoscience 44: 227–237.
- Hughes SJ. 1979. Relocation of species of *Endophragma* auct. with notes on relevant generic names. New Zealand Journal of Botany 17: 139–188.
- Hughes SJ. 1980. *Endophragma cerasii*. Fungi Canadenses 162: 1–2.
- Ismail Y, Johnson RH, Wells MV, Pusavat J, Douglas K, Arsura EL. 1993. Invasive sinusitis with intracranial extension caused by *Curvularia lunata*. Archives of Internal Medicine 153: 1604–1606.
- Iwen PC, Hinrichs SH, Rupp ME. 2002. Utilization of the internal transcribed spacer regions as molecular targets to detect and identify human pathogens. Medical Mycology 40: 87–109.
- Jain BL. 1962. Two new species of *Curvularia*. Transactions of the British Mycological Society 45: 539–544.
- Jiang YL, Zhang TY. 2007. Notes on soil dematiaceous hyphomycetes. Mycosystema 26: 17–21.
- Kamalam A, Ajithadass K, Sentamilselvi G, Thambiah AS. 1992. Paronychia and black discoloration of a thumb caused by *Curvularia lunata*. Mycopathologia 118: 83–84.
- Kwon-Chung KJ, Bennett JW. 1992. Medical Mycology. Lea & Febiger, USA.
- Lopes JO, Jobim NM. 1998. Dermatophytosis of the toe web caused by *Curvularia lunata*. Revista do Instituto de Medicina Tropical de São Paulo 40: 327–328.
- Luttrell ES. 1963. Taxonomic criteria in *Helminthosporium*. Mycologia 55: 643–674.
- Manamgoda DS, Cai L, McKenzie EHC, Crous PW, Madrid H, et al. 2012. A phylogenetic and taxonomic re-evaluation of the *Bipolaris-Cochliobolus-Curvularia* complex. Fungal Diversity 56: 131–144.
- Mathur RL, Mathur BL. 1959. A new species of *Curvularia* from the leaves of *Jasminum sambac*. Current Science 28: 448–449.
- McGinnis MR, Rinaldi MG, Winn RE. 1986. Emerging agents of phaeohyphomycosis: pathogenic species of *Bipolaris* and *Exserohilum*. Journal of Clinical Microbiology 24: 250–259.
- Nelson RR, Hodges CS. 1965. A new species of *Curvularia* with a protuberant conidial hilum. Mycologia 57: 822–825.
- Nylander JAA. 2004. MrModeltest v. 2. Software distributed by the author. Evolutionary Biology Centre, Uppsala University, Sweden.
- O'Donnell K, Sarver BAJ, Brandt M, Chang DC, Noble-Wang J, et al. 2007. Phylogenetic diversity and microsphere array-based genotyping of human pathogenic fusaria including isolates from the 2005–06 multistate contact lens-associated U.S. keratitis outbreaks. Journal of Clinical Microbiology 45: 2235–2248.
- Olivier C, Berbee ML, Shoemaker RA, Loria R. 2000. Molecular phylogenetic support from ribosomal DNA sequences for origin of *Helminthosporium* from *Leptosphaeria*-like loculoascomycete ancestors. Mycologia 92: 736–746.
- Park MS, Romanosky CE, Pryor BM. 2008. A reexamination of the phylogenetic relationship between the causal agents of carrot black rot, *Alternaria radicina* and *A. carotiincultae*. Mycologia 100: 511–527.
- Pryor BM, Bigelow DM. 2003. Molecular characterization of *Embellisia* and *Nimbya* species and their relationship to *Alternaria*, *Ulocladium* and *Stemphylium*. Mycologia 95: 1141–1154.
- Pryor BM, Gilbertson RL. 2000. Molecular phylogenetic relationships amongst *Alternaria* species and related fungi based upon analysis of nuclear ITS and mt SSU rDNA sequences. Mycological Research 104: 1312–1321.
- Revankar SG, Sutton DA. 2010. Melanized fungi in human disease. Clinical Microbiology Reviews 23: 884–928.
- Ronquist S, Huelsenbeck JP. 2003. MrBayes 3: Bayesian phylogenetic inference under mixed models. Bioinformatics 19: 1572–1574.
- Schoch CL, Seifert KA, Huhndorf S, Robert V, Spouge JL, et al. 2012. Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for fungi. Proceedings of the National Academy of Sciences USA 109: 6241–6246.
- Schoch CL, Sung GH, López-Giráldez F, Townsend JP, Miadlikowska J, et al. 2009. The Ascomycota tree of life: a phylum-wide phylogeny clarifies the origin and evolution of fundamental reproductive and ecological traits. Systematic Biology 58: 224–239.
- Seifert K, Morgan-Jones G, Gams W, Kendrick B. 2011. The genera of hyphomycetes. CBS Biodiversity Series 9. CBS-KNAW Fungal Biodiversity Centre, Utrecht, The Netherlands.
- Sivanesan A. 1984. The bitunicate ascomycetes and their anamorphs. Strauss & Kramer, Liechtenstein.
- Sivanesan A. 1987. Graminicolous species of *Bipolaris*, *Curvularia*, *Drechslera*, *Exserohilum* and their teleomorphs. Mycological Papers 158: 154–185.
- Sun G, Oide S, Tanaka E, Shimizu K, Tanaka C, Tsuda M. 2003. Species separation in *Curvularia 'geniculata'* group inferred from Brn1 gene sequences. Mycoscience 44: 239–244.
- Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. 2011. MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. Molecular Biology and Evolution 28: 2731–2739.
- Tandon RN, Bilgrami KS. 1962. A new pathogenic species of genus *Curvularia*. Current Science 31: 254.
- Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG. 1997. The ClustalX Windows interface: flexible strategies for multiple alignment aided by quality analysis tools. Nucleic Acids Research 25: 4876–4882.
- Tsuda M, Ueyama A. 1981. *Pseudocochliobolus australiensis*, the ascigerous state of *Bipolaris australiensis*. Mycologia 73: 88–96.
- Tsuda M, Ueyama A, Nishihara N, 1977. *Pseudocochliobolus nisikadoi*, the perfect stage of *Helminthosporium coicis*. Mycologia 69: 1109–1120.
- Vilgalys R, Hester M. 1990. Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several species of *Cryptococcus*. Journal of Bacteriology 172: 4238–4246.
- White TJ, Bruns TD, Lee S, Taylor J. 1990. Amplification and direct sequencing of fungal ribosomal genes for phylogenetics. In: Gelfand M, Sninsky JI, White TJ (eds), PCR protocols: a guide to methods and applications: 315–322. Academic Press, USA.
- Yanagihara M, Kawasaki M, Ishizaki H, Anzawa K, Udagawa S, et al. 2010. Tiny keratotic brown lesions of the interdigital web between the toes of a healthy man caused by *Curvularia* species infection and a review of cutaneous *Curvularia* infections. Mycoscience 51: 224–233.
- Zhang M, Zhang TY, Wu YM. 2004. A new name and a new variety in *Curvularia*. Mycosystema 23: 177–178.
- Zhang Y, Crous PW, Schoch CL, Hyde KD. 2012. Pleosporales. Fungal Diversity 53: 1–221.
- Zhang Y, Schoch CL, Crous PW, Gruyter J de, Woudenberg JHC, et al. 2009. Multi-locus phylogeny of Pleosporales: a taxonomic, ecological and evolutionary re-evaluation. Studies in Mycology 64: 85–102.