

A new gall crab species (*Brachyura*, *Cryptochiridae*) associated with the free-living coral *Trachyphyllia geoffroyi* (*Scleractinia*, *Merulinidae*)

Sancia E.T. van der Meij¹

¹ Department of Marine Zoology, Naturalis Biodiversity Center, Darwinweg 2, 2333 CR Leiden, The Netherlands

Corresponding author: Sancia E.T. van der Meij (Sancia.vanderMeij@naturalis.nl)

Academic editor: S. De Grave | Received 21 January 2015 | Accepted 31 March 2015 | Published 27 April 2015

<http://zoobank.org/E6B958CD-5FED-45F2-BDD8-90BBA1D32087>

Citation: van der Meij SET (2015) A new gall crab species (*Brachyura*, *Cryptochiridae*) associated with the free-living coral *Trachyphyllia geoffroyi* (*Scleractinia*, *Merulinidae*). *ZooKeys* 500: 61–72. doi: 10.3897/zookeys.500.9244

Abstract

A new species of gall crab is described from the free-living stony coral *Trachyphyllia geoffroyi*. Specimens were collected during field work in Lembeh Strait (Indonesia) and off Kudat (Malaysian Borneo). This new species, here named *Lithoscaptus semperi* sp. n., is the ninth species assigned to the genus. It can be separated from its congeners by not having the internal orbital angle extending beyond the external orbital angle, and by the stout female P2 merus with prominent distomesial projection. In addition, the carapace surface appears smooth, despite having small tubercles on the anterior half, and is without noticeable spines, other than those on the frontal margin. The distinctive carapace pattern in life is a diagnostic character in male specimens.

Keywords

Cospeciation, host specificity, Indonesia, Malaysia, Thoracotremata

Introduction

During field work in Indonesia and Malaysia an undescribed gall crab species was encountered living in dwellings in free-living *Trachyphyllia geoffroyi* (Audouin, 1826) corals. This scleractinian species is usually found on soft substrate of reef bases near coral reefs, where it can occur in large numbers (Fisk 1983, Best and Hoeksema 1987). The polyps of *T. geoffroyi* are fleshy and a large mantle can extend beyond the perimeter of the skeleton.

Trachyphyllia geoffroyi was classified in its own family, Trachyphylliidae Verrill, 1901, but this taxon was recently synonymised with Merulinidae Verrill, 1865 (Huang et al. 2014). The sister genera of *Trachyphyllia* Milne Edwards & Haime, 1849 are *Coelestrea* Verrill, 1866 and *Dipsastraea* de Blainville, 1830, which include coral species that formerly belonged to *Goniastrea* Milne Edwards & Haime, 1848 and *Favia* Milne Edwards, 1857. Corals belonging to these genera are host to cryptochirids of the genus *Lithoscaptus* A. Milne-Edwards, 1862 (Fize and Serène 1957, Kropp 1990).

Semper (1881) mentioned gall crabs associated with Indo-Pacific and Atlantic “*Trachyphyllia*”, but no formally described gall crab has been recorded living in association with *T. geoffroyi*. This new gall crab species, here named *Lithoscaptus semperi* sp. n., is the ninth assigned to the genus.

Methods

Gall crabs were collected in Indonesia (Lembah Strait, N Sulawesi – February 2012) and Malaysia (off Kudat, N Borneo – September 2012). Corals were searched for gall crabs, taken to the field laboratory and subsequently split with hammer and chisel. The crabs were preserved in 80% ethanol, after being photographed with a digital SLR camera equipped with a macro lens to register colour patterns. All crab specimens are deposited in the Crustacea collection of Naturalis Biodiversity Center in Leiden, the Netherlands (formerly Rijksmuseum van Natuurlijke Historie, collection coded as RMNH.Crus.D).

Drawings were made with a stereomicroscope with camera lucida. Carapace lengths and widths were measured to the nearest 0.1 mm using an eyepiece micrometre, with the crabs positioned on a level surface. Abbreviations used: CL, carapace length; CW, carapace width (at widest point); MXP3, third maxilliped; ovig., ovigerous; P, pereopod; G, male gonopod. Carapace measurements are given as CL × CW, in mm.

Taxonomy

Cryptochiridae Paul'son, 1875

Lithoscaptus A. Milne-Edwards, 1862

Lithoscaptus semperi sp. n.

<http://zoobank.org/65F0D837-961A-42B7-8F9E-2C806DD54238>

Figs 1–3

Type locality. Tigabu Isl. (06°53'51"N, 117°27'36"E), Kudat, Sabah (N Borneo), Malaysia.

Coral host holotype. *Trachyphyllia geoffroyi* (Audouin, 1826).

DNA barcoding. A COI sequence (partially, Folmer et al. 1994) of paratype RMNH.Crus.D.54331 has been deposited in GenBank under accession number KP688583.

Type material. Holotype. RMNH.Crus.D.56962, ovig. female, 6.4×4.6 . **Allotype** (with holotype), male, 3.6×2.5 . Collected by the author from 13 m depth on 8 September 2012. **Paratype.** RMNH.Crus.D.54331, Lubani Rock, Kudat, Sabah (N Borneo), Malaysia ($06^{\circ}53'45.0''\text{N}$, $117^{\circ}23'15.8''\text{E}$), 10–15 m, 07.ix.2012, 1 ovig. female, 6.2×4.7 , leg. SET van der Meij.

Material examined. Indonesia: RMNH.Crus.D.56957, Aer Perang, Lembah Strait ($01^{\circ}28'25''\text{N}$, $125^{\circ}14'02''\text{E}$), ca. 10 m, 02.ii.2012, 1 female, leg. BT Reijnen; RMNH.Crus.D.56958, Tanjung Labuhankompeni, Lembah Strait ($01^{\circ}25'55''\text{N}$, $125^{\circ}11'10''\text{E}$), 28 m, 04.ii.2012, 1 female, leg. BW Hoeksema; RMNH.Crus.D.56959, Kelapadua, Lembah Strait ($01^{\circ}26'19''\text{N}$, $125^{\circ}12'49''\text{E}$), 20 m, 09.ii.2012, 2 juvenile males, leg. BW Hoeksema; RMNH.Crus.D.54250, Tanjung Nanas I, Lembah Strait ($01^{\circ}27'39''\text{N}$, $125^{\circ}13'35''\text{E}$), 25–30 m, 17.ii.201, 1 ovig. female, 1 female, leg. BW Hoeksema; **Malaysia:** RMNH.Crus.D.54259, Lubani Rock, Kudat ($06^{\circ}53'45''\text{N}$, $117^{\circ}23'15''\text{E}$), 10–15 m, 07.ix.2012, 1 ovig. female (slightly damaged), leg. BW Hoeksema; RMNH.Crus.D.54280, Lubani Rock, Kudat ($06^{\circ}53'45''\text{N}$, $117^{\circ}23'15''\text{E}$), 10–15 m, 07.ix.2012, 1 ovig. female, 1 male, leg. BW Hoeksema; RMNH.Crus.D.56960, Lubani Rock, Kudat ($06^{\circ}53'45''\text{N}$, $117^{\circ}23'15''\text{E}$), 10–15 m, 07.ix.2012, 1 male, leg. SET van der Meij; RMNH.Crus.D.56961, Lubani Rock, Kudat ($06^{\circ}53'45''\text{N}$, $117^{\circ}23'15''\text{E}$), 10–15 m, 07.ix.2012, 1 ovig. female, leg. SET van der Meij; RMNH.Crus.D.54312, Tigabu Is., Kudat ($06^{\circ}53'51''\text{N}$, $117^{\circ}27'36''\text{E}$), 9 m, 08.ix.2012, 1 ovig. female (damaged), 1 male, leg. SET van der Meij; RMNH.Crus.D.56963, Fairway Shoal, Kudat ($07^{\circ}07'06''\text{N}$, $117^{\circ}30'42''\text{E}$), 12 m, 10.ix.2012, 1 male, leg. BT Reijnen; RMNH.Crus.D.56964, Belaruan, Kudat ($07^{\circ}01'50''\text{N}$, $117^{\circ}00'41''\text{E}$), ca. 15m, 20.ix.2012, 1 male, leg. BW Hoeksema; RMNH.Crus.D.54258, Tajau, Kudat ($06^{\circ}59'36''\text{N}$, $116^{\circ}50'27''\text{E}$), 21 m, 25.ix.2012, 1 female, 1 male, leg. BW Hoeksema. All material was collected from the scleractinian coral *Trachyphyllia geoffroyi*.

Description of female holotype. Carapace (Fig. 1A) rectangular, longer than broad, CL 1.4 times longer than CW; widest near midlength, dorsal surface in lateral view strongly convex in both directions, deflected anteriorly (Fig. 1B); anterior half of carapace with small, sharp tubercles, posterior half smooth with few, rounded granules, cardiointestinal region slightly inflated. Frontal margin armed with small anteriorly directed spines. Frontal margin on ventral side features few, small tubercles. Pterygostomial region fused to carapace.

Eyestalk exposed dorsally, slightly granular, small spines on mesial margin. Cornea anterolateral. Lateral margin of stalk at same level as anterolateral angle; distal margin with small spines (Fig. 1A, C). Distal segment of antennules with protruding article, visible from ventral side (Fig. 1C, D).

Antennular peduncle dorsal surface with small, sharp tubercles, slightly inflated distomesially; apex extending beyond tip of eyestalk; spines on mesial margin larger than those on distal margin. Ventral surface smooth, slightly tapering anteriorly in ventral view (Fig. 1C).

MXP3 (Fig. 1E) exopod rectangular; ischium subtriangular, smooth, mesial and distal margins straight, anteromesial lobe with few simple setae; merus with distolateral




Figure 1. Ovigerous female holotype (6.4×4.6) of *Lithoscaptus semperi* sp. n. (RMNH.Crus.D.56962) **A** habitus, dorsal view **B** carapace, lateral view **C** anterior margin of carapace, ventral view **D** close-up of antennule **E** MXP3 **F** left P1 (cheliped) **G** left P2 **H** left P3 **I** left P4 **J** left P5. Scale bars 1 mm; A–B, D–E, F–J share scale bars.

projection, simple setae; distal portion of carpus with short, simple setae, dactylus with bundle of setae.

P1 (chelipeds, Fig. 1F) slender; carpus length twice height, scattered small tubercles on dorsal surface, simple setae; propodus length twice height, somewhat granu-

lated, few, scattered setae, fingers slender, mesial surface of fingers smooth, cutting edge entire, tips of fingers crossing.

P2 (Fig. 1G) longer, coarser than P1; ischium without setae; merus stout, plump, smooth with few, small rounded tubercles on distal half of dorsal surface, simple setae on lateral surface, numerous plumose setae on dorsal surface; joint between merus, carpus not extending more than at right angle; carpus smooth with small rounded tubercles on dorsal surface, simple setae on dorsal surface; propodus slightly shorter than carpus, surface smooth with small rounded tubercles on dorsal surface, simple setae on lateral and dorsal surface; dactylus half-length of propodus, smooth, sharp, curved ventrally.

P3 (Fig. 1H) ischium without setae; merus length 1.5 times height, rounded, few rounded tubercles on distal half of dorsal surface, simple setae along dorsal, lateral surface; joint between merus, carpus not extending more than at right angle; carpus length 2.5 times height, rounded tubercles on dorsal surface, simple setae on lateral and dorsal surface; propodus length twice height, rounded tubercles on dorsal surface, scattered simple setae; dactylus similar length as propodus, smooth, sharp, slightly curved ventrally.

P4 (Fig. 1I) similar to P3, less coarse; ischium without setae; merus length 1.5 times height, small rounded tubercles close to joint with carpus, carpus length 2.5 times height, rounded tubercles on distal half of dorsal surface, scattered simple setae; propodus half-length carpus, rounded tubercles on distal half of dorsal surface, few scattered simple setae; dactylus similar length as propodus, smooth, sharp, straight.

P5 (Fig. 1J) ischium without setae; merus, carpus, propodus, dactylus all of equal length, all with short simple setae; carpus, propodus slender compared to merus; dactylus smooth, sharp, slightly curved ventrally.

P3, P4 decreasing in size from P2.

Abdomen enlarged, lateral margins fringed with setae (Fig. 1A, B).

Gonopore (vulva); reniform, size half the height of sternite 6.

Description of male allotype. Carapace (Fig. 2A) subrectangular to trapezoid, CL 1.5 times longer than CW, widest near anterior half, convex in lateral view, deflected anteriorly, with broad W-shaped depression (Fig. 2A, B). Anterior half of carapace and carapace margins with small spines, posterior half of carapace smooth.

Ocular peduncles with small spines on distal margin, cornea elliptical, longer than broad; antennal article extending beyond eyestalk, with spines along margins (Fig. 2C). Antennule slender compared to holotype, distal segment of antennules with protruding article (Fig. 2D).

MXP3 (Fig. 2E) exopod rectangular; ischium smooth, triangular, few scattered simple setae on distal and lateral margins, merus with distolateral projection, simple setae; propodus, dactylus of similar length, latter with bundle of short setae.

P1 (chelipeds, Fig. 2G) stout; merus length twice height, smooth; carpus with rounded and conical tubercles, simple setae on dorsal surface; propodus stout, with conical tubercles, simple setae on dorsal surface; fingers slender, mesial surfaces of dactyl slightly gaping, tips of fingers crossing.




Figure 2. Male allotype (3.6 × 2.5) of *Lithoscaptus semperi* sp. n. (RMNH.Crus.D.56962) **A** habitus, dorsal view **B** carapace, lateral view **C** anterior margin of carapace, ventral view **D** close-up of antennule **E** MXP3 **F** thoracic sternites **G** right P1 (cheliped) **H** right P2 **I** right P3 **J** right P4 **K** right P5. Scale bar 1 mm; A–C, D–E, F–K share scale bars.

P2 (Fig. 2H) ischium without setae; merus relatively stout, smooth, length twice height, simple short setae on lateral and dorsal surface; carpus, propodus of similar length; carpus with few rounded tubercles and setae on dorsal surface; propodus smooth except for rounded tubercles on dorsal surface, few setae on lateral, dorsal surface, dactylus smooth, sharp, curved ventrally.

P3 and P4 (Fig. 2I, J) similar to P2, somewhat smaller; ischium without setae; merus smooth, simple short setae on lateral and dorsal surface; carpus, propodus of

same length, few rounded tubercles and setae on dorsal surface; dactylus smooth, sharp, curved ventrally.

P5 (Fig. 2K) ischium with few setae; merus, carpus, propodus smooth, with simple short setae on dorsal and lateral surface; dactylus smooth, sharp, curved.

P3, P4 decreasing in size from P2.

Abdomen teardrop-shaped, widest at 4th somite; telson slightly pointed with few simple setae (Fig. 2F).

Gonopod 1 almost straight, tapering, apex sharply pointed. Distal margin with 2-3 non-plumose short simple setae, medial margin without setae (examined in RMNH. Crus.D.56964).

Colour. Female (Fig. 3A–B): Overall off-white. Pereiopods opaque, carpus, dactylus P1 and P2 translucent violet, sometimes with a pale orange line. Eyes with wide longitudinal brownish-red lines. Male (Fig. 3C–D): Carapace opaque with an off-white distinctive pattern over the whole carapace surface. Pereiopods opaque, P1 carpus, dactylus translucent violet, sometimes with a pale orange line. Eyes brown-red. In juvenile males (Fig. 3E), the carapace pattern is pale orange, pereiopods off-white.

Placement in genus. The placement of *Lithoscaptus semperi* sp. n. in the genus *Lithoscaptus* is somewhat tentative. The first (partial) molecular reconstruction of relationships within the Cryptochiridae shows that the genus *Lithoscaptus* is paraphyletic (van der Meij and Reijnen 2014). However, following the diagnosis of *Lithoscaptus* by Kropp (1990), the new species best fits the genus, except for the absence of a proximal tooth on the cutting edge of P1 dactylus and the presence of a distomesial projection of P2 merus in females. Kropp (1994) noted that his new species, *L. prionotus*, had the pterygostomial region not fused to the carapace, unlike other species in the genus. It is likely that the characters defining the genus need to be redefined, or that certain species need to be moved to a new genus.

Comparisons. Eight species of *Lithoscaptus* are currently recognised (Ng et al. 2008: 212, Davie 2015). *Lithoscaptus semperi* sp. n. can be distinguished from *L. nami* (Fize & Serène, 1957), *L. tri* (Fize & Serène, 1956) and *L. pardalotus* Kropp, 1995 by not having the internal orbital angle extending beyond the external orbital angle. The new species can be separated from *L. grandis* (Takeda & Tamura, 1983), *L. paradoxus* A. Milne-Edwards, 1862 and *L. prionotus* Kropp, 1994 by the smooth appearance of surface of the carapace, despite the small tubercles on the anterior half of the carapace, and the lack of noticeable spines other than the small spines on the frontal carapace margin. *Lithoscaptus pacificus* (Edmonson, 1933) and *L. helleri* (Fize & Serène, 1957) lack the stout merus with prominent distomesial projection of P2 (female specimens). The off-white carapace colour and translucent violet colour on P1 and P2 in females, and the distinctive carapace pattern in males differs from patterns found on other *Lithoscaptus* species.

Distribution. The known distribution of *L. semperi* sp. n. includes northern Borneo and North Sulawesi. Specimens were collected at water depths between 9 and approximately 30 meters. Its host *Trachyphyllia geoffroyi* was described from the Gulf of Suez (Egypt), but this species has a wide distribution that includes the Red Sea, East




Figure 3. Colour in life of *Lithoscaptus semperi* sp. n. **A–B** non-ovigerous female (4.5 × 3.2; RMNH. Crus.D.54258) dorsal view and ventral view **C–D** male (2.5 × 1.9; RMNH. Crus.D.54258) dorsal view and ventral view **E** juvenile male (2.0 × 1.6; RMNH. Crus.D.56959) dorsal view **F** in-situ photograph of dwellings (left male, right female) of *L. semperi* sp. n. in *Trachyphyllia geoffroyi* on Lubani Rock reef, Kudat (Malaysia). Photos by BT Reijnen/SET van der Meij.

Africa, Seychelles, Maldives, Nicobar Isls., 'East Indies', China Sea, Philippines, Japan, Australia and New Caledonia (Scheer and Pillai 1983). Based on the widespread distribution of *T. geoffroyi*, a wider distribution range than the two presently recorded locations is expected for *L. semperi* sp. n.

Coral host. *Lithoscaptus semperi* sp. n. is so far strictly associated with *T. geoffroyi* (Fig. 3F). It is the first record of associated fauna for this coral host. Colonies of *T. geoffroyi* are free-living, have flabello-meandroid colony shapes and fleshy polyps. Cryptochirids have previously been recorded to inhabit free-living corals; crabs of the genus *Fungicola* are associated with free-living - and attached - mushroom corals (Fungiidae), whereas *Troglocarcinus corallicola* is associated with a wide range of Atlantic corals, including the free-living coral *Manicina areolata* (Mussidae) (Fize and Serène 1957, van der Meij 2014, 2015).

Remarks. Fize and Serène (1957: p. 163) report on *Cryptochirus coralliodytes* from *Trachyphyllia* based on a record of Semper (1881: p. 221) who writes: "*I found them [C. coralliodytes] in the Philippine Archipelago in cavities in Goniastrea Bournoni [= Goniastrea retiformis (de Lamarck, 1816)], in an undetermined true Astræa, which was unfortunately lost, also in an undescribed Trachyphyllia; finally I received a new form through A. Agassiz from the West Indian seas, which may perhaps form a distinct genus, though it is very nearly allied to the first. It also lives in a Trachyphyllia.*" The coral genus *Trachyphyllia* is described from the Red Sea and has a widespread Indo-Pacific distribution; however, it does not occur in the Atlantic Ocean. The most similar Atlantic species would be *Manicina areolata* (Linnaeus, 1758). Furthermore, on p. 453 (note 103 belonging to p. 221) Semper writes: "*This crab, living in Trachyphyllia, a West Indian coral, is extremely like Cryptochirus, and perhaps belongs to the same genus; this can only be determined by future and more exact examination. But the 'cave dwelling' of this West Indian crab is perfectly unlike that of the Eastern species, which is found from the Red Sea as far as the Pacific Ocean; it is not cylindrical, but has one side quite flat, so that its transverse section is almost exactly a half-circle; the underside of the crab rests against the flat side of the cavity.*" The gall crab *Troglocarcinus corallicola* Verrill, 1908 has been recorded from a wide range of hosts, including *M. areolata* (Kropp and Manning 1987, van der Meij 2014). As mentioned by Semper (1881), the dwelling of *T. corallicola* in *M. areolata* is shaped like a half-circle (see e.g. Van der Meij 2014: Fig. 1B); therefore, it seems plausible that Semper was referring to the coral *M. areolata* when he discussed a West Indian *Trachyphyllia*. Alternatively, Semper could have been referring to the Atlantic genus *Colpophyllia* because Milne Edwards and Haime (1849), who established *Trachyphyllia*, compared their new genus with *Colpophyllia* (see Huang et al. [2014] for a discussion on the genus *Trachyphyllia*). Like *M. areolata*, *Colpophyllia natans* (Houttuyn, 1772) also hosts *T. corallicola* (see van der Meij 2014). It remains unclear whether Semper found gall crabs in Indo-Pacific corals currently recognized as *Trachyphyllia geoffroyi*. Semper is not known to have formally described any gall crab species (Ng et al. 2008).

Etymology. Named after the German naturalist Carl Gottfried Semper (1832–1893), who was the first to mention gall crabs occurring in *Trachyphyllia*.

Acknowledgements

The Tun Mustapha Park Expedition (TMPE) 2012 was jointly organized by WWF-Malaysia, Universiti Malaysia Sabah (UMS), Sabah Parks and Naturalis Biodiversity Center, The Netherlands. TMPE was funded by the Ministry of Science, Technology and Innovation (MOSTI) and USAID Coral Triangle Support Partnership (CTSP). The research permits were granted by the Economic Planning Unit, Prime Minister's Department Malaysia and Sabah Biodiversity Centre. Field work in Lembeh Strait in 2012 took place during a Marine Biodiversity Workshop based at the Bitung Field Station (LIPI), co-organized by Yosephine Tuti of the Research Centre of Oceanography of the Indonesian Institute of Sciences (PPO-LIPI) in Jakarta, Markus Lasut of Universitas Sam Ratulangi in Manado, North Sulawesi (Indonesia), and Bert Hoeksema of Naturalis Biodiversity Center. Funding for the fieldwork in Indonesia was provided by the LB Holthuisfonds (Naturalis), and the Schure-Beijerinck-Poppingfonds (KNAW). Bastian Reijnen and Bert Hoeksema (Naturalis) collected material used in this study. Erik-Jan Bosch (Naturalis) made the beautiful line drawings. The COI barcode was produced as part of the Naturalis Barcoding Project. I thank Roy Kropp and Peter Ng for their constructive comments on an earlier version of the manuscript.

References

- Audouin JV (1826) Explication sommaire des planches de polypes de l'Égypte et de la Syrie, publiées par Jules-César Savigny. In: Savigny JC (Ed.) Description de l'Égypte, ou recueil des observations et des recherches qui ont été faites en Égypte pendant l'expédition de l'armée française... Histoire naturelle Vol. Tome 1, 4 partie: 339. Imprimerie Impériale, Paris.
- Best MB, Hoeksema BW (1987) New observations on scleractinian corals from Indonesia: 1. Free-living species belonging to the Faviina. Zoologische Mededelingen, Leiden 61: 387–403.
- Blainville HM de (1830) Zoophytes. In: Levrault FG (Ed.) Dictionnaire des sciences naturelles, dans lequel on traite méthodiquement des différens êtres de la nature, considérés soit en eux-mêmes, d'après l'état actuel de nos connoissances, soit relativement à l'utilité qu'en peuvent retirer la médecine, l'agriculture, le commerce et les arts. Tome 60. Le Normat, Paris.
- Davie P (2015) *Lithoscaptus*. Accessed through: World Register of Marine Species at <http://marinespecies.org/aphia.php?p=taxdetails&id=439231> [visited 26 March 2015]
- Edmondson CH (1933) *Cryptochirus* of the central Pacific. Bernice P. Bishop Museum Occasional Papers 10(5): 1–23.
- Fisk DA (1983) Free-living corals: distributions according to plant cover, sediments, hydrodynamics, depth and biological factors. Marine Biology 74: 287–294. doi: 10.1007/BF00403453
- Fize A, Serène R (1956) Note préliminaire sur quatre espèces nouvelles d'Hapalocarcinidés avec quelques remarques au sujet du *Cryptochirus rugosus* Edmonson. Bulletin de la Société Zoologique de France 80(5–6): 379–382.

- Fize A, Serène R (1957) Les Hapalocarcinides du Viet-Nam. Mémoires de l'Institut Océanographique de Nhatrang 10: 1–202, pls. 1–18.
- Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R (1994) DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology* 3: 294–297.
- Houttuyn M (1772) *Natuurlyke historie of uitvoerige beschryving der dieren, planten en mineraalen* 17. Houttuyn, Amsterdam, 614 pp, pls. 126–138.
- Huang D, Benzoni F, Fukami H, Knowlton N, Smith ND, Budd AF (2014) Taxonomic classification of the reef coral families Merulinidae, Montastracidae and Diploastracidae (Cnidaria: Anthozoa: Scleractinia). *Zoological Journal of the Linnean Society* 171: 277–355. doi: 10.1111/zoj.12140
- Kropp RK (1990) Revision of the genera of gall crabs (Crustacea: Cryptochiridae) occurring in the Pacific Ocean. *Pacific Science* 44: 417–448.
- Kropp RK (1994) The gall crabs (Crustacea: Decapoda: Brachyura: Cryptochiridae) of the Rumphius Expeditions revisited, with descriptions of three new species. *Raffles Bulletin of Zoology* 42: 521–538.
- Kropp RK (1995) *Lithoscaptus pardalotus*, a new species of coral-dwelling gall crab (Crustacea: Brachyura: Cryptochiridae) from Belau. *Proceedings of the Biological Society of Washington* 108: 637–642.
- Kropp RK, Manning RB (1987) The Atlantic gall crabs, family Cryptochiridae (Crustacea: Decapoda: Brachyura). *Smithsonian Contributions to Zoology* 462: 1–21. doi: 10.5479/si.00810282.462
- Lamarck JBPA de M de (1816) *Histoire naturelle des animaux sans vertèbres*. Vol. 2, Verdier, Paris.
- Linnaeus C (1758) *Systema naturae* (ed. 10), 1. Laurentii Salvii, Holmiae, 824 pp.
- Meij SET van der (2014) Host species, range extensions, and an observation of the mating system of Atlantic shallow-water gall crabs (Decapoda: Cryptochiridae). *Bulletin of Marine Science* 90: 1001–1010. doi: 10.5343/bms.2014.1017
- Meij SET van der (2015) Host relations and DNA reveal a cryptic gall crab species (Crustacea: Decapoda: Cryptochiridae) associated with mushroom corals (Scleractinia: Fungiidae). *Contributions to Zoology* 84: 39–57.
- Meij SET van der, Reijnen BT (2014) The curious case of *Neotroglocarcinus dawydoffi* (Brachyura, Decapoda): biogeographic patterns resulting from isolation. *Systematics and Biodiversity* 12: 503–512. doi: 10.1080/14772000.2014.946979
- Milne-Edwards A (1862) Faune carcinologique de l'île de la Réunion. In: Maillard L (Ed.) *Notes sur l'île de la Réunion*. Vol. Annexe F. Paris, 16 pp.
- Milne Edwards H (1857) *Histoire naturelle des coralliaires ou polypes proprement dits* 2. Librairie Encyclopédique de Roret, Paris, 631 pp.
- Milne Edwards H, Haime J (1848) Note sur la classification de la deuxième tribu de la famille des Astréides. *Académie Des Sciences, Paris, Comptes Rendus* 27: 490–497.
- Milne Edwards H, Haime J (1849) *Recherches sur les polypiers*. Quatrième mémoire. Monographie des astréides(1). Tribu II. Astréens (Astreinae). *Annales Des Sciences Naturelles, Série 3, Zoologie* 11: 233–312.

- Ng PKL, Guinot D, Davie PJF (2008) *Systema Brachyurorum*: Part I. An annotated checklist of extant brachyuran crabs of the world. The Raffles Bulletin of Zoology Supplement No. 17: 1–286.
- Paul'son O (1875) *Studies on Crustacea of the Red Sea with notes regarding other seas. Part I. Podophthalmata and Edriophthalmata (Cumacea)*. S.V. Kul'zhenko, Kiev, xiv, 144 pp, plates 1–22. [In Russian, English translation by the Israel Program for Scientific Translations, Jerusalem, 1961]
- Scheer G, Pillai CSG (1983) Report on the stony corals from the Red Sea. *Zoologica, Stuttgart* 45(133): 1–198, pls. 1–41.
- Semper C (1881) *Animal life as affected by the natural conditions of existence*. D. Appleton and company, New York. [Originally published in German as: Semper K (1880) *Die natürlichen Existenzbedingungen der Tiere*. Brockhaus, Leipzig]
- Takeda M, Tamura Y (1983) Coral-inhabiting Crabs of the Family Hapalocarcinidae from Japan A Small Collection made at Kushimoto and Koza, Kii Peninsula IX. *Bulletin of the National Science Museum, Tokyo, series A (Zoology)* 9: 1–12.
- Verrill AE (1865) Classification of polyps (extract condensed from Synopsis of the Polyps and Corals of the North Pacific Exploring Expedition under Commodore C. Ringgold and Captain John Rodgers, U.S.N.). *Communications of the Essex Institute* 4: 145–152.
- Verrill AE (1866) Synopsis of the polyps and corals of the North Pacific Exploring Expedition, under Commodore C. Ringgold and Captain John Rodgers, U.S.N., from 1853 to 1856. Collected by Dr. Wm. Stimpson, naturalist to the expedition. With descriptions of some additional species from the west coast of North America. Pt. 3: Madreporaria. *Proceedings of the Essex Institute* 5(3): 17–50, pls. 1–2.
- Verrill AE (1901) Variations and nomenclature of Bermudian, West Indian and Brazilian reef corals, with notes on various Indo-Pacific corals. *Transactions of the Connecticut Academy of Arts and Sciences* 11: 63–168, pls. 10–35.
- Verrill AE (1908) *Brachyura and Anomura: their distribution, variations, and habits: Decapod Crustacea of Bermuda. I*. *Transactions of the Connecticut Arts and Sciences* 13: 299–474.