

UNTERSUCHUNGEN AN NIEDERLÄNDISCHEN MOOREN

K. WESTERWOLDE *).

Researches of Bogs in Westerwolde, province of Groningen

by

H. J. ESHUIS (Utrecht).

The bogs of S. E. Groningen are part of the great peat-marshes extending from S. E. Drente as far as N.W. Germany inclusive. So far as the territory of Westerwolde is concerned, people have begun digging off very early. According to the map by Kraayenhoff in 1816 nearly the whole peat-marsh westward from the line Blijham—Termaarsch had already been reclaimed, only a few parts still being covered with the original peat-layer (cf. map, fig. 1). The digging off east of the above line commences at the beginning of the 19th century on the borderland of Groningen and Drente.

Borings were performed in three places and the samples pollen-analytically and stratigraphically examined.

I. „Veenhuizer stukken”.

The first profile is from the „Veenhuizer stukken” north of Termaarsch. Peat has been dug off for the greater part here, in some places down to the sand. The bog still existing is about 1,50 m thick, this bigness probably having been greater: the stratigraphic investigations established that probably a portion has already been dug off.

The percentage of *Pinus* (cf. diagram I) is very great at the commencement of the pollen-diagram (98%), *Betula* hardly occurs (2%). The extension of *Corylus* is already very great (71%) and increases at once to a maximum (140%). Beyond the top of the diagram the *Corylus*-percentage decreases rapidly; nevertheless it keeps occurring largely (40%) in the whole profile up to 25 cm below the surface and then it descends to 20%.

As soon as the *Corylus*-curve begins to decline after its maximum,

*) See Recueil des Travaux botaniques néerlandais Vol. XXIX (1932), p. 1 and Vol. XXXII (1935) p. 430, also in Mededeelingen van het Botanisch Museum en Herbarium v. d. Rijksuniversiteit te Utrecht no. 1, 23 and 24.




Fig. 1.

the percentage of *Pinus* also falls off very quickly and it continues to move with slight deviations round an average of 5% up to the surface.

Meanwhile *Alnus* has augmented and as in most of the N.W. European profiles, *Alnus* also here has the greatest percentage of the pollen in the younger layers of the bog. Although in post-glacial times *Alnus* has been a tree greatly coming to the front in these parts, it is not in general characteristic of a definite period.




Diagram I: „Veenhuizer stukken”.

1 = Wood; 2 = Andromeda; 3 = Calluna; 4 = Erica; 5 = Monocotyledones; 6 = Scheuchzeria; 7 = Rhynchospora; 8 = Hypnaceae; 9 = Sphagnum.

Its extension in Central Europe is not nearly so great as in these regions (Rudolph 1930) and therefore the authors who have examined the bogs of N.W. Germany (Koch, Overbeck & Schmitz, Schubert and others) do not take a separate *Alnus* period for granted. For the growth of *Alnus* is determined by edaphic factors. Tüxen (1931) pointed to the fact that *Alnus* is a tree with typical demands of standing-place. Especially it grows on a very wet soil. The alder-marsh is poor in plant nutritious matter, it furnishes, however, good pasture-ground and therefore most of the *Alnetum* which have existed have changed into grass-land. Along most of the brooks of Westerwolde *Alnus* is sure to have had a great extension, the nature of the soil which was necessary for its growth being present. Soon after its appearance *Alnus* has spread maximally (81%). The percentage decreases when approaching the surface where it is yet 61.

Of the components of the *Quercetum mixtum* *Quercus* and *Tilia* appear at the same time, in the second place also *Ulmus*. *Tilia* and *Ulmus* are found only in small quantities, *Quercus* is the principal component of the *Quercetum mixtum*. A short time after the appearance of the *Quercetum mixtum* *Fagus* appears in very small quantities only, which quantities, however, increase quite near the surface. There also appears *Carpinus*. In the same time the *Ericaceae* have a maximal development (380%); *Betula* originally occurring in small quantities only at last also somewhat increases.

The whole profile practically includes only two periods:

1. a *Pinus-Corylus*-period, which terminates at the beginning of the diagram; *Pinus* and *Corylus* have a maximum, *Alnus* appears.

2. a *Quercetum mixtum-Corylus*-period; *Alnus* has a great extension, *Fagus* appears, later on also *Carpinus*; at the end of the diagram *Alnus* and *Corylus* decrease, *Betula* increases.

Though the classical terminology of the postglacial climate by Blytt & Sernander has been waved aside by Von Post (1930), Grosz (1930), K. Bertsch (1935) and others, we may use their nomenclature without the climatological connotation. Von Post pointed to the fact that the terminology established by Blytt & Sernander, originally only meant for Scandinavia, however conveyed to other regions of Europe, is now even obsolete for Sweden too. In the first place because neither in Sweden nor elsewhere the sole definite border-line between its periods, the subboreal-subatlantic limit, was a turning point for forest-evolution in the same degree as it was for bog-stratigraphy,

while the terms boreal, atlantic etc. were also inadequate for the other stages of development. Secondly, because more recent investigations, especially in Sweden, have shown that the changes of climate had been considerably more complicated than the old scheme suggested.

As a general division of the postarctic epoch V o n P o s t proposed in accordance with the main features of European forest-development a system of three periods:

1. The stage of the approach of the warm period, characterized by the appearance and the first increase of relatively heat-loving trees of different kinds.
2. The stage of the culmination of these forest-elements.
3. The stage of the decrease of the characteristic trees of the warm period and the appearance or the return of the predominant forest-constituents of the present day.

Summarizing the research of the last and best developed profiles we will trace how far the division of V o n P o s t is useful here.

If we use the terminology of B l y t t & S e r n a n d e r only as a division of time, therefore without the climatological connota-



Tree-stumps, found in the peat.

(photo A. Palle)

tion, we may suppose after comparison with the results of the investigation of neighbouring German bogs that the *Pinus-Corylus*-period is boreal, the *Quercetum mixtum-Corylus* period, falling past the crossing of the *Pinus*- and *Alnus*-curves, atlantic.

As we were saying the peat has been dug off down to the sand in some places. Fairly many tree-stumps have come out during the process, obviously originating from *Pinus* and having two forms. One kind (on the right side of the photo) has a normal vertical root-system, among whose parts there are great clods of sand. The tap-root of the other kind (on the left side of the photo), however, has been reduced and the lateral roots have developed horizontally; often the roots have curved upwards partly. This latter kind of tree-stump contains peat among the roots.

Evidently the stump coming from the sand is the older one, it stood in the diluvial underground. When the soil became wetter and peat began being formed, the tap-root remained rudimentary and the lateral roots have strongly developed, that the roots should find sufficient oxygen for their existence.

The conic form of the stumps of *Pinus* in the bog is striking. Their origin will have been as follows: The foot of the tree getting surrounded by peat, also the horizontal roots in the long run became short of oxygen and finally the tree died. The atmosphere and the water now acted most upon the border-line of air and peat (at A in fig. 2) and this process gradually decreases down to the lower end of the tree (B in fig. 2). When the decay has pro-




Fig. 2




Fig. 3

ceeded far enough the tree snaps off on the border-line of peat and air, and the conically formed lower end sticks with the roots in the soil, thus being no longer open to the influence of the weather (cf. fig. 3). The tree which keeps lying on the peat decays totally and this explains why mostly only tree-stumps and seldom whole trees are found in the peat. This, however, principally

holds good for *Pinus*; of *Quercus* often stems are found and no roots.

It is evident from the preceding, that the „Veenhuizer stukken” found their origin in the fact, that the soil of a forest changed into a marsh. In Germany this is called „Versumpfungsmoor”.

As to the further stratigraphic research, this has produced little that is of value. Of the principal elements of the peat in the older layers only wood and Monocotyledones prove to be recognizable; a nearer classification, however, was impossible in general. Exceptions to this are offered by *Scheuchzeria* and *Rhynchospora*: the former was easily to be recognized by its fibrous structure, of the latter fruits were found. The whole, however, gives an impression of wood-peat; gradually the trees must abandon the contest against the growth of the bog.

More upwards, in the atlanticum, first *Sphagnum* appears and next *Ericaceae* (twigs, leaves and flowers of *Calluna* and *Erica*, a seed of *Andromeda*): so the peat-formation has become from meso- oligotrophic. These layers belong to the old peat, which appears from the high degree of weathering; young *Sphagnum*-peat fails all over. Perhaps the upper part of the peat already belongs to the border-layer, i.e. the peat underlying the Grenz-Horizont (the percentage of *Ericaceae* is great here); it is not clear, however. We must take it for granted that the young *Sphagnum*-peat, if it has ever been present, has been dug off all over.

II. „Hoornder veen”.

The second profile was bored in the „Hoornder veen”, 3 km west of Wedde. Here too nearly the whole peat has been dug off and the land has been made fit for agriculture. The remaining part rises above the surroundings like an island.

The pollen-analytic research brought to the light that this profile is older than the preceding one; it begins in the preboreal. Just as the „Veenhuizer stukken” the bog lies on sand. The bottom-sample is very poor in pollen of trees, on the other hand many *Cyperaceae*-pollen were found: in onze preparation 2 *Betula*-, 2 *Pinus*-, and 4 *Corylus*-, but 65 *Cyperaceae*-pollen (cf. diagram II). From this we might conclude that at this time there were no forests (cf. *Firbas* 1935).

The late-glacial, i.e. the time which passed between the recession of the ice from the inmost Baltic final-moraine and the second Finnish *Salpausselkä*, was divided by *Firbas* into three periods:

1. an arctic period, which is characterized by forestlessness.




Diagram II: „Hoorder veen”.

1 = Charcoal; 2 = Vaccinium Oxycoccus; 3 = Erica; 4 = Calluna;
 5 = Andromeda; 6 = Monocotyledones; 7 = Eriophorum vaginatum;
 8 = Rhynchospora; 9 = Sphagnum.

2. a subarctic period, in which pure *Betula*- and *Pinus*-forests are found.

3. a proper preboreal period, in which we found the first appearance of the more demanding trees: *Corylus*, *Alnus* and the components of the *Quercetum mixtum*.

In the researched profile the first period is clearly to recognize. The second period, however fails; in the research, at least, it has not been found. The cause of this may be the too great distances between the researched samples; for in the sandy bottem-layers each cm represents a much longer period than in the higher parts of the bog, so that proportionally in the deeper layers the distance between the samples must be much smaller.

When the pollen-diagram begins, *Betula* and *Pinus* still dominate and though *Betula* still prevails, this tree already is decreasing. For a short time *Pinus* still rises, then reaches a maximum (71%) and afterwards is present during the whole profile, just as *Betula*, but in lower percentages.

Meanwhile the heat-loving trees, which at the beginning of the diagram already were found in small quantities, have increased greatly: *Alnus* soon reaches a high value (67%) and keeps moving round this same value as an average during the further development of the bog.

The boreal, which has begun about the *Pinus*-top, has its ending at the crossing of the *Pinus*- and *Alnus*-curves and is succeeded by the atlanticum. The *Corylus*-curve rises to a maximum of 91% and further remains in the profile up to the surface with an average value of 31%. In the „Hoornder veen” a *Corylus*-top lies about the middle of the atlanticum; in the „Veenhuizer stukken” this top still lies in the boreal. Koch (1929) found in the Münsterland in general a *Corylus*-top in the boreal and in the atlanticum a secondary rising of *Corylus*, which sometimes was greater than the first maximum. Van Raalte & Wassink (1932) found in Zwarte Meer now a top in the boreal, now in the atlanticum, now in both. Van Dobben (1932) found in the bog of Valthermond a *Corylus*-top in the atlanticum. Evidently now the one, now the other top may fail. It is possible too that one of both tops is not found on account of too great distances between the samples.

Before *Corylus* has its maximal value, the *Quercetum mixtum* already reaches a top. As to the order of appearance of the component parts of the *Quercetum mixtum* this must be thought as follows: first *Quercus* appears, then *Ulmus* and finally *Tilia*. *Quercus* first reaches a top (and together with *Quercus* the Quer-

cetum mixtum), then *Tilia* and finally *Ulmus*.

Two other tops concur with the *Corylus*-top: the group of *Varia* has a maximum of 42%, the *Ericaceae* even exceed 650%. At that time the surroundings may have been thinly wooded.

Even before the Grenz-Horizont now and again *Fagus* appears in small percentages; only passed the Grenz-Horizont *Fagus* rises above 10%, be it only for a short time. Then also the first *Carpinus* appears.

The history of the forests may be resumed here as follows:

1. a forestless period.
2. a short *Betula*-*Pinus*-period succeeded by
3. a *Quercetum mixtum*-*Corylus*-period in which *Alnus* immediately has a high value, *Quercus*, *Tilia*, *Ulmus*, and *Corylus* in turns reach maxima, *Betula* sometimes occurs in still important percentages and *Fagus* begins to appear.

A *Fagus*-period such as the German bogs mostly show and such as Ten Houten (1935) found in the „Korenburger veen” also, does not occur here.

Just as with the other two profiles, the stratigraphic research has been done less thoroughly, because when we aim at getting acquainted with the forest-history, it is of less importance than the pollen-analytic research.

In the preboreal and in the boreal only remnants of Monocotyledones are found. Shortly after the crossing of the *Pinus*- and *Alnus*-curve, in the atlanticum therefore, *Sphagnum* begins to help in the formation of the peat. Later on it is especially *Eriophorum vaginatum* and *Calluna* (twigs, leaves and flowers) and some less important *Ericaceae* (*Vaccinium Oxycoccus*, *Erica* and *Andromeda*) that compose the bulk of the samples. The main constituent is old *Sphagnum*-peat.

Remarkable is the occurrence of charcoal, which may be found from the atlanticum up to the Grenz-Horizont. The Grenz-Horizont is easily observed, and lies at 17 cm below the surface. The upper 7 cm of the peat have greatly decayed and are full of roots of the recent vegetation.

III. „Veeler veen”.

The last profile was bored in the „Veeler veen”, 6 km east of Wedde. At the time when Krayenhoff drew his map these regions were totally consisting of bog yet. Little by little large fragments have been reclaimed, indeed, but still there is original bog, which is being worked until the present day.

The formation of the peat has begun in moss-hags, great basins

in the ground from 10 to 20 m diameter. When these moss-hags had been filled, the bog kept growing in an upward direction, at the same time overgrowing the space between the moss-hags, so that at last the whole soil was covered with a layer of peat. By sounding it was possible to find the middle of a moss-hag, so that a profile of 4,38 m could be bored.

The bog lies on sand. A typical forestless period with high values for the Cyperaceae was not found, alas. The diagram




Diagram III: „Veeler veen”.

1 = Wood; 2 = Monocotyledones; 3 = Andromeda; 4 = Calluna; 5 = Erica; 6 = Menyanthes; 7 = Viola palustris; 8 = Myriophyllum; 9 = Nuphar; 10 = Batrachium; 11 = Betula nana; 12 = Eriophorum vaginatum; 13 = Potamogeton; 14 = Scheuchzeria; 15 = Scirpus; 16 = Carex; 17 = Bryales; 18 = Hypnaceae; 19 = Sphagnum imbricatum; 20 = Sphagnum spec.; 21 = Selaginella.

(cf. diagram III) begins at the time of the subarctic *Betula*- and *Pinus*-forests, the second period of *Firbas*. As remnants of the former tundra-vegetation in these layers were found *Selaginella selaginoides* (microspores) and *Betula nana* (leaf-fragments). In the „Soesterveen” *Selaginella* was found in the first, forestless period of *Firbas*; in the same period and at the rise of the subarctic forests also *Betula nana*. In the „Veeler veen” both are found somewhat later. Probably they found in the bog a refuge, which met their demands in the best way. The research of Vriezenveen („Bruine Haar”) by Florschütz & Wassink (1935), where *Selaginella* is found as late as the *Pinus-Corylus*-period, proves that also later on *Selaginella* may be found.

Betula, which at first with an average value of 90% far exceeds *Pinus*, decreases after some time and is then almost completely crowded out by *Pinus*. Near this change of domination a maximum of the *Cyperaceae* is found. This does not attain the 100% but points to a smaller density of forests during a short time or to a stronger growth of *Carices*.

Now there appear in succession *Quercus*, *Corylus* and *Alnus*, so here the third period of *Firbas* is reached. The preboreal ends more or less when the *Pinus*-curve reaches a top. In the boreal a rise of *Corylus* appears. *Pinus* soon declines, *Alnus* increases, and as in the former profiles the atlanticum begins with the crossing of the *Pinus*- and the *Alnus*-curve at about 40%.

Of the components of the *Quercetum mixtum* only *Quercus* is a rather important element. *Tilia* and *Ulmus* have a small share in the composition of the forests; the greatest percentage of *Tilia* is 3, of *Ulmus* 4. The *Quercetum mixtum* reaches a top with *Corylus*. Except as for *Fagus*, the diagram has a normal course.

Fagus, however, is to be found here in the atlanticum already at a depth of 1,75 m below the surface soon finding a maximum of extension there; next it is soon reduced again to nought and not until 0,75 m below the surface it appears again. In general *Fagus* begins in the Netherlands shortly before or after the Grenz-Horizont, at least with somewhat important percentages (Florschütz & others 1932, Florschütz & Wassink 1935, Ten Houten 1935). An exception to this is the bog of „Boerendijk” (Vriezenveen), which was explored by Florschütz & Wassink. The authors suppose *Fagus* to have settled during the atlanticum here and there on the sand, so that only at the outskirts it sent pollen into the bog.

Vermeer—Louman (1934) also mentions some finds of *Fagus* from old layers. In a boring near Sloten (province of

N. Holland) she found the empirical limit of *Fagus*-pollen (i.e. the beginning of the cohesive curve) in the boreal. In the atlanticum *Fagus* attains 4% of the total at a depth of 4 m; then the percentage gets low again. The rational limit of the pollen (i.e. the beginning of the continued rise of the curve) lies at 2,35 m, the bottom of the young *Sphagnum*-peat.

Polak (1929) too found already regularly *Fagus* in the old *Sphagnum*-peat of the „Riekerpolder”.

Ernst (1934) who also noticed the great fluctuation in the empirical and the rational limit of *Fagus*-pollen, distinguishes three types of curves of *Fagus*-pollen with regard to the Grenz-Horizont:

1. the *Fagus*-curve begins below the Grenz-Horizont, while the absolute, empirical and rational limit coincide. The curve rises without a minimum to a top.

2. the *Fagus*-curve begins as sub 1, while also the absolute, empirical and rational limit coincide. The curve makes a top immediately, but soon falls again, after which only above the Grenz-Horizont a maximum is formed.

3. the absolute, empirical and rational pollen-limits are separated by great distances. Only above the Grenz-Horizont the formation of a top begins.

Ernst points to the fact that *Fagus* must have spread irregularly and that at first only here and there it has obtained a foothold. Edaphic factors will have played a part here. The distribution by animals and men, which according to Ernst possibly has contributed to the spreading of *Fagus*, probably is of little importance. What factors, however, cause the minimum with the second type, to which the present case of the „Veeleer veen” belongs, Ernst does not point out. As *Fagus* according to F. Bertsch (1935) occurs only in a specific climate (*Fagus* wants an oceanic and shuns the continental climate), we might expect here a change of climate. The three types of Ernst, however, are not found regionally divided but in almost all districts explored, from the Ems as far as in the north-east of Denmark close together. So a factor of climate is impossible.

Though we cannot indicate a further distinct cause for the minimum in the *Fagus*-curve of the second type, it is evident that the importance of the Grenz-Horizont, at least with regard to the appearance of *Fagus* is only slight.

Summarizing we may distinguish in the „Veeleer veen”:

1. a *Betula*-*Pinus*-period,
2. a *Pinus*-*Betula*-period, during which *Quercetum mixtum*,

Corylus and *Alnus* begin to appear.

3. a *Quercetum mixtum-Corylus*-period, in which *Alnus* has a great extension and *Fagus* a small top.

4. a *Quercetum mixtum-Corylus-Fagus*-period, during which the quantity of *Fagus* increases for the second time, but in small density.

Distinguishing the climate-periods according to Von Post in this diagram meets with difficulties. The first period, the approach of the warm period with the appearance and the first extension of the heat-loving trees goes in the diagram nearly as far as the *Pinus-Betula*-period inclusive. This period is succeeded by the *Quercetum mixtum-Corylus*-period in which both *Quercus* and *Corylus* show a top, a culmination therefore of heat-loving trees, which is typical of the second period of Von Post. Limiting the third period, however, is more difficult. For a typical decrease of the characteristic trees of the second period (components of the *Quercetum mixtum* and *Corylus*) does not actually occur. Perhaps it is right to draw this line between the *Quercetum mixtum-Corylus*-period and the *Quercetum mixtum-Corylus-Fagus*-period as according to F. Bertsch *Fagus* might point to the fact that the climate became rather atlantic.

The stratigraphic research brought to the light that the peat in the moss-hag is caused by the change of a pool into land. From the sand as far as 3,20 m *Hypnaceae* are found which point to this fact. Further seeds of *Potamogeton*, *Nuphar luteum*, *Myriophyllum alterniflorum*, *Batrachium*, *Scirpus* and *Menyanthes*. Also *Carex* and *Scheuchzeria* were found in rather large quantities. Remarkable, however, is the presence of *Sphagnum* in the first eutrophic vegetation.

From 3,20 m up to 1,55 m as principal finds may be mentioned: *Scheuchzeria*, *Carex*, *Scirpus* and other *Monocotyledones* which could not be determined. Above this layer begins the old peat which, however, is not composed all over of *Sphagnum*, but for the greater part of *Bryales indeterminatae*. Rather soon there appear *Ericaceae*, first *Erica*, next *Calluna*.

Between 1,40 m and 1,30 m there was a layer of water in the bog, hence the interruption of *Erica* and *Bryales*.

A distinct border-layer cannot be distinguished; this proceeds unobservably from the old peat which ends at 1,10 m, where the young peat begins. The latter consists of not-determined *Sphagna* and especially of *Sphagnum imbricatum*. There are to be found *Ericaceae* (*Erica*, *Calluna* and *Andromeda*) and especially *Eriophorum vaginatum*.

To conclude a short comparison may follow of the profiles researched here with the bogs that are nearest: the bog of Valthermond (Van Dobben 1932) and the bogs of Emsland (Koch 1934 a, b, c). On most points these researches exhibit a great resemblance with the outcomes at Westerwolde.

The bog near Valthermond has developed in about an equal thickness as the „Veeler veen”. Of both the forest-history may be retraced as far as in the preboreal. The further development of the forests is nearly the same with both of them, with these exceptions, however, that Van Dobben found the absolute limit of the *Fagus*-pollen only after the Grenz-Horizont, and that he observed a decline of the *Corylus*-percentage after the subboreal-subatlantic contact. The stratigraphic research brought to the light that as far as the limit of the boreal and the atlanticum the peat was formed by *Hypnaceae*. This way of originating points to the change of a pool into land, so the same as happened with the formation of the „Veeler veen”. Here this was more evident, however, through the finds of *Potamogeton*, *Nuphar*, *Myriophyllum* and *Batrachium*.

Koch, on the other hand, nowhere found as origin of the peat-formation the change of a pool into land but always the change of a forest into marsh, probably by the rising of the ground-water; in the same way, therefore, as has been described as to the „Veenhuizer stukken” and the „Veeleer veen” outside the moss-hags. The bottom-layers of Koch's bogs always are wood-peat.

Koch's most fully developed profiles begin in the preboreal, in the subarctic period, in which no *Selaginella* was found indeed, but where *Betula nana* was shown by pollen-statistical numerations. The exactness of this method to demonstrate *Betula nana* is called in question, however; at any rate leaf-fragments are of more importance and these were not found by Koch.

Koch's oldest profile which is from the „Walchumer Moor” (8—10 km S.S.E. of Boertange) shows in the preboreal and the boreal repeatedly change of domination between *Betula* and *Pinus*. Possibly these changes would have been found in the „Veeler veen” and the „Hoornder veen” too, if the distances between the researched samples had been taken shorter. With Koch these are 5 cm, with the present researches 15 cm.

As for the further process of the forest-history this agrees especially with that of the „Veeler veen”, which is so to say part of the bogs of Emsland.

Fagus in most of the profiles appears before the Grenz-Horizont

and then takes an extension (e.g. in the „Walchumer Moor“) of 6 to 7% in the atlanticum and subboreal (in the „Veeler veen“ 10%). After the Grenz-Horizont *Fagus* reaches values of 25%, percentages not found at Valthermond nor at Westerwolde. Anyhow, in S.E. Groningen *Fagus* is already a very old tree.

BIBLIOGRAPHY.

- Bertsch, F. 1935. Das Pfrunger Ried und seine Bedeutung für die Florensgeschichte Südwestdeutschlands.
Beih. Bot. Centralbl. Bd. LIV, Abt. B, 185—243.
- Bertsch, K. 1935. Das deutsche Wald im Wechsel der Zeiten. Biologie in Einzeldarstellungen I. Tübingen.
- Brinkmann, P. 1934. Zur Geschichte der Moore, Marschen und Wälder Nordwestdeutschlands III. Das Gebiet der Jade.
Diss. Frankfurt a. M. 1933; Bot. Jahrb. Bd. LXVI, 1934.
- Dobben, W. H. van. 1932. Resultate von Untersuchungen an einigen niederländischen Mooren C. Valthermond.
Rec. Trav. Bot. Néerl. Vol. XXIX, 12—15 (also in Mededeelingen v. h. Botanisch Museum en Herbar. Utrecht No. 1).
- Ernst, O. 1933. Zur Geschichte der Moore, Marschen und Wälder Nordwestdeutschlands IV. Untersuchungen in Nordfriesland.
Diss. Frankfurt a. M., Schriften nat.wiss. Vereins für Schleswig—Holstein, Bd. XX, 211—334.
- Firbas, F. 1935. Die Vegetationsentwicklung des mitteleuropäischen Spätglacials.
Bibl. Botanica H. 112.
- Florschütz, F. und G. Vermeulen. 1932. Resultate von Untersuchungen an einigen niederländischen Mooren A. Soesterveen.
Rec. Trav. Bot. Néerl. Vol. XXIX, 1—6 (also in Mededeelingen v. h. Botanisch Museum en Herbar. Utrecht No. 1).
- Florschütz, F. und E. C. Wassink. 1935. Untersuchungen an niederländischen Mooren H. Vriezenveen.
Rec. Trav. Bot. Néerl. Vol. XXXII, 438—449 (also in Mededeelingen v. h. Botanisch Museum en Herbar. Utrecht No. 24).
- Grosz, H. 1930. Das Problem der nacheiszeitlichen Klima- und Florensentwicklung in Nord- und Mitteleuropa.
Beih. Bot. Centralbl. Bd. XLVII, 1—110.
- Houten, J. G. ten. 1935. Untersuchungen an niederländischen Mooren E. Korenburgerveen.
Rec. Trav. Bot. Néerl. Vol. XXXII, 430—437 (also in Mededeelingen v. h. Botanisch Museum en Herbar. Utrecht No. 23).
- Koch, H. 1929. Paläobotanische Untersuchungen einiger Mooren des Münsterlandes.
Diss. Frankfurt a. M., Beih. Bot. Centralbl. Bd. XLVI, Abt. B, 1—70.
- Koch, H. 1930. Stratigraphische und pollenfloristische Studien an drei nordwestdeutschen Mooren.
Planta Bd. II, 509—527.

- Koch, H. 1934a. Ein Profil aus dem Bourtanger Moor als Beispiel zur Moor- und Waldgeschichte an der Mittelems.
Ber. Deutsch. Bot. Ges. Bd. LII, 101—109.
- Koch, H. 1934b. Mooruntersuchungen im Emsland und im Hümmling.
Intern. Rev. d. ges. Hydrobiol. u. Hydrogr. Bd. 31, 109—156.
- Koch, H. 1934c. Untersuchungen zur Geschichte des Waldes an der Mittelems.
Bot. Jahrb. Bd. LXVI, 567—595.
- Overbeck, F. und H. Schmitz. 1931. Zur Geschichte der Moore, Marschen und Wälder Nordwestdeutschlands I. Das Gebiet von der Niederweser bis zur unteren Ems.
Mitt. Prov. Stelle Naturdenkmalpfl. Hannover, H. 3.
- Polak, B. 1929. Een onderzoek naar de botanische samenstelling van het Hollandsche veen.
Diss. Amsterdam.
- Post, L. von. 1930. Problems and working-lines in the postarctic forest-history of Europe.
Proc. 5th Intern. Bot. Congr. Cambridge, 48—54.
- Raalte, M. H. van und E. C. Wassink. 1932. Resultate von Untersuchungen an einigen niederländischen Mooren B. Zwarte Meer.
Rec. Trav. Bot. Néerl. Vol. XXIX, 6—12 (also in Mededeelingen v. h. Botanisch Museum en Herbar. Utrecht No. 1).
- Rudolph, K. 1930. Grundzüge der nacheiszeitlichen Waldgeschichte Mitteleuropas.
Beih. Bot. Centralbl. Bd. XLVII, Abt. B, 111—176.
- Schröder, D. 1930. Pollenanalytischen Untersuchungen in den Worpsweder Mooren.
Abh. nat.wiss. Vereins Bremen Bd. XXVIII, 13—30.
- Schubert, E. 1933. Zur Geschichte der Moore, Marschen und Wälder Nordwestdeutschlands II. Das Gebiet der Oste und Niederelbe.
Mitt. Prov. Stelle Naturdenkmalpfl. Hannover, H. 4.
- Tüxen, R. 1931. Die Grundlagen der Urlandschaftsforschung.
Nachr. Niedersachs. Urgesch. 1931, Nr. 5, 59—105.
- Vermeer—Louman, G. G. 1934. Pollenanalytisch onderzoek van de West-Nederlandsche bodem.
Diss. Amsterdam.