

Karyotypes of four species of Xenodontini snakes (Serpentes: Dipsadidae) and implications for taxonomy

Camila Falcione^{1,4}, Alejandra Hernando¹, Diego Andrés Barrasso², Diego Omar Di Pietro³

¹ Laboratorio de Herpetología, Facultad de Ciencias Exactas y Naturales y Agrimensura, Universidad Nacional del Nordeste, Av. Libertad 5470 (3400), Corrientes, Argentina

² Instituto de Diversidad y Evolución Austral (CONICET), Boulevard A. Brown 2915 (U9120ACD), Puerto Madryn, Chubut, Argentina

³ Sección Herpetología, División Zoología Vertebrados, Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata, Av. 122 y 60 s/n (B1900FWA) La Plata, Argentina

⁴ E-mail: camilafalcione@hotmail.com

Key words: karyosystematics, *Lygophis*, variability

Abstract

The karyotypes of four Xenodontini snake species, *Lygophis dilepis*, *L. meridionalis*, *L. flavifrenatus* and *L. anomalus*, are here described for the first time. We studied specimens from northeastern Argentina using conventional and silver (Ag-NOR) staining. While the typical ophidian karyotype is $2n = 36$, we found that the karyotype of the studied species is $2n = 34$, with metacentric and submetacentric chromosome pairs. The Ag-NOR staining revealed that nucleolar organizer regions (NORs) are located on one pair of microchromosomes. In *L. dilepis* and *L. anomalus* the 4th chromosome pair is heteromorphic, and we suggest that it might be considered as the ZW sex chromosome pair. The optimization of available karyological data on a molecular phylogenetic tree of the tribe Xenodontini shows that the diploid numbers $2n = 28$, 30 and 34 represent putative synapomorphy for *Erythrolamprus*, *Xenodon* and *Lygophis*, respectively. Our results provide new insights which fill gaps in our knowledge on the cytology in the genus *Lygophis* and identified a possible diagnostic character for the genus.

Contents

Introduction	265
Materials and methods	267
Results	268
Discussion	268
Acknowledgements	271
References	271
Appendix	273

Introduction

In recent years, phylogenetic studies based on morphological and molecular data have resulted in modifications in the systematic and phylogenetic relationships

of Serpentes (Zaher, 1999; Vidal *et al.*, 2000, 2010; Zaher *et al.*, 2009; Grazziotin *et al.*, 2012; Pyron *et al.*, 2013). The monophyletic tribe Xenodontini (Vidal *et al.*, 2000, 2010; Grazziotin *et al.*, 2012) is one of the South American snake radiations comprising about 70 species (Uetz and Jirí, 2013). According to the classification considered the clade is included in the subfamily Xenodontinae (Zaher *et al.*, 2009) or in the subfamily Dipsadinae of the family Colubridae (Pyron *et al.*, 2013).

The Xenodontini (sensu Vidal *et al.*, 2010 and Grazziotin *et al.*, 2012) is composed by the genera *Lygophis* Fitzinger 1843, *Erythrolamprus* Boie 1826 and *Xenodon* Boie 1826, clade also recover in other phylogenetic analysis (Zaher *et al.*, 2009; Pyron *et al.*, 2013, see nomenclatural discussion in Curcio *et al.*, 2009). The genus *Lygophis* was resurrected by Zaher *et al.* (2009), and confirmed by other studies (Vidal *et al.*, 2010; Grazziotin *et al.*, 2012; Pyron *et al.*, 2013). *Lygophis* comprises eight species grouped in the 'anomalus' and the 'lineatus' morphological groups, with three and five species respectively (Dixon, 1985, Michaud and Dixon, 1987).

The cytogenetics of the Xenodontini is poorly known. Chromosomal data are restricted to six *Erythrolamprus* species and four *Xenodon* species (Beçak, 1968; Beçak and Beçak, 1969; Beçak *et al.*, 1971, 1975; Gutiérrez *et al.*, 1984). Karyological information has been obtained using conventional cytological staining protocols. Localization of the nucleolar organizer regions (NORs) was only carried out on *E. poecilogyrus schotti* (Trajtengertz *et al.*, 1995).

Although information is scarce, the Xenodontini appear to be a karyologically diverse tribe. In fact, four

Table 1. Centromeric index (CI) and chromosome type of macrochromosome pairs (1-8) of *Lygophis* snakes analyzed in this study. a, the heteromorphic pair. Abbreviations: M = metacentric, SM = submetacentric.

Chromosome pair number	Sex		1	2	3	4 ^a	5	6	7	8	
<i>L. flavifrenatus</i>	♂	CI	47.62 ±1.08	46.37 ±2.23	45.56 ±2.98	46.27 ±1.99		46.68 ±1.56	46.99 ±1.90	44.90 ±1.92	45.69 ±1.69
		Type	M	M	M	M		M	M	M	M
<i>L. dilepis</i>	♀	CI	47.19 ±3.51	35.71 ±2.44	47.76 ±0.97	47.10 ±1.49	34.04 ±2.11	47.15 ±1.13	47.51 ±1.21	45.35 ±2.22	46.61 ±1.53
		Type	M	SM/M	M	M	SM	M	M	M	M
	♂	CI	47.30 ±2.63	35.95 ±2.15	47.77 ±1.09	45.33 ±2.27		45.74 ±2.49	47.01 ±2.23	46.94 ±1.99	46.33 ±1.46
		Type	M	SM/M	M	M		M	M	M	M
<i>L. meridionalis</i>	♀	CI	49.16 ±0.45	43.52 ±2.23	46.09 ±0.77	31.73 ±3.22	48.27 ±1.42	47.47 ±1.55	45.88 ±2.31	45.01 ±1.08	44.64 ±2.40
		Type	M	M	M	SM	M	M	M	M	M
<i>L. anomalus</i>	♀	CI	48.94 ±1.31	44.05 ±0.67	45.49 ±1.47	42.59 ±0.49	33.10 ±1.17	43.16 ±0.33	42.26 ±1.54	40.73 ±1.86	47.64 ±1.53
		Type	M	M	M	M	SM	M	M	M	M
	♂	CI	47.76 ±0.69	39.40 ±1.09	46.90 ±1.13	46.43 ±1.41		45.05 ±1.91	31.85 ±1.56	41.37 ±1.40	38.43 ±1.58
		Type	M	M	M	M		M	SM	M	SM/M

Table 2. Intra and intergeneric variation in the diploid number, chromosome formula, and sex chromosome morphology in species of tribe Xenodontini. Abbreviations: M = macrochromosome, m = microchromosome.

Species	2n	Chromosome formula	Sex chromosome morphology		Reference
			Z	W	
<i>Erythrolamprus aesculapii venustissimus</i>	28	16+4+8 (♂,♀)	Metacentric	Submetacentric	Beçak et al., 1966; Beçak and Beçak, 1969
<i>E. epinephelus</i>	28	16+4+8 (♂,♀)	-	-	Gutiérrez et al., 1984
<i>E. almadensis</i>	28	27+1+0 (♀)	Metacentric	Acrocentric	Beçak et al., 1975
		28+0+0 (♂)			
<i>E. bizona</i>	28	18+2+8 (♀)	Submetacentric	Submetacentric	Gutiérrez et al., 1984
<i>E. miliaris</i>	28	19+1+8 (♀)	Metacentric	Acrocentric	Beçak and Beçak, 1969
		20+0+8 (♂)			
<i>E. poecilogyus schotti</i>	32	-	-	-	Beçak et al., 1971; Trajtengertz et al., 1995
<i>Xenodon merremi</i>	30	16+0+14 (♂,♀)	Metacentric	Metacentric	Beçak, 1968
<i>X. neuwedii</i>	30	14+2+14 (♂,♀)	Metacentric	Metacentric	Beçak and Beçak, 1969
<i>X. severus</i>	30	14M+16m	-	-	Beçak et al., 1971
<i>X. rabdocephalus</i>	34	14+8+12 (♂,♀)	Submetacentric	Metacentric	Gutiérrez et al., 1984
<i>Lygophis anomalus</i>	34	16+0+18 (♂,♀)	Metacentric	Submetacentric	This study
<i>L. dilepis</i>	34	16+0+18 (♂,♀)	Metacentric	Submetacentric	This study
<i>L. meridionalis</i>	34	16+0+18 (♀)	-	-	This study
<i>L. flavifrenatus</i>	34	16+0+18 (♂)	-	-	This study

diploid numbers (2n = 28, 30, 32, 34), eight karyotype formulas, the ZW sex determination system, and a remarkable intra- and inter-generic karyotypic variability were described by previous studies (Beçak, 1968; Beçak and Beçak, 1969; Beçak et al., 1971, 1975; Gutiérrez et al., 1984). In *Erythrolamprus*, one karyo-

type 2n = 32 and five karyotypes 2n = 28 which differ in micro- and macro-chromosome number and macro-chromosomes morphology have been found (Beçak and Beçak, 1969; Beçak et al., 1971, 1975; Gutiérrez et al., 1984). Three species of *Xenodon* have the same diploid number of 2n = 30 including 14 or 16 macro-




Fig. 1. Mitotic karyotypes of four *Lygophis* species. A) *Lygophis dilepis* (female), B) *L. dilepis* (male), C) *L. meridionalis* (female), D) *L. flavifrenatus* (male), E) *L. anomalus* (female) and F) *L. anomalus* (male). Rounded boxes illustrate the heteromorphic pair. Scale bar = 10 μ m.

chromosomes and microchromosomes, while *X. rabdocephalus* ($2n = 34$) exhibits 22 macro-chromosomes and a reduced micro-chromosome complement of 12 (Beçak, 1968; Beçak and Beçak, 1969; Beçak *et al.*, 1971; Gutiérrez *et al.*, 1984).

Cytogenetic characters can be used to infer evolutionary relationships if they are analysed together with other independent characters (morphological, molecular, immunological, isozyme) (Sites and Reed, 1994). Indeed, such an approach has been demonstrated to be important to understand the diversity and evolution in snakes (Oguiura *et al.*, 2009; Mezzasalma *et al.*, 2014).

Hitherto no cytological characters were known for the genus *Lygophis*. To fill this information vacuum,

we documented the karyotype and the location of Ag-NORs in four species: *L. dilepis* Cope, 1862, *L. meridionalis* (Schenkel, 1902) and *L. flavifrenatus* Cope, 1862 (all belonging to the ‘*lineatus*’ group *sensu* Michaud and Dixon, 1987) and *L. anomalus* (Günther, 1858) (belonging to the ‘*anomalus*’ group *sensu* Dixon, 1985).

Material and methods

Chromosome analyses were carried out in males and females of *L. flavifrenatus*, *L. meridionalis*, *L. dilepis* and *L. anomalus* (see details in the Appendix). Voucher




Fig. 2. Metaphases after silver staining (Ag-NORs). Arrows indicate the NORs on a pair of microchromosomes. A) *Lygophis dilepis*, B) *L. meridionalis*, C) *L. flavifrenatus*. Scale bar = 10 μ m.

specimens are deposited in the Colección Herpetológica de la Universidad Nacional del Nordeste (UNNEC), Corrientes, Argentina.

Four hours prior to animal dissection, specimens were injected intraperitoneally with 0.1% colchicine (1 ml/100 g body weight). The euthanasia method proposed by Beaupre *et al.* (2004) was used. Chromosomes were obtained from intestinal epithelium by dispersion of cells on the hot stage. Chromosome preparations were stained conventionally using a 10% Giemsa solution at pH 6.8. The NORs were detected using the silver staining (Ag-NOR) technique applied by Howell and Black (1980). The Ag-NORs banding staining was performed only for species which provided a sufficient number and quality of metaphase plate.

To calculate the centromeric index (CI), the arms of macrochromosomes were measured on ten metaphase plates of each specimen using the software Micro-Measure version 3.3 (Reeves and Tear, 2000). The chromosomal formula was determined following Puccinini-Seale (1981): ($2n = I + II + III$), with I = metacentric or submetacentric macrochromosomes, II = telocentric or subtelocentric macrochromosomes and III = microchromosomes. Chromosomes that measured around 1 micron (μ m) were classified as microchromosomes.

To reconstruct karyotype evolution within the Xenodontini clade, the diploid number (from our results and from the literature) of Xenodontini species was optimized on the most recent phylogenetic hypothesis of Squamata by Pyron *et al.* (2013) using the par-

simony criterion with TNT software (Goloboff *et al.*, 2008).

Results

The diploid chromosome complements of *L. dilepis*, *L. meridionalis*, *L. flavifrenatus* and *L. anomalus* were similar (Fig. 1A-F). The karyotype was $2n = 34$ (16 + 0 + 18) and consisted of eight pairs of metacentric or submetacentric macrochromosomes, gradually decreasing in size (Table 1) and nine pairs of microchromosomes. No secondary constrictions were observed.

In females of *L. dilepis*, *L. meridionalis* and *L. anomalus* the 4th chromosomal pair was heteromorphic. In *L. dilepis* and *L. anomalus*, the larger element was metacentric and the smaller submetacentric (Fig. 1A, E). In *L. meridionalis*, the submetacentric corresponded to the larger chromosome and the metacentric to the smaller (Fig. 1C). In males of *L. dilepis*, *L. flavifrenatus* and *L. anomalus*, the homologous of pair 4 were metacentric and of similar size (Fig. 1B, D, F).

The NORs were detected on one pair of microchromosomes in *L. dilepis*, *L. flavifrenatus*, *L. meridionalis* (Fig. 2A-C).

Discussion

In snakes, chromosome numbers range from $2n = 24$ to $2n = 52$, with $2n = 36$ being the most frequent num-

ber (16 macrochromosomes and 20 microchromosomes) (Beçak and Beçak, 1969; Oguiura *et al.*, 2009). Karyotypes differing from this formula and with the chromosome number varying from $2n = 28$ to $2n = 34$ have been documented in 14 Xenodontini species belonging to the genera *Erythrolamprus* and *Xenodon*, as reported by several studies (Beçak, 1968; Beçak and Beçak, 1969; Beçak *et al.*, 1971; 1975; Gutiérrez *et al.*, 1984), and to the genus *Lygophis*, as here studied (Table 2).

Five *Erythrolamprus* species share the diploid number $2n = 28$ but with variations in the number of bi- and unarmed macro- and microchromosomes. *Erythrolamprus aesculapii venustissimus*, *E. epinephelus*, *E. bizona* and *E. miliaris* have karyotypes with 20 macrochromosomes and eight microchromosomes whereas 28 macrochromosomes and no microchromosomes were described in *E. almadensis* (Beçak and Beçak, 1969; Beçak *et al.*, 1975; Gutiérrez *et al.*, 1984). When considering all currently known chromosomal data for *Erythrolamprus* (Table 2), two out of the 10 pairs of macrochromosome are unarmed in *E. aesculapii venustissimus* (pairs 9 and 10) and *E. epinephelus* (pairs 6 and 9) and one in *E. bizona* (pair 6). In *E. miliaris* and *E. almadensis* all macrochromosomes are biarmed. In *E. poecilogyrus schotti*, $2n = 32$, and there is no clear distinction between macro- and microchromosomes (Beçak *et al.*, 1971; Trajtengertz *et al.*, 1995). In the genus *Xenodon*, *X. merremi* and *X. newiedii* have a similar $2n = 30$ karyotype with eight macrochromosome pairs and seven microchromosome pairs (Beçak, 1968; Beçak and Beçak, 1969). Although *X. severus* also exhibits $2n = 30$, it has seven macro- and eight microchromosomes (Beçak *et al.*, 1971). According to Beçak *et al.* (1971), in the karyotype of *X. severus*, a translocation that occurred between macrochromosomes can explain the numeric differences in the karyotypes of *Xenodon* species. Moreover, *X. rabdocephalus* exhibits the highest diploid number known in Xenodontini ($2n = 34$). Its karyotype includes 22 macrochromosome pairs of which 14 are biarmed and 8 unarmed, coupled to 12 microchromosomes (Gutiérrez *et al.*, 1984). The chromosome complement of *X. rabdocephalus* has more unarmed macrochromosome pairs but one or two fewer microchromosome pairs than the karyotype of $2n = 30$ *Xenodon* species (Beçak, 1968; Beçak and Beçak, 1969; Beçak *et al.*, 1971).

Lygophis dilepis, *L. meridionalis*, *L. flavifrenatus* and *L. anomalus* have a similar karyotype consisting of 34 chromosomes (16 biarmed macrochromosome

pairs and 18 microchromosomes). Although $2n = 34$ has also been reported for *X. rabdocephalus*, differences can be noted between these two taxa, since the *Lygophis* karyotype consists of eight pairs of biarmed macrochromosomes and three additional microchromosomes pairs when compared with the karyotype of *X. rabdocephalus*.

Considering that the karyotype $2n = 36$ (16 macrochromosomes and 20 microchromosomes) occurred in the common ancestor of snakes (Oguiura *et al.*, 2009), several mechanisms have been proposed to explain chromosomal evolution of Xenodontini. In *Erythrolamprus* ($2n = 28$ and 32) and *Xenodon* ($2n = 30$ and 34) the reduction in the diploid number resulted from unequal translocations between macro- and microchromosomes (Gutiérrez *et al.*, 1984) and Robertsonian fusions of microchromosomes (Beçak and Beçak, 1969). Additional centric fissions of macrochromosomes probably played an important role in the differentiation of *Xenodon* karyotypes (Gutiérrez *et al.*, 1984). On the other hand, we suggest that chromosomal inversion and reduction in the number of microchromosomes were involved in the karyotype evolution of *Lygophis* ($2n = 34$). Lower microchromosomes number also have observed in others Dipsadidae snakes (*Hydrodynastes*: $2n = 24$, *Philodryas serra*: $2n = 28$, *Thamnodynastes strigatus*: $2n = 32$ and *T. hypoconia*: $2n = 34$) (Beçak and Beçak, 1969).

The Boidae genera *Eryx*, *Acrantophis* and *Sanzinia* ($2n = 34$) and *Micrurus* species (Elapidae) from Central America ($2n = 26$ to $2n = 34$) tend to reduce the chromosome number probably due to fusion microchromosomes processes (Beçak and Beçak, 1969; Gutiérrez and Bolaños, 1979; Mengden and Stock, 1980; Oguiura *et al.*, 2009). From an ancestral karyotype $2n = 48$ in Pseudoxyrhophiinae (Lamprophiidae), the chromosomal diversification and reduction of the chromosome number may have occurred as a result of translocations of microchromosomes to macrochromosomes, tandem fusions and centric fissions and fusions (Mezzasalma *et al.*, 2014). Recently, it has been proposed that during diversification of Squamata the decrease in number of microchromosomes may have occurred by repeated fusions between macro- and/or other micro-chromosomes (Uno *et al.*, 2012).

The presence of ZW sex chromosomes in the 4th position was reported in many colubrids, elapids, and viperids (Beçak and Beçak, 1969; Singh, 1972; Gutiérrez *et al.*, 1979; Mengden and Stock, 1980; Beçak and Beçak, 1981; Ota, 1999; Aprea *et al.*, 2003, 2006) and has been considered a putative synapomorphy of the




Fig. 3. Diploid number optimization within Xenodontini clade based on the molecular phylogenetic hypothesis presented by Pyron et al. (2013). The outgroup consisted of the sister clade (*Uromacer catesbyi* + *U. frenatus* + *U. oxyrhynchus*). The gray branches represent ambiguity and the question symbols (?) the missing entries. On the right are the karyograms of the included species for which chromosomal information is available.

superfamily Colubroidea (Oguiura et al., 2009). The degree of heteromorphy of the ZW chromosomes is remarkable; they can be similar morphologically or differ in shape and/or size and in heterochromatin distribution (Beçak and Beçak, 1969; Singh, 1972; Mengden and Stock, 1980; Olmo, 1986; Oguiura et al., 2009). A ZZ:ZW system among Xenodontini was also reported for *X. merremi*, *X. neuwedii*, *X. rabdocephalus*, *E. almadensis*, *E. miliaris* and *E. bizona* (Table 2) (Beçak, 1968; Beçak and Beçak, 1969; Beçak et al., 1975; Gutiérrez et al., 1984).

In *L. dilepis* and *L. anomalus* females metaphases exhibited a metacentric/submetacentric heteromorphic 4th pair, and the respective males possessed two homo-

morphic metacentric chromosomes on the same position. Although based on the analysis of Giemsa-stained metaphases, these observations lead us to suggest that the 4th pair could be related to the ZW system of sex determination. Beçak and Beçak (1969) and Matsubara et al. (2006) suggested that, in snakes, heteromorphic sex chromosomes are the result of pericentric inversions, heterochromatinization and deletion of euchromatic regions.

Lygophis dilepis, *L. flavifrenatus* and *L. meridionalis* have a similar NORs location (one pair of microchromosomes). The same position has been reported for Xenodontini in *E. poecilogyrus schotti* (Trajten-gertz et al., 1995) and in Old and New World snakes

(Camper and Hanks, 1995; Aprea *et al.*, 2006). As such, this number and location of NORs appears to be prevalent in Serpentes (Camper and Hanks, 1995; Aprea *et al.*, 2006). However, in Xenodontini, further studies are necessary to understand the distribution of this character within this group.

Of the thirteen species of the Xenodontini tribe cytogenetically described to date eleven are included in the phylogeny proposed by Pyron *et al.* (2013) and the available data only allow evaluation of the chromosome number and morphology. Our optimization shows that diploid numbers represent putative synapomorphies for each genera (Fig. 3): *Lygophis* $2n = 34$, *Xenodon* $2n = 30$ and *Erythrolamprus* $2n = 28$ (with an increase in *E. poecilogyrus* to $2n = 32$). Unfortunately, we have no karyological data for the sister clade (*Uromacer catesbyi* + *U. frenatus* + *U. oxyrhynchus*) and for this reason it was not possible to evaluate the ancestral diploid number of the study group.

This study provides new data on the karyology of the Xenodontini. Yet, more studies on the karyology as well as on phylogenetic relationships in the Xenodontini species are necessary. Moreover, karyological information is available for three species that have not yet been included in molecular phylogenetic studies: *E. bizona* ($2n = 28$), *L. dilepis* ($2n = 34$), and *X. rabdocephalus* ($2n = 34$). Consequently, this information could not be optimized.

Although based on only four species, our study reinforces that there is remarkable variation in the karyotypes of Xenodontini. The karyotype $2n = 34 = 16 + 0 + 18$ is shared among the studied *Lygophis* species and could thus be a diagnostic character for the genus. Increasing the taxonomic sampling in combination with the analysis of the banding patterns will allow us to further elucidate chromosomal evolution in Xenodontini.

Acknowledgements

We thank Dirección de Recursos Naturales of the Corrientes Province (Argentina) and Dirección de Fauna y Parques of the Formosa Province (Argentina) for providing collecting permits, Brigitte Marazzi for constructive comments and language review, Rodrigo Cajade for valuable suggestions and comments to this manuscript, José Augusto Ruiz García, Lucila Curi and María Teresa Sandoval for help us with the images. This work was supported by the Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Argentina; and Secretaría General de Ciencia y Técnica, Universidad Nacional del Nordeste (SGCyT-UNNE), Argentina. We also wish thank to the two anonymous reviewers for their helpful comments.

References

- Aprea G, Odierna G, Andreone F, Glaw F, Vences M. 2003. Unusual karyotype in the Malagasy colubrid snake *Mimophis mahfalensis*. *Amphibia-Reptilia* 24: 215-219.
- Aprea G, Gentilli A, Zuffi MAL, Odierna G. 2006. The karyology of *Vipera aspis*, *V. atra*, *V. hugyi*, and *Cerastes vipera*. *Amphibia-Reptilia* 27: 113-119.
- Beaupre SB, Jacobson ER, Lillywhite HB, Zamudio K. 2004. *Guidelines for use of amphibians and reptiles in field and laboratory research, 2nd Edition*. Revised by the Herpetological Animal Care and Use Committee (HACC) of the American Society of Ichthyologists and Herpetologists, Lawrence, Kansas.
- Beçak W. 1968. Karyotypes, sex chromosomes and chromosomal evolution in snakes. Pp. 53-95 in: Bucherl W, Buckley E, Deulofeu V, eds, *Venomous Animals and their Venoms*. Vol. I. Academic Press, New York.
- Beçak W, Beçak ML. 1969. Cytotaxonomy and chromosomal evolution in Serpentes. *Cytogenetics* 8: 247-262.
- Beçak W, Carneiro SM, Beçak ML. 1971. Cariologia comparada em seis especies da colubrideos (Serpentes). *Ciencia e Cultura* 23: 123.
- Beçak ML, Beçak W, Chen TR, Shoffner RN. 1975. *Chromosome Atlas: Fish, Amphibians, Reptiles and Birds*. Vol. 3. Benirschke K, Hsu TC, eds. Springer-Verlag, Berlin.
- Beçak W, Beçak ML. 1981. Behaviour of the ZW Sex Bivalent in the Snake *Bothrops jararaca*. *Chromosoma* 83: 289-293.
- Camper DJ, Hanks BG. 1995. Variation in the Nucleolus Organizer Region among New World Snakes. *Journal of Herpetology* 29: 468-471.
- Curcio FF, Piacentini V, Fernandes DS. 2009. On the status of the snake genera *Erythrolamprus* Boie, *Liophis* Wagler and *Lygophis* Fitzinger (Serpentes, Xenodontinae). *Zootaxa* 2173: 66-68.
- Dixon JR. 1985. A review of *Liophis anomalus* and *Liophis elegantissimus*, and the description of a new species (Serpentes: Colubridae). *Copeia* 1985: 565-573.
- Goloboff PA, Farris JS, Nixon KC. 2008. TNT, a free program for phylogenetic analysis. *Cladistics* 24: 1-13.
- Grazziotin FG, Zaher H, Murphy RW, Scrocchi G, Benavides MA, Zhang YP, Bonatto SL. 2012. Molecular phylogeny of the New World Dipsadidae (Serpentes: Colubroidea): a reappraisal. *Cladistics* 28: 437-459.
- Gutiérrez JM, Bolaños R. 1979. Cariotipos de las principales serpientes coral (Elapidae: *Micrurus*) de Costa Rica. *Revista de Biología Tropical* 27: 57-73.
- Gutiérrez JM, Taylor RT, Bolaños R. 1979. Cariotipos de diez especies de serpientes costarricenses de la familia Viperidae. *Revista de Biología Tropical* 27: 309-319.
- Gutiérrez JM, Solórzano A, Cerdas L. 1984. Estudios cariológicos de cinco especies de serpientes costarricenses de la familia Colubridae. *Revista de Biología Tropical* 32: 263-267.
- Howell WM, Black DA. 1980. Controlled silver-staining of nucleolus organizer regions with a protective colloidal developer: a 1-step method. *Experientia* 3: 1014-1015.
- Matsubara K, Tarui H, Toriba M, Yamada K, Nishida-Umehara C, Agata K, Matsuda Y. 2006. Evidence for different origin of sex chromosomes in snakes, birds, and mammals and step-wise differentiation of snake sex chromosomes. *Proceedings of the National Academy of Sciences of the United States of America* 103: 18190-18195.

- Mengden GA, Stock D. 1980. Chromosomal evolution in Serpentes; a comparison of G and C chromosome patterns of some colubrid and boid genera. *Chromosoma* 79: 52-61.
- Mezzasalma M, Andreone F, Branch WR, Glaw F, Guarino FM, Nagy ZT, Odierna G, Aprea G. 2014. Chromosome evolution in pseudoxyrhophiine snakes from Madagascar: a wide range of karyotypic variability. *Biological Journal of the Linnean Society* 112: 450-460.
- Michaud EJ, Dixon JR. 1987. Taxonomic revision of the *Liophis lineatus* complex (Reptilia: Colubridae) of Central and South America. *Milwaukee Public Museum Contributions in Biology and Geology* 71: 1-26.
- Oguiura N, Ferrarezzi H, Batistic RF. 2009. Cytogenetics and molecular data in snakes: a phylogenetic approach. *Cytogenetics and Genome Research* 127: 128-142.
- Olmo E. 1986. A. Reptilia. Pp. 1-100 in: John B, ed., *Animal cytogenetics*, Vol. 4: Chordata 3. Gebrüder Bornträger, Berlin.
- Olmo E, Signorino G. 2005. Chromorep: a reptile chromosomes database. Accesible at <http://193.206.118.100/professori/chromorep.pdf>. Accessed: February 2015.
- Ota H. 1999. Karyotype of *Pareas iwasakii*: The first chromosomal description of a Pareatine snake (Colubridae). *Japanese Journal of Herpetology* 18: 16-18.
- Peccinini-Seale D. 1981. New developments in vertebrate cytology. IV. Cytogenetic studies in reptiles. *Genetica* 56: 123-148.
- Pyron RA, Burbrink FT, Wiens JJ. 2013. A phylogeny and revised classification of Squamata, including 4161 species of lizards and snakes. *BMC Evolutionary Biology* 13: 93.
- Reeves A, Tear J. 2000. *MicroMeasure for Windows, version 3.3*. Available at: <http://www.colostate.edu/Depts/Biology/MicroMeasure>.
- Singh L. 1972. Evolution of karyotypes in snakes. *Chromosoma* 38: 185-236.
- Sites JW, Reed K. 1994. Chromosomal evolution, speciation, and systematics: some relevant issues. *Herpetologica* 50: 237-249.
- Trajtengertz I, Beçak ML, Ruiz IR. 1995. Ribosomal cistrons in *Bothrops neuwiedi* (Serpentes) subspecies from Brazil. *Genome* 38: 601-606.
- Uetz P, Jirí H. 2013. The Reptile Database. Accessible at <http://www.reptile-database.org/>. [accessed 1 April 2015]
- Uno Y, Nishida C, Tarui H, Ishishita S, Takagi C, Nishimura O, Ishijima J, Ota H, Kosaka A, Matsubara K, Murakami Y, Kuratani S, Ueno N, Agata K, Matsuda Y. 2012. Inference of the protokaryotypes of amniotes and tetrapods and the evolutionary processes of microchromosomes from comparative gene mapping. *PLoS ONE* 7: e53027.
- Vidal N, Kindl SG, Wong A, Hedges SB. 2000. Phylogenetic relationships of Xenodontine snakes inferred from 12s and 16s ribosomal RNA sequences. *Molecular Phylogenetics and Evolution* 14: 389-402.
- Vidal N, Dewynter MI, Gower DJ. 2010. Dissecting the major American snake radiation: A molecular phylogeny of the Dipsadidae Bonaparte (Serpentes, Caenophidia). *Comptes Rendus Biologies* 333: 48-55.
- Zaher H. 1999. Hemipenial morphology of the South American xenodontine snakes, with a proposal for a monophyletic Xenodontinae and a reappraisal of colubroid hemipenes. *American Museum of Natural History* 240: 1-168.
- Zaher H, Grazziotin FG, Cadle JE, Murphy RW, Moura-Leite JC, Bonatto SL. 2009. Molecular phylogeny of advanced snakes (Serpentes, Caenophidia) with an emphasis on South American Xenodontines: a revised classification and descriptions of new taxa. *Papéis Avulsos de Zoologia* 28: 1-23.

Received: 29 July 2015

Revised and accepted: 27 November 2015

Published online: 5 July 2016

Editor: J. van Rooijen

Appendix

Species names, sex, collection numbers, and localities of specimens of *Lygophis* sampled in this study. UNNEC = Colección Herpetológica de la Universidad Nacional del Nordeste, Corrientes, Argentina.

Lygophis anomalus (n = 5): UNNEC-13001 (♀) Paso de los Libres (29°34'04,5''S, 57°25'45''W), Corrientes province, Argentina; UNNEC-13002 (♀) La Cruz (29°8'21,6''S, 56°52'35,5''W), Corrientes province, Argentina; UNNEC-11047 (♂) Mercedes (28°41'22''S, 57°28'32''W), Corrientes province, Argentina; UNNEC-10843 (♂) Mercedes (28°42'12''S, 57°28'26''W), Corrientes province, Argentina; UNNEC-11046 (♀) Medanos (33°25'54''S, 59°04'17''W), Entre Rios province, Argentina.

Lygophis dilepis (n = 6): UNNEC-09736 (♂) Fontana (25°20'18''S, 59°41'17''W), Formosa province, Argen-

tina; UNNEC-10102 (♂) Corrientes Capital (27°28'09''S, 58°46'56''W), Corrientes province, Argentina; UNNEC-10204 (♀), UNNEC-10211 (♀), and UNNEC-10212 (♀) ParajePerichón (27°25'45''S, 58°44'45''W), Corrientes province, Argentina; UNNEC-11831 (♂) Calchaquí (29°57'55''S, 60°19'49''W), Santa Fé province, Argentina.

Lygophis flavifrenatus (n = 2): UNNEC-11013 (♂) San Roque (28°52'37''S, 58°28'09''W), Corrientes province, Argentina; UNNEC-11263 (♂) Corrientes Capital (27°28'09''S, 58°46'56''W), Corrientes province, Argentina.

Lygophis meridionalis (n = 2): UNNEC-10209 (♀) Isla Apipé (27°31'12''S, 56°44'32''W), Corrientes province, Argentina; UNNEC-11263 (♀) Concepción (28°23'01''S, 57°51'57''W), Corrientes province, Argentina.