



New combinations in *Cenchrus* and *Urochloa* (Gramineae)

A.P.M. van der Zon¹

Key words

Brachiaria
Cenchrus
Gramineae
Pennisetum
sub-Saharan grasses
Urochloa

Abstract Recent taxonomic innovations have led to the inclusion of most of the *Brachiaria* species in *Urochloa* and of all *Pennisetum* species in *Cenchrus*. Many of the necessary combinations have been made for the West African species, but still nine new combinations and seven lectotypifications are presented here.

Published on 25 September 2019

INTRODUCTION

During the preparation of the *Gramineae* of West and Northern sub-Saharan Africa (WNSA, http://grasses-africa.linnaeus.naturalis.nl/linnaeus_ng/) new taxonomic insights in the delimitation of genera were encountered. Recent molecular phylogenetic studies (Chemisquy et al. 2010) have led to the amalgamation of the genera *Cenchrus* L. and *Pennisetum* Rich. and the inclusion of most of the species of *Brachiaria* (Trin.) Griseb. in the genus *Urochloa* P.Beauv. (Veldkamp 1996, Sosef 2016). The generic names of *Cenchrus* and of *Urochloa* have priority and so the species of *Pennisetum* and most of *Brachiaria* need to be transferred to them. Recently, Verloove (2002) and Sosef (2016) made new combinations, but a number of new combinations for species in the West African region still needs to be made. Therefore, nine new combinations are presented here, of which seven are lectotypified.

For on-line specimen information JStor Plants was consulted (plants.jstor.org), specimens seen only as image are denoted with an *.

TAXONOMY

Cenchrus gracilescens (Hochst.) Zon, *comb. nov.*

Pennisetum gracilescens Hochst. (1855) 199. — Lectotype (designated here): *Schimper 1411* (lecto P P00442918; isolecto G G00022563*, K K000281181, K00281182, L L0044929, P P00442250, S S14-20436*), Ethiopia, Dsche Dselia.

Lectotypification — Hochstetter (1855) designated the collection as type, but did not indicate a holotype among the duplicates.

Cenchrus laxius (Clayton) Zon, *comb. nov.*

Beckeropsis laxior Clayton (1963) 118. — *Pennisetum laxior* (Clayton) Clayton (1978) 580. — Type: *Keay FHI22678* (holo K K000281173), Nigeria, Idanre summit, Orusun.

Cenchrus nodiflorus (Franch.) Zon, *comb. nov.*

Pennisetum nodiflorum Franch. (1895) 363. — Lectotype (designated here): *Hens b32* (lecto P P00442930; isolecto G G0022556*, G00022557*, L L0822667, P P00442929*), Democratic Republic of Congo, Stanley pool, Kinshasa.

Lectotypification — Franchet (1895) cited two specimens: *Hens b32*, DRC, Stanley Pool, and *Dybowski s.n.* (PP02256337*), Congo, Brazzaville. *Hens b32* is selected as lectotype, based on a specimen in P, because it has most duplicates.

Cenchrus nubicum (Hochst.) Zon, *comb. nov.*

Gymnotrix nubica Hochst. (1844) 251. — *Pennisetum nubicum* (Hochst.) K.Schum. ex Engl. (1894) 58. — Lectotype (designated here): *Kotschy 13* (lecto W W0030467; isolecto G G00022329*, K K000281238, S S09-13390*), Sudan, Kordofan, Monte Beci.

Lectotypification — Hochstetter (1844) cited three specimens: *Kotschy 13*, *Kotschy 152* (BM, G G00022327*, HAL HAL0106782*, K K000281237*, L L0539824, L0539825, L0539827, M M0104086*, W W0030467*, WAG WAG0001536), Arasch Cool, Sudan and *Kotschy 368* (G G00022330*, S S09-13390, W W0030468*), Sudan. *Kotschy 13* from Vienna was selected.

Cenchrus pirottai (Chiov.) Zon, *comb. nov.*

Pennisetum pirottai Chiov. (1903) 37. — *Beckeropsis pirottai* (Chiov.) Stapf & C.E.Hubb. (1933) 269. — Lectotype (designated by Phillips 1995): *Ter-racciano & Pappi 2814* (lecto FT FT000357*; isolecto G G00022918*), Eritrea Barca, Agorda.

Note — Phillips (1995: 275) gives the FT collection as holotype. This collection should be considered as the lectotype, because Chiovenda (1903) did not select a holotype among the duplicates.

Cenchrus schweinfurthii (Pilg.) Zon, *comb. nov.*

Pennisetum schweinfurthii Pilg. (1901) 121. — Lectotype (designated here): *Schweinfurt 1500* (lecto K K000281247; isolecto B B 10 0167808*, BM BM000923366*, G G00022542*, US US00140920*, W W18890245022*), Sudan, Gallaba.

Note — Pilger (1901) did not indicate a holotype, therefore a lectotype is selected following Phillips (1995), who mentioned the K specimen as isotype.

¹ Naturalis Biodiversity Center, P.O. Box 9517, 2300 RA Leiden, The Netherlands; tonvanderzon@gmail.com.

Urochloa falcifera (Trin.) Zon, *comb. nov.*

Panicum falciferum Trin. (1826) 127. — *Brachiaria falcifera* (Trin.) Stapf (1919) 517. — Type: *Sabine s.n.* (not seen), Ghana, Accra.

Urochloa ovalis (Stapf) Zon, *comb. nov.*

[*Panicum ovale* R.Br. (1814) lxii, *nom. nud.*] — *Brachiaria ovalis* Stapf (1919) 546. — Lectotype (designated here): *Salt s.n.* (lecto K K000282117*), Abyssinia.

Note — Stapf (1919) cited four specimens, *Hildebrandt 337* (BM BM000923194*), Eritrea, Habab; *Salt s.n.* (K K000282117*), Abyssinia; *Thomson s.n.* (K K000282115*), British Somaliland, Haud, and *Appleton s.n.* (K K000282116*), British Somaliland, Bohotle? and Upper Sheikh. I selected *Salt s.n.* as lectotype, because this specimen was used by Brown (1814).

Urochloa serrifolia (Hochst.) Zon, *comb. nov.*

Panicum serrifolium Hochst. (1855) 196. — *Brachiaria serrifolia* (Hochst.) Stapf (1919) 548. — Lectotype (designated by Clayton 1982): *Schimper 2171* (lecto K K000282206), Ethiopia, Dschadscha.

Note — Hochstetter (1855) refers to the collections of Schimper from Abyssinia as original material. Clayton (1982: 596) gives as type: *Schimper 2171* (K), ?Ethiopia, Dschadscha. This should be considered as the lectotypification.

REFERENCES

- Brown R. 1814. List of new and rare plants collected in Abyssinia during the years 1805 and 1810. In: Salt H (ed), A voyage to Abyssinia and travels into the interior of that country, etc., Appendix 4: lxii. F.C. and J. Rivington, London.
- Chemisquy MA, Giussani LM, Scataglini, MA, et al. 2010. Phylogenetic studies favour the unification of Pennisetum, Cenchrus and Odontelytrum (Poaceae): a combined nuclear, plastid and morphological analysis, and nomenclatural combinations in Cenchrus. *Annals of Botany* 106: 107–130.
- Chiavenda E. 1903. *Familia Graminaceae* (2). *Annuario del Reale Istituto Botanico di Roma* 8 (1): 37–38, t. 2.
- Clayton W. 1963. Angiosperms of the Cambridge Annobon Island expedition. *Bulletin of the British Museum (Natural History), Botany* 3: 118.
- Clayton W. 1978. New species and combinations in African and Indian grasses. *Studies on Gramineae XLIII*. *Kew Bulletin* 32: 580.
- Clayton W. 1982. *Flora of Tropical East Africa, Gramineae* (part 3): 579–600, 675–690. Balkema, Rotterdam.
- Engler HGA. 1894. Über die Gliederung der Vegetation von Usambara und der angrenzenden Gebiete. *Abhandlungen der Königlichen Akademie der Wissenschaften in Berlin*: 1–86.
- Franchet AR. 1895. *Flore du Congo Français*. *Bulletin de la Société d'Histoire Naturelle d'Autun* 8: 363.
- Hochstetter CFF. 1844. *Gramina nova Africanas proponit et describit*. *Flora* 27: 251.
- Hochstetter CFF. 1855. *Plantas novas Africanas proponit et describit*. *Flora* 38: 196, 199.
- Phillips S. 1995. *Poaceae* (Gramineae). In: Hedberg I, Edwards S (eds), *Flora of Ethiopia and Eritrea* 7: 221–229, 261–275. The National Herbarium, Addis Ababa University, Ethiopia & The Department of Systematic Botany, Uppsala University, Sweden.
- Pilger RKF. 1901. *Gramineae africanas*. *Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie* 30: 121.
- Sosef M. 2016. Taxonomic novelties in central African grasses (Poaceae), *Panicaceae* 1. *Plant Ecology and Evolution* 149: 356–365.
- Stapf O. 1919. In: Oliver D (ed), *Flora Tropical Africa* 9: 517, 546, 548. Crown Agents for Oversea Governments and Administrations, London.
- Stapf O, Hubbard CE. 1933. Notes on African grasses: XIII. *Bulletin of Miscellaneous Information* 1933: 269–302.
- Trinius CB. 1826. *De Graminibus Paniceis*. *Impensis Academiae Imperialis Scientiarum, Petropoli*.
- Veldkamp JF. 1996. Proposal to conserve the name *Brachiaria* (Trin.) Griseb. (Gramineae) with a conserved type. *Taxon* 45: 319–320.
- Verloove F. 2002. New combinations in *Cenchrus* (Panicaceae, Poaceae) in Europe and the Mediterranean area. *Willdenowia* 42: 59–60.