



Changing the game: resolving systematic issues in key *Fusarium* species complexes

L. Lombard¹, M. Sandoval-Denis^{1,2}, L. Cai³, P.W. Crous^{1,2,4}

Published on 18 December 2019.

The genus *Fusarium* represents a ubiquitous group of fungi found in most natural substrates as either pathogens or non-pathogens associated with other fungi, plants and animals including humans (Leslie & Summerell 2006, Aoki et al. 2014). These fungi are also well-known for their ability to produce a cocktail of secondary metabolites that include important mycotoxins that pose a significant health risk to animals, humans and plants (Marasas et al. 1985, Logrieco et al. 2002, O'Donnell et al. 2018). Therefore, it is not unusual for this genus to be rated among the top 10 most economically and scientifically important plant pathogenic genera (Dean et al. 2012). However, as a large number of *Fusarium* species complexes include numerous cryptic species, several taxonomic/classification systems, most not subject to the International Code of Nomenclature for algae, fungi, and plants (ICN; Thurland et al. 2018), have been applied, sometimes resulting in erroneous assumptions. Furthermore, the lack of living ex-type cultures for several important *Fusarium* species has been a serious stumbling block in providing Latin binomials for several economically important cryptic *Fusarium* species.

Recent developments in the taxonomy of *Fusarium* and allied genera resulted in improved phylogenetic backbones for many species complexes (Guarnaccia et al. 2018, Sandoval-Denis et al. 2018, Maryani et al. 2019a, Lombard et al. 2019b, Sandoval-Denis & Crous 2019). However, much effort is still needed to fully discern the complexity of such a diverse group of taxa. This dedicated volume accommodates five papers mainly focusing on providing taxonomic stability to two important *Fusarium* species complexes (*F. oxysporum* (FOSC) and *F. incarnatum-equiseti* (FIESC)), and the related genus *Neocosmospora*. The intention of this volume is to highlight the importance of nomenclatural type studies to provide a stable platform for phylogenetic research to allow for the naming of cryptic species.

Fusarium oxysporum is probably the most commonly encountered species of *Fusarium*, and ranked fifth in the top 10 list of plant pathogenic fungi (Dean et al. 2012). This soil-borne asexual fungus includes both pathogenic (plants and animals, including humans) and non-pathogenic strains that display a complex phylogenetic structure of cryptic species. In the paper by Lombard et al. (2019a), *F. oxysporum* s.str. is epitypified to allow for the formal naming of 15 of these cryptic taxa. In addition, the various subspecific classification systems used for the FOSC are discussed, highlighting their limitations.

Panama disease, a devastating *Fusarium* wilt of both wild and cultivated banana, is associated with species in the FOSC (Maryani et al. 2019a), which have been well studied in recent years (Fourie et al. 2011, Ordóñez et al. 2015). However, limited information is available on non-FOSC *Fusarium* species also isolated in association with pathogenic *Fusarium* species causing Panama disease of banana. An extensive survey of local banana varieties in Indonesia resulted in a collection of 20 non-FOSC that were shown to belong to the *F. fujikuroi* (FFSC), FIESC and *F. sambucinum* (FSSC) species complexes, including five novel species for which Koch's postulates were tested (Maryani et al. 2019b).

Members of the FIESC include saprobes, fungicolous species, secondary invaders and pathogens of animals, humans and plants (O'Donnell et al. 2009, 2012, Sandoval-Denis et al. 2018). Using multi-locus phylogenies and Genealogical Concordance Phylogenetic Species Recognition (GCPSR), over 30 phylogenetic species have been recognised in this complex (O'Donnell et al. 2009, 2012, Villani et al. 2016, Maryani et al. 2019b, Santos et al. 2019, Torbati et al. 2019) for which less than 10 phylogenetic species have been provided with Latin binomials. This problem relates to the lack of nomenclatural types for several *Fusarium* species in this complex. In a survey of more than 22 plant species collected in eight provinces of China a total of 77 isolates were found to include 14 species in the FIESC, which included nine novel species provided with Latin binomials (Wang et al. 2019). Following this, Xia et al. (2019) provided more taxonomic stability to the FIESC through the epitypification of *F. compactum*, *F. incarnatum* and *F. scirpi*, and neotypification of *F. camptoceras*. This allowed for naming of 20 newly resolved and previously known phylogenetic species in the FIESC that lacked formal descriptions.

The genus *Neocosmospora*, previously known as the *F. solani* species complex, is ubiquitous, including saprobes, insect symbiont species as well as plant endophytes and important pathogens (animals, humans and plants) (Guarnaccia et al. 2018, Sandoval-Denis et al. 2018, Sandoval-Denis & Crous 2019). This genus encompasses a poorly characterised species-rich diversity, only now becoming evident through modern phylogenetic studies. However, several species in this genus lack nomenclatural types, placing a serious constraint on providing Latin binomials to new as well as already recognised phylogenetic species. In the monograph of Sandoval-Denis et al. (2019), a large number of type specimens and representative cultures were examined by means of morphology and phylogenetic inference. Through epi-, neo- and lectotypification, 68 species are now accepted in *Neocosmospora*, which also includes 13 new combinations and 29 new species. A further 11 recognised phylogenetic species remain undescribed and 17 doubtful or excluded taxa are also briefly discussed.

¹ Westerdijk Fungal Biodiversity Institute, Uppsalalaan 8, 3584 CT Utrecht, The Netherlands; corresponding author e-mail: l.lombard@wi.knaw.nl.

² Faculty of Natural and Agricultural Sciences, Department of Plant Sciences, University of the Free State, P.O. Box 339, Bloemfontein 9300, South Africa.

³ State Key Laboratory of Mycology, Institute of Microbiology, Chinese Academy of Sciences (CAS), Beijing, 100101, China.

⁴ Wageningen University and Research Centre (WUR), Laboratory of Phytopathology, Droevendaalsesteeg 1, 6708 PB Wageningen, The Netherlands.

Given the ubiquitous nature of *Fusarium*, and its importance as plant, human and animal pathogen, the incredibly high number of undescribed species that still exist in nature pose a serious threat for future food and fibre security. Furthermore, the high number of 'known' *Fusarium* species that are not tied to genetically well-characterised ex-type cultures is also a serious cause for concern. Papers in this volume provide an onset in addressing this issue, and clearly highlight the importance of nomenclatural type studies in resolving taxonomic difficulties surrounding important species complexes in the genera *Fusarium* and *Neocosmospora*. Making these types and associated DNA barcodes easily accessible through accessible public collections and databases is therefore of the utmost importance for future studies.

REFERENCES

- Aoki T, O'Donnell K, Geiser DM. 2014. Systematics of key phytopathogenic *Fusarium* species: current status and future challenges. *Journal of General Plant Pathology* 80: 189–201.
- Dean R, Van Kan JAL, Pretorius ZA, et al. 2012. The top 10 fungal pathogens in molecular plant pathology. *Molecular Plant Pathology* 13: 414–430.
- Fourie G, Steenkamp ET, Ploetz RC, et al. 2011. Current status of the taxonomic position of *Fusarium oxysporum* formae specialis *cubense* within the *Fusarium oxysporum* complex. *Infection, Genetics and Evolution* 11: 533–542.
- Guarnaccia V, Sandoval-Denis M, Aiello D, et al. 2018. *Neocosmospora perseae* sp. nov., causing trunk cankers on avocado in Italy. *Fungal Systematics and Evolution* 1: 131–140.
- Leslie JF, Summerell BA. 2006. *The Fusarium laboratory manual*. Blackwell Publishing Professional, Ames.
- Logrieco A, Mulè G, Moretti, et al. 2002. Toxigenic *Fusarium* species and mycotoxins associated with maize ear rot in Europe. *European Journal of Plant Pathology* 108: 597–609.
- Lombard L, Sandoval-Denis M, Lamprecht SC, et al. 2019a. Epitypification of *Fusarium oxysporum* – clearing the taxonomic chaos. *Persoonia* 43: 1–47.
- Lombard L, Van Doorn R, Crous PW. 2019b. Neotypification of *Fusarium chlamydosporum* – a reappraisal of a clinically important species complex. *Fungal Systematics and Evolution* 4: 183–200.
- Marasas WFO, Nelson PE, Toussoun TA. 1985. *Toxigenic Fusarium species. Identity and mycotoxicology*. The Pennsylvania State University Press, USA.
- Maryani N, Lombard L, Poerba YS, et al. 2019a. Phylogeny and genetic diversity of the banana *Fusarium* wilt pathogen *Fusarium oxysporum* f. sp. *cubense* in the Indonesian centre of origin. *Studies in Mycology* 92: 155–194.
- Maryani N, Sandoval-Denis M, Lombard L, et al. 2019b. New endemic *Fusarium* species hitch-hiking with pathogenic *Fusarium* strains causing Panama disease in small-holder banana plots in Indonesia. *Persoonia* 43: 48–69.
- O'Donnell K, Humber RA, Geiser DM, et al. 2012. Phylogenetic diversity of insecticolous *Fusaria* inferred from multilocus DNA sequence data and their molecular identification via FUSARIUM-ID and *Fusarium* MLST. *Mycologia* 104: 427–445.
- O'Donnell K, McCormick SP, Busman M, et al. 2018. Marasas et al. 1984 "Toxigenic *Fusarium* species: Identity and mycotoxicology" revisited. *Mycologia* 110: 1058–1080.
- O'Donnell K, Sutton DA, Rinaldi MG, et al. 2009. Novel multilocus sequence typing scheme reveals high genetic diversity of human pathogenic members of the *Fusarium incarnatum-equiseti* and *F. chlamydosporum* species complexes within the United States. *Journal of Clinical Microbiology* 47: 3851–3861.
- Ordonez N, Seidl MF, Waalwijk C, et al. 2015. Worse comes to worse: Bananas and Panama disease. When plant and pathogen clones meet. *PLoS Pathogens* 11: e1005197.
- Sandoval-Denis M, Crous PW. 2019. Removing chaos from confusion: assigning names to common human and animal pathogens in *Neocosmospora*. *Persoonia* 41: 109–129.
- Sandoval-Denis M, Guarnaccia V, Polizzi G, et al. 2018. Symptomatic Citrus trees reveal a new pathogenic lineage in *Fusarium* and two new *Neocosmospora* species. *Persoonia* 40: 1–25.
- Sandoval-Denis M, Lombard L, Crous PW. 2019. Back to the roots: a reappraisal of *Neocosmospora*. *Persoonia* 43: 90–185.
- Santos ACDS, Trindade JVC, Lima CS, et al. 2019. Morphology, phylogeny, and sexual stage of *Fusarium caatingaense* and *Fusarium pernambucanum*, new species of the *Fusarium incarnatum-equiseti* species complex associated with insects in Brazil. *Mycologia* 111: 244–259.
- Thurland NJ, Wiersma JH, Barbie FR, et al. (eds). 2018. *International Code of Nomenclature for algae, fungi, and plants (Shenzhen Code)*. Koeltz Botanical Books. [Regnum Vegetabile 159].
- Torbati M, Arzanlou M, Sandoval-Denis M, et al. 2019. Multigene phylogeny reveals new fungicolous species in the *Fusarium tricinctum* species complex and novel hosts in the genus *Fusarium* from Iran. *Mycological Progress* 18: 119–133.
- Villani A, Moretti A, De Saeger S, et al. 2016. A polyphasic approach for characterization of a collection of cereal isolates of the *Fusarium incarnatum-equiseti* species complex. *International Journal of Food Microbiology* 234: 24–35.
- Wang MM, Chen Q, Diao YZ, et al. 2019. *Fusarium incarnatum-equiseti* complex from China. *Persoonia* 43: 70–89.
- Xia JW, Sandoval-Denis M, Crous PW, et al. 2019. Numbers to names – restyling the *Fusarium incarnatum-equiseti* species complex. *Persoonia* 43: 186–221.