



Naturalis Repository

A new genus for *Philydorerythrocerum* and *P. fuscipenne* (Aves: Furnariidae)

George Sangster, Michael G. Harvey, Jimmy Gaudin & Santiago Claramunt

DOI:

<https://doi.org/10.11646/zootaxa.5361.2.11>

Downloaded from

[Naturalis Repository](#)

Article 25fa Dutch Copyright Act (DCA) - End User Rights

This publication is distributed under the terms of Article 25fa of the Dutch Copyright Act (Auteurswet) with consent from the author. Dutch law entitles the maker of a short scientific work funded either wholly or partially by Dutch public funds to make that work publicly available following a reasonable period after the work was first published, provided that reference is made to the source of the first publication of the work.

This publication is distributed under the Naturalis Biodiversity Center 'Taverne implementation' programme. In this programme, research output of Naturalis researchers and collection managers that complies with the legal requirements of Article 25fa of the Dutch Copyright Act is distributed online and free of barriers in the Naturalis institutional repository. Research output is distributed six months after its first online publication in the original published version and with proper attribution to the source of the original publication.

You are permitted to download and use the publication for personal purposes. All rights remain with the author(s) and copyrights owner(s) of this work. Any use of the publication other than authorized under this license or copyright law is prohibited.

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the department of Collection Information know, stating your reasons. In case of a legitimate complaint, Collection Information will make the material inaccessible. Please contact us through email: collectie.informatie@naturalis.nl. We will contact you as soon as possible.



A new genus for *Philydor erythrocerum* and *P. fuscipenne* (Aves: Furnariidae)

GEORGE SANGSTER^{1*}, MICHAEL G. HARVEY^{2,3}, JIMMY GAUDIN⁴ & SANTIAGO CLARAMUNT⁵¹Naturalis Biodiversity Center, Darwinweg 2, PO Box 9517, 2300 RA Leiden, Netherlands✉ g.sangster@planet.nl; <https://orcid.org/0000-0002-2475-7468>²Department of Biological Sciences, The University of Texas at El Paso, El Paso, TX 79968, USA✉ mgh272@gmail.com; <https://orcid.org/0000-0001-8050-6068>³Biodiversity Collections, The University of Texas at El Paso, El Paso, TX 79968, USA⁴34, avenue Antoine de Saint-Exupéry, 17 000 La Rochelle, France✉ jimmy.gdn.lr.taxo@gmail.com; <https://orcid.org/0000-0001-6519-1053>⁵Department of Ecology and Evolutionary Biology, University of Toronto, 25 Harbord Street, Room 537, Toronto, ON M5S 3G5, Canada✉ s.claramunt@utoronto.ca; <https://orcid.org/0000-0002-8926-5974>

*Corresponding author


A major task of modern taxonomy is to ensure that taxonomic groups above the species category are monophyletic (Hennig 1966; Cracraft 1981). Recent phylogenetic studies of the ovenbirds (Furnariidae) revealed numerous non-monophyletic genera and resulted in a step-wise revision of the group, including the description of six new genera (Chesser & Brumfield 2007; Chesser *et al.* 2009; Claramunt *et al.* 2010; Derryberry *et al.* 2010a, 2010b; Claramunt 2014; see also Claramunt *et al.* 2013). The first comprehensive species-level phylogeny of the group was based on three mitochondrial genes, one nuclear intron and two nuclear exons (Derryberry *et al.* 2011). This study revealed *Philydor* von Spix, 1824 as a taxonomically problematic genus, showing a fivefold polyphyly. This has been partially addressed in subsequent taxonomies by the placement of *P. erythropterum* (P.L. Sclater, 1856) and *P. rufum* (Vieillot, 1818) in the genus *Dendroma* Swainson, 1837 (Chesser *et al.* 2020; Gill *et al.* 2023; Remsen *et al.* 2023) and *P. ruficaudatum* (d'Orbigny & Lafresnaye, 1838) and *P. lichtensteini* (Cabanis & Heine, 1860) in an expanded genus *Anabacerthia* Lafresnaye, 1840b (Dickinson & Christidis 2014; Gill *et al.* 2023; Remsen *et al.* 2023). However, *Philydor*, as currently delimited (Gill *et al.* 2023; Remsen *et al.* 2023), is still polyphyletic because *P. erythrocerum* (von Pelzeln, 1859) and *P. fuscipenne* Salvin, 1866 are not closely related to the type species of *Philydor* (*P. atricapillus* zu Wied-Neuwied, 1821). In the phylogeny by Derryberry *et al.* (2011), *P. erythrocerum* and *P. fuscipenne* were sister to the monotypic genus *Megaxenops* Reiser, 1905, with strong support, whereas *P. atricapillus* and *P. pyrrhodes* (Cabanis, 1849) were inferred to be sister to the monotypic genus *Heliobletus* Reichenbach, 1853, again with strong support (Fig.1).

A more recent study by Harvey *et al.* (2020) used phylogenetic analysis of 2389 ultra-conserved genomic elements (UCEs) to estimate the relationships among suboscine passerines, including Furnariidae. In this study, the relationships among foliage-gleaners were incongruent with those inferred by Derryberry *et al.* (2011) (Fig. 1). However, *Philydor* (sensu Gill *et al.* 2023; Remsen *et al.* 2023) was again polyphyletic, with *P. erythrocerum* and *P. fuscipenne* inferred as the sister-group of the two species of *Anabazenops* Lafresnaye, 1840a (*A. dorsalis* (Sclater & Salvin, 1880) and *A. fuscus* (Vieillot, 1816)), and *P. atricapillus* as the sister species of *Heliobletus*, which together were the sister group of *P. pyrrhodes*.

Phylogenetic analysis clearly supports the removal of *P. erythrocerum* and *P. fuscipenne* from *Philydor*, which would become restricted to *P. atricapillus*, *P. novaesi* Teixeira & Gonzaga, 1983, and possibly *P. pyrrhodes*. Differences in structure (see below) strongly suggest that it would be unwise to group *P. erythrocerum* and *P. fuscipenne* with either of their inferred sister-taxa, *Megaxenops* or *Anabazenops* (Fig. 2), which would unnecessarily expand the morphological diversity of these genera. Thus, a separate genus is warranted for *P. erythrocerum* and *P. fuscipenne*.

No generic name appears to have been proposed for *P. erythrocerum* and *P. fuscipenne* (Sclater 1890; Sharpe 1901; Ridgway 1911; Cory & Hellmayr 1925; Peters 1951; Wolters 1977). Therefore, we suggest:

(a) Derryberry *et al.* (2011)



(b) Harvey *et al.* (2020)




FIGURE 1. Phylogenetic trees (cladograms) of *Philydor* and select representatives from related genera based on (a) Derryberry *et al.* (2011) and (b) Harvey *et al.* (2020). Numbers above branches refer to (a) posterior probability values and (b) bootstrap values. Nomenclature follows Gill *et al.* (2023).




FIGURE 2. External morphology of five species of foliage-gleaner (Philydorinae). Illustrations by Tim Worfolk (used with permission from *Birds of the world* / Lynx Edicions).

Neophilydor new genus

Type species: *Anabates erythrocerus* von Pelzeln, 1859 (currently *Philydor erythrocerum*).

Diagnosis: Small-sized foliage-gleaners (14–17 cm) most closely related to either *Megaxenops* or *Anabazenops*. Differs from both genera in having more elongated wings (hand-wing index 18–20 versus 12–17 in *Megaxenops* and *Anabazenops*, Claramunt *et al.* 2013), a generally shorter bill (length from nares ranges: 10.4–12.6 mm versus 12.5–16.9 mm, Claramunt *et al.* 2013), and generally shorter tarsus (17.8–20.8 mm versus 20.0–24.9 mm). The bill is overall straight with a slightly downcurved culmen in *Neophilydor*, contrasting with the heavy and deep bills of *Megaxenops* and *Anabazenops*. Bills of *Megaxenops* and *Anabazenops* have a straight culmen and, in *Megaxenops* and *A. fuscus*, an upturned tomium. The two species of *Neophilydor* further differ from *Megaxenops* in the presence of a pale postocular stripe (absent in *Megaxenops*), and darker, browner wings (bright rufous in *Megaxenops*). The species of *Neophilydor* further differ from *Anabazenops* by their smaller size, slender build (stocky in *Anabazenops*), and claw of hallux decidedly shorter than digit (claw equally long in *Anabazenops fuscus*; Ridgway 1911).

Differs from *Philydor* (*P. atricapillus*, *P. novaesi*, *P. pyrrhodes*) in generally less rounded tail (R1/R6 ratio: 1.09–1.25 versus 1.18–1.60).

Included taxa: *Neophilydor erythrocerum* **new comb.** (von Pelzeln, 1859) and *Neophilydor fuscipenne* **new comb.** (Salvin, 1866).

Etymology: Derived from the Greek νεος (*neos*) meaning new and *Philydor*, the genus of similar-looking foliage-gleaners in which the two species were formerly included. The gender of the name is neuter.

Acknowledgements

We thank Lynx Edicions for permitting the use of paintings of ovenbirds from the *Handbook of the Birds of the World*. Terry Chesser and Pamela Rasmussen provided helpful comments and edits that improved the quality of our paper.

References

- Cabanis, J. (1849) Vogel. In: Schomburgk, R. (Ed.), *Reisen in Britisch-Guiana in den Jahren 1840–1844: nebst einer Fauna und Flora Guiana's nach Vorlagen von Johannes Müller, Ehrenberg, Erichson, Klotzsch, Troschel, Cabanis und Andern*. Vol. 3. J.J. Weber, Leipzig, pp. 662–765.
- Cabanis, J. & Heine, F. (1860) *Museum Heineanum, Verzeichniss der ornithologischen Sammlung des Oberamtmann Ferdinand Heine auf Gut St. Burchard vor Halberstadt, mit kritischen Anmerkungen und Beschreibung der neuen Arten systematisch bearbeitet von Dr. Jean Cabanis erstem Custos der Königlichen zoologischen Sammlung zu Berlin und Ferdinand Heine Stud. Philos. Theil 2*. Schreibögel. R. Frantz, Halberstadt, 175 pp.
- Chesser, R.T. & Brumfield, R.T. (2007) *Tarphonimus*, a new genus of ovenbird (Aves: Passeriformes: Furnariidae) from South America. *Proceedings of the Biological Society of Washington*, 120, 337–339.
[https://doi.org/10.2988/0006-324X\(2007\)120\[337:TANGOO\]2.0.CO;2](https://doi.org/10.2988/0006-324X(2007)120[337:TANGOO]2.0.CO;2)
- Chesser, R.T., Billerman, S.M., Burns, K.J., Cicero, C., Dunn, J.L., Kratter, A.W., Lovette, I.J., Mason, N.A., Rasmussen, P.C., Remsen, J.V., Stotz, D.F. & Winker, K. (2020) Sixty-first supplement to the American Ornithological Society's check-list of North American birds. *Auk*, 137, ukaa030.
<https://doi.org/10.1093/auk/ukaa030>
- Chesser, R.T., Claramunt, S., Derryberry, E. & Brumfield, R.T. (2009) *Geocerthia*, a new genus of terrestrial ovenbird (Aves: Passeriformes: Furnariidae). *Zootaxa*, 2213 (1), 64–68.
<https://doi.org/10.11646/zootaxa.2213.1.4>
- Claramunt, S. (2014) Phylogenetic relationships among Synallaxini spinetails (Aves: Furnariidae) reveal a new biogeographic pattern across the Amazon and Paraná river basins. *Molecular Phylogenetics and Evolution*, 78, 223–231.
<https://doi.org/10.1016/j.ympev.2014.05.011>
- Claramunt, S., Derryberry, E.P., Chesser, R.T., Aleixo, A. & Brumfield, R.T. (2010) Polyphyly of *Campylorhamphus*, and description of a new genus for *C. pucherani* (Dendrocolaptinae). *Auk*, 127, 430–439.
<https://doi.org/10.1525/auk.2009.09022>
- Claramunt, S., Derryberry, E.P., Cadena, C.D., Cuervo, A.M., Sanín, C. & Brumfield, R.T. (2013) Phylogeny and classification of *Automolus* foliage-gleaners and allies (Furnariidae). *Condor*, 115, 375–385.
<https://doi.org/10.1525/cond.2013.110198>
- Cory, C.B. & Hellmayr, C.E. (1925) Catalogue of the birds of the Americas and the adjacent islands in Field Museum of Natural History. Part IV. Furnariidae, Dendrocolaptidae. *Field Museum of Natural History, Zoology Series*, 13 (4), 1–390.
<https://doi.org/10.5962/bhl.title.2975>
- Cracraft, J. (1981) Toward a phylogenetic classification of the recent birds of the world (Class Aves). *Auk*, 98, 681–714.
- Derryberry, E., Claramunt, S., Chesser, R.T., Aleixo, A., Cracraft, J., Moyle, R.G. & Brumfield, R.T. (2010a) *Certhiasomus*, a new genus of woodcreeper (Aves: Passeriformes: Dendrocolaptidae). *Zootaxa*, 2416 (1), 44–50.

<https://doi.org/10.11646/zootaxa.2416.1.2>

- Derryberry, E., Claramunt, S., O'Quin, K.E., Aleixo, A., Chesser, R.T., Remsen, J.V. & Brumfield, R.T. (2010b) *Pseudasthenes*, a new genus of ovenbird (Aves: Passeriformes: Furnariidae). *Zootaxa*, 2416 (1), 61–68.
<https://doi.org/10.11646/zootaxa.2416.1.4>
- Derryberry, E.P., Claramunt, S., Derryberry, G., Chesser, R.T., Cracraft, J., Aleixo, A., Pérez-Eman, J., Remsen, J.V. & Brumfield, R.T. (2011) Lineage diversification and morphological evolution in a large-scale continental radiation: the Neotropical ovenbirds and woodcreepers (Aves: Furnariidae). *Evolution*, 65, 2973–2986.
<https://doi.org/10.1111/j.1558-5646.2011.01374.x>
- Dickinson, E.C. & Christidis, L. (2014) *The Howard and Moore Complete Checklist of the Birds of the World. Vol. 2. Passerines. 4th Edition*. Aves Press, London, 752 pp.
- Gill, F., Donsker, D. & Rasmussen, P., eds. (2023) IOC world bird list. Version 13.1. Available from: <https://www.worldbirdnames.org/new/> (accessed 16 October 2023)
<https://doi.org/10.14344/IOC.ML.13.1>
- Harvey, M.G., Bravo, G.A., Claramunt, S., Cuervo, A.M., Derryberry, G.E., Battilana, J., Seeholzer, G.F., McKay, J.S., O'Meara, B.C., Faircloth, B.C., Edwards, S.V., Pérez-Emán, J., Moyle, R.G., Sheldon, F.H., Aleixo, A., Smith, B.T., Chesser, R.T., Silveira, L.F., Cracraft, J., Brumfield, R.T. & Derryberry, E.P. (2020) The evolution of a tropical biodiversity hotspot. *Science*, 370, 1343–1348.
<https://doi.org/10.1126/science.aaz6970>
- Hennig, W. (1966) *Phylogenetic Systematics*. University of Illinois Press, Urbana, Illinois, 263 pp.
- Lafresnaye, F. (1840a) *Anabazenops*. In: d'Orbigny, C. (Ed.), *Dictionnaire Universel d'Histoire Naturelle. Vol. 1*. Published by the author, Paris, pp. 411.
- Lafresnaye, F. (1840b) *Anabacerthia*. In: d'Orbigny, C. (Ed.), *Dictionnaire Universel d'Histoire Naturelle. Vol. 1*. Published by the author, Paris, pp. 412.
- d'Orbigny, A. & Lafresnaye, F. (1838) Synopsis Avium. *Magasin de Zoologie*, 8 (2), 1–34.
- von Pelzeln, A. (1859) Über neue Arten der Gattungen *Synallaxis*, *Anabates* und *Xenops* in der kaiserlichen ornithologischen Sammlung, nebst Auszügen aus Johann Natterer's nachgelassenen Notizen über die von ihm in Brasilien gesammelten Arten der Subfamilien: Furnariinae und Synallaxinae. *Sitzungsberichte Der Kaiserlichen Akademie Der Wissenschaften In Wien, Mathematisch-Naturwissenschaftliche Classe*, 34, 99–134.
- Peters, J.L. (1951) *Check-list of Birds of the World. Vol. 7*. Museum of Comparative Zoology, Cambridge, Massachusetts, 318 pp.
- Reichenbach, L. (1853) *Handbuch der Speciellen Ornithologie*. Expedition der Vollständigsten Naturgeschichte, Dresden und Leipzig, Lief. 3, 1–336.
- Reiser, O. (1905) [Vorläufigen Bericht über die ornithologische Ausbeute während der von der kais. Akademie der Wissenschaften im Jahre 1903 nach Brasilien entsendeten Expedition]. *Anzeiger der Kaiserlichen Akademie der Wissenschaften, Mathematisch-Naturwissenschaftliche Classe*, 42, 320–324.
- Remsen, J.V., Areta, J.I., Bonaccorso, E., Claramunt, S., Jaramillo, A., Lane, D.F., Pacheco, J.F., Robbins, M.B., Stiles, F.G. & Zimmer, K.J. (2023) A classification of the bird species of South America. Version 3 January 2023). American Ornithological Society. Available from: <http://www.museum.lsu.edu/~Remsen/SACCBaseline.htm> (accessed 16 October 2023)
- Ridgway, R. (1911) The birds of North and Middle America. *Bulletin of the United States National Museum*, 50 (5), 1–859.
- Salvin, O. (1866) Descriptions of eight new species of birds from Veragua. *Proceedings of the Zoological Society of London*, 1866, 67–76.
- Slater, P.L. (1856) On some additional species of birds received in collections from Bogota. *Proceedings of the Zoological Society of London*, Part 24, 25–31.
<https://doi.org/10.1111/j.1096-3642.1857.tb01190.x>
- Slater, P.L. (1890) *Catalogue of the Birds in the British Museum. Vol. 15. Catalogue of the Passeriformes, or Perching Birds, in the Collection of the British Museum. Tracheophonae*. Trustees of the British Museum (Natural History), London, 371 pp.
- Slater, P.L. & Salvin, O. (1880) On new birds collected by Mr. C. Buckley in Eastern Ecuador. *Proceedings of the Zoological Society of London*, 1880, 155–161.
<https://doi.org/10.1111/j.1469-7998.1880.tb06545.x>
- Sharpe, R.B. (1901) *A Hand-list of the Genera and Species of Birds. Vol. 3*. Trustees of the British Museum, London, 367 pp.
- von Spix, J.B. (1824) *Avium Species Novae Brasiliam. Vol. 1*. Hübschmann, München, 90 pp.
- Swainson, W. (1837) *The Natural History and Classification of Birds. Vol. 2*. Longman, Rees, Orme, Brown, Green, Longman and Taylor, London, 398 pp.
- Teixeira, D.M. & Gonzaga, L.P. (1983) Um novo Furnariidae do nordeste do Brasil: *Philydor novaesi* sp. nov. (Aves, Passeriformes). *Boletim do Museu Paraense Emílio Goeldi, Nova Série Zoologia*, 124, 1–22.
- Vieillot, L.P. (1816) *Analyse d'une Nouvelle Ornithologie Elementaire*. Deterville, Paris, 70 pp.
- Vieillot, L.P. (1818) In: *Nouveau Dictionnaire d'Histoire Naturelle. Vol. 26*. Abel Lanoe, Paris, 584 pp.
- zu Wied-Neuwied, M. (1821) *Reise nach Brasilien in den Jahren 1815 bis 1817. Vol. 2*. H.L. Brönnner, Frankfurt am Main, 345 pp.
<https://doi.org/10.5962/bhl.title.85967>
- Wolters, H.E. (1977) *Die Vogelarten der Erde. Lief. III*. Paul Parey, Hamburg, 80 pp.