



Naturalis Repository

A new genus for *Pycnopygius cinereus*/*P. ixoides* (Aves: Meliphagidae)

George Sangster, Petter Zahl Marki, Jimmy Gaudin, Martin Irestedt & Knud A. Jonsson

DOI:

<http://dx.doi.org/10.11646/zootaxa.5330.1.10>

Downloaded from

[Naturalis Repository](#)

Article 25fa Dutch Copyright Act (DCA) - End User Rights

This publication is distributed under the terms of Article 25fa of the Dutch Copyright Act (Auteurswet) with consent from the author. Dutch law entitles the maker of a short scientific work funded either wholly or partially by Dutch public funds to make that work publicly available following a reasonable period after the work was first published, provided that reference is made to the source of the first publication of the work.

This publication is distributed under the Naturalis Biodiversity Center 'Taverne implementation' programme. In this programme, research output of Naturalis researchers and collection managers that complies with the legal requirements of Article 25fa of the Dutch Copyright Act is distributed online and free of barriers in the Naturalis institutional repository. Research output is distributed six months after its first online publication in the original published version and with proper attribution to the source of the original publication.

You are permitted to download and use the publication for personal purposes. All rights remain with the author(s) and copyrights owner(s) of this work. Any use of the publication other than authorized under this license or copyright law is prohibited.

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the department of Collection Information know, stating your reasons. In case of a legitimate complaint, Collection Information will make the material inaccessible. Please contact us through email: collectie.informatie@naturalis.nl. We will contact you as soon as possible.



A new genus for *Pycnopygius cinereus*/*P. ixoides* (Aves: Meliphagidae)

GEORGE SANGSTER^{1*}, PETTER ZAHL MARKI², JIMMY GAUDIN³, MARTIN IRESTEDT⁴ & KNUD A. JONSSON⁴

¹Naturalis Biodiversity Center, Darwinweg 2, PO Box 9517, 2300 RA Leiden, the Netherlands.

✉ g.sangster@planet.nl; <https://orcid.org/0000-0002-2475-7468>

²Division of Research and Innovation, University of Agder, Universitetsveien 25, 4630 Kristiansand, Norway.

<https://orcid.org/0000-0001-6990-9915>

³34, avenue Antoine de Saint-Exupéry, 17 000 La Rochelle, France.

<https://orcid.org/0000-0001-6519-1053>

⁴Department of Bioinformatics and Genetics, Swedish Museum of Natural History, P.O. Box 50007, SE-10405, Stockholm, Sweden.

<https://orcid.org/0000-0003-1680-6861>; <https://orcid.org/0000-0002-1875-9504>

*Corresponding author

The honeyeaters (Meliphagidae) are an Australasian group of 196 species (Gill *et al.* 2023). This group shows great phenotypic, ecological and behavioural diversity and has been termed a “spectacular avian radiation” (Joseph *et al.* 2014), deservedly so because comparative studies have shown that the honeyeaters are more functionally diverse than other related and co-distributed non-nectarivorous meliphagoid clades (Marki *et al.* 2019). Phylogenetic analysis of DNA sequences and continued sampling of additional species has led to a much better understanding of the composition and phylogenetic relationships of this group (Driskell & Christidis 2004; Gardner *et al.* 2010; Nyári & Joseph 2011; Andersen *et al.* 2014, 2019; Joseph *et al.* 2014; Marki *et al.* 2017; McCullough *et al.* 2019). Multiple genera have been identified as para- or polyphyletic groups, including *Certhionyx*, *Anthochaera* and *Phylidonyris* (Driskell & Christidis 2004; Joseph *et al.* 2014), *Lichenostomus* (Gardner *et al.* 2010; Nyári & Joseph 2011; Joseph *et al.* 2014), *Xanthotis*, *Melidectes* and *Glycifohia* (Andersen *et al.* 2014), *Meliphaga* (Marki *et al.* 2017; McCullough *et al.* 2019), and *Lichmera* and *Pycnopygius* (Marki *et al.* 2017). As a result, multiple genera have been resurrected from synonymy, and other genera have been renamed or newly described (e.g. Nyári & Joseph 2011; Andersen *et al.* 2014, 2022; Joseph *et al.* 2014; McCullough *et al.* 2019).

Pycnopygius cinereus (Sclater, 1873) and *P. ixoides* (Salvadori, 1878) were originally described in the genus *Ptilotis* Swainson, 1837 (Sclater 1873; Salvadori 1878). In the first two decades of the 20th century, these species were grouped with several other honeyeaters in the genus *Ptilotis*, whereas *P. stictocephalus* Salvadori, 1876 was placed in the monotypic genus *Pycnopygius* Salvadori, 1880 (Dubois 1902, Ogilvie-Grant 1915). Subsequently, *Pycnopygius cinereus* and *P. ixoides* were included in *Pycnopygius* (Stresemann 1923; Mayr & Rand 1937; Mayr 1941). This is now universally followed (e.g. Salomonsen 1967; Wolters 1979; Sibley & Monroe 1990; Beehler & Pratt 2016; Clements *et al.* 2023; Gill *et al.* 2023).

Recent phylogenetic studies have revealed the genus *Pycnopygius* (sensu Salomonsen 1967; Gill *et al.* 2023) to be a polyphyletic assemblage (Marki *et al.* 2017; Hay *et al.* 2022). The phylogeny of Marki *et al.* (2017) placed *P. cinereus* and *P. ixoides* sister to a clade formed by *Prosthemadera novaeseelandiae* J.F. Gmelin, 1788 and two species of *Anthornis* G.R. Gray, 1840 [i.e., *A. melanura* (Sparrman, 1786) and *A. melanocephala* (G.R. Gray, 1843)], with poor support. However, these four species, together with *Certhionyx variegatus* Lesson, 1830, formed a well-supported clade (PP 0.99). The divergence of the two species of *Pycnopygius* from *Prosthemadera* and *Anthornis* was dated at 14 million years before present (Marki *et al.* 2017). *P. stictocephalus* was placed very distantly from *P. cinereus* and *P. ixoides* and was resolved as the sister taxon of *Grantiella picta* (Gould, 1838) with full support (PP 1.0).

The phylogeny of Hay *et al.* (2022) also showed *Pycnopygius* to be polyphyletic. In that study, *Pycnopygius cinereus* and *P. ixoides* were sister to *Certhionyx variegatus*, whereas *P. stictocephalus* again was sister to *G. picta*. Support for these relationships was not indicated.

The polyphyly of *Pycnopygius* means that *P. cinereus* and *P. ixoides* should be placed in a different genus than *P. stictocephalus*. The type species of *Pycnopygius* is *P. stictocephalus*, so that species remains in *Pycnopygius*. *P. cinereus*

and *P. ixoides* are not readily included in an existing genus. Due to differences in topology found by Marki *et al.* (2017) and Hay *et al.* (2022), the sister taxon of *P. cinereus* and *P. ixoides* remains uncertain. This, in combination with the deep divergence, uncertain support in Hay *et al.* (2022), and considerable morphological differences between *P. cinereus*/*P. ixoides* and *C. variegatus* (Fig. 1), means that placement of *P. cinereus* and *P. ixoides* in the genus *Certhionyx* is contra-indicated. Similarly, placement of *P. cinereus* and *P. ixoides* with the morphologically and genetically distinctive *Prosthemadera* and *Anthornis* in a single genus is neither well-supported nor helpful (Fig. 1). We conclude that *P. cinereus* and *P. ixoides* are best placed in their own genus. Examination of the relevant literature did not identify any available genus-group name for *Pycnopygius cinereus* and *P. ixoides* (Dubois 1902; Sharpe 1909; Salomonsen 1967; Wolters 1979). Therefore, we propose a new genus-group name:

***Pinarostola* new genus**

Type species: *Ptilotis cinerea* Sclater, 1873 (currently *Pycnopygius cinereus*).

Diagnosis: Medium-sized honeyeaters (18–22 cm). Both species differ from *Prosthemadera novaeseelandiae* in (i) smaller size (*P. novaeseelandiae* is 27–32 cm); (ii) crown dark olive-grey or blackish-brown, with grey or blackish-brown streaking or scaling (crown appearing dark blue or green-violet in *P. novaeseelandiae*); (iii) ear-coverts, chin, throat and underparts grey or dusky brownish-grey (appearing dark blue or green-violet in *P. novaeseelandiae*); (iv) side of throat without a pair of white ball-like tufts (present in *P. novaeseelandiae*); (v) head, neck, wings, tail and underparts without iridescence (present in *P. novaeseelandiae*); (vi) neck feathers concolorous with upperparts (neck with white lacy collar extending to upper mantle and side of neck in *P. novaeseelandiae*); (vii) lower belly, thighs and vent grey or dusky brownish-grey (black-brown in *P. novaeseelandiae*); and (viii) lesser and median wing-coverts olive-brown or dark brown (white in *P. novaeseelandiae*).




FIGURE 1. External morphology of five species of honeyeater (Meliphagidae), illustrating the differences in structure and plumage between two species of “*Pycnopygius*” and their inferred sister taxa (*Certhionyx* in one study, and *Prosthemadera*+*Anthornis* in another study). (a) *Certhionyx variegatus* female (Andrew Allen/iNaturalist), (b) *Certhionyx variegatus* male (hednota/iNaturalist), (c) *Pycnopygius ixoides* (Bradley Hacker/ML106934401), (d) *Pycnopygius cinereus* (Benoit Segerer/iNaturalist), (e) *Prosthemadera novaeseelandiae* (Nik Baines/iNaturalist), (f) *Anthornis melanura* male (digitaltrails/iNaturalist), (g) *Anthornis melanura* female (Glen Fergus/Wikimedia).

Both species differ from *Anthornis melanura* and *A. melanocephala* in (i) sexual dimorphism in plumage lacking (present in *Anthornis*); (ii) tail straight along axis (fairly narrow at base and becoming broader towards tip in *A. melanura*); (iii) head and neck without iridescent purple sheen (present in male *Anthornis*); (iv) lores grey or blackish-brown with fine blackish speckling (black in male *Anthornis*); (v) upperparts olive-brown or dark brown (olive-green in male *Anthornis*); (vi) underparts grey or dusky brownish-grey (olive-green in males and olive-brown in females of *Anthornis*); (vii) primaries and secondaries dark olive-brown or brown (black with bluish sheen in male *A. melanura*, with greenish sheen

in male *A. melanocephala*); and (viii) tail dark olive-brown or brown (dark brown to black-brown in *A. melanura*, dark brown with bluish sheen in male *A. melanocephala*).

Both species differ from *Certhionyx variegatus* in (i) small fleshy wattle or flange that is noticeably serrated or “toothed” hanging down from the lower eyelid lacking (present in *C. variegatus*); (ii) sexual dimorphism in plumage lacking (present in *C. variegatus*); (iii) mandible nearly straight (distinctly decurved in *C. variegatus*); (iv) bill blackish (bluish in *C. variegatus*); and (v) plumage grey, olive-brown, brown or blackish-brown with slight streaking or scaling (plumage very distinctly pied black and white in male *C. variegatus*).

Included taxa: *Pinarostola cinerea* **comb. nov.** (Sclater, 1873) and *Pinarostola ixoides* **comb. nov.** (Salvadori, 1878).

Etymology: Derived from the Greek πιναρός (*pinaros*) meaning dirty, and the Greek στολή (*stolē*) meaning dress or clothing. The genus name (‘dirty clothing’) refers to the dull, uneven, grey to brown plumage of the two species. The gender is feminine.

Acknowledgements

We thank the photographers, Andrew Allen, Nik Baines, digitaltrails, Glen Fergus, Bradley Hacker, hednota and Benoit Segerer, for making their work available for scientific use. Les Christidis and an anonymous referee provided constructive criticism that helped improve the paper.

Literature Cited

- Andersen, M.J., Naikatini, A. & Moyle, R.G. (2014) A molecular phylogeny of Pacific honeyeaters (Aves: Meliphagidae) reveals extensive paraphyly and an isolated Polynesian radiation. *Molecular Phylogenetics and Evolution*, 71, 308–315. <https://doi.org/10.1016/j.ympev.2013.11.014>
- Andersen, M.J., McCullough, J.M., Friedman, N.R., Peterson, A.T., Moyle, R.G., Joseph, L. & Nyari, A.S. (2019) Ultraconserved elements resolve genus-level relationships in a major Australasian bird radiation (Aves: Meliphagidae). *Emu*, 119, 218–232. <https://doi.org/10.1080/01584197.2019.1595662>
- Andersen, M.J., Gregory, S.M., Dickinson, E.C., Watling, D. & Scofield, R.P. (2022) A new genus-group name for *Gymnomyza* Reichenow, 1914 (Passeriformes: Meliphagidae) a preoccupied name in Diptera. *Avian Systematics*, 1, 27–29.
- Beehler, B.M. & Pratt, T.K. (2016) *Birds of New Guinea: Distribution, Taxonomy, and Systematics*. Princeton University Press, Princeton and Oxford, 668 pp. <https://doi.org/10.1515/9781400880713>
- Clements, J.F., Schulenberg, T.S., Iliff, M.J., Fredericks, T.A., Gerbracht, J.A., Lepage, D., Billerman, S.M., Sullivan, B.L. & Wood, C.L. (2022) *The eBird/Clements Checklist of Birds of the World. Version 2022*. Available from: <https://www.birds.cornell.edu/clementschecklist/download/> (accessed 31 July 2023)
- Driskell, A.C. & Christidis, L. (2004) Phylogeny and evolution of the Australo-Papuan honeyeaters (Passeriformes, Meliphagidae). *Molecular Phylogenetics and Evolution*, 31, 943–960. <https://doi.org/10.1016/j.ympev.2003.10.017>
- Dubois, A.J.C. (1902) *Synopsis avium: nouveau manuel d'ornithologie. Part I*. H. Lamertin, Bruxelles, 729 pp.
- Gardner, J.L., Trueman, J.W.H., Ebert, D., Joseph, L. & Magrath, R.D. (2010) Phylogeny and evolution of the Meliphagoidea, the largest radiation of Australasian songbirds. *Molecular Phylogenetics and Evolution*, 55, 1087–1102. <https://doi.org/10.1016/j.ympev.2010.02.005>
- Gill, F., Donsker, D. & Rasmussen, P. (Eds.) (2023) *IOC world bird list. Version 13.1*. Available from; <https://doi.org/10.14344/IOC.ML.13.1> (accessed 31 July 2023)
- Gmelin, J.F. (1788) *Systema Naturae, per regna tria Naturae: secundum Classes, Ordines, Genera, Species, cum Characteribus, Differentiis, Synonymis, Locis. Vol. 1. Pars 1. 13th Edition*. J.P. Delamollière, Lugduni [Lyon], 500 pp.
- Gould, J. (1838) *A Synopsis of the Birds of Australia, and the Adjacent Islands, Pt. 4*. Published by the author, London, 36 pp.
- Gray, G.R. 1840. *A List of the Genera of Birds, with an Indication of the Typical Species of each Genus*. R. & J.E. Taylor, London, 80 pp. <https://doi.org/10.5962/bhl.title.13777>
- Gray, G.R. (1843) List of the birds hitherto recorded as found in New Zealand, Chatham, and Auckland Islands, with their synonyma. In: Dieffenbach, E. (Ed.), *Travels in New Zealand. Vol. 2*. John Murray, London, pp. 186–201.
- Hay, E.M., McGee, M.D. & Chown, S.L. (2022) Geographic range size and speciation in honeyeaters. *BMC Ecology and Evolution*, 22, 86. <https://doi.org/10.1186/s12862-022-02041-6>
- Joseph, L., Toon, A., Nyári, Á.S., Longmore, N.W., Rowe, K., Haryoko, T., Trueman, J. & Gardner, J.L. (2014) A new synthesis of the molecular systematics and biogeography of honeyeaters (Passeriformes: Meliphagidae) highlights biogeographical

- and ecological complexity of a spectacular avian radiation. *Zoologica Scripta*, 43, 235–248.
<https://doi.org/10.1111/zsc.12049>
- Lesson, R.P. (1830) *Traite d'ornithologie, ou tableau methodique des ordres, sous-ordres, familles, tribus, genres, sous-genres, et races. Vol. 1.* F.G. Levrault, Paris, 659 pp.
<https://doi.org/10.5962/bhl.title.69307>
- Marki, P.Z., Jønsson, K.A., Irestedt, M., Nguyen, J.M., Rahbek, C. & Fjeldså, J. (2017) Supermatrix phylogeny and biogeography of the Australasian Meliphagides radiation (Aves: Passeriformes). *Molecular Phylogenetics and Evolution*, 107, 516–529.
<https://doi.org/10.1016/j.ympev.2016.12.021>
- Marki, P.Z., Kennedy, J.D., Cooney, C.R., Rahbek, C. & Fjeldså, J. (2019) Adaptive radiation and the evolution of nectarivory in a large songbird clade. *Evolution*, 73, 1226–1240.
<https://doi.org/10.1111/evo.13734>
- Mayr, E. (1941) *List of New Guinea birds. A Systematic and Faunal List of the Birds of New Guinea and Adjacent Islands.* American Museum of Natural History, New York, New York, 260 pp.
<https://doi.org/10.5962/bhl.title.68262>
- Mayr, E. & Rand, A.L. (1937) Results of the Archbold expeditions 14. Birds of the 1933–1934 Papuan Expedition. *Bulletin of the American Museum of Natural History*, 73, 1–248.
- McCullough, J.M., Joseph, L., Moyle, R.G. & Andersen, M.J. (2019) Ultraconserved elements put the final nail in the coffin of traditional use of the genus *Meliphaga* (Aves: Meliphagidae). *Zoologica Scripta*, 48, 411–418.
<https://doi.org/10.1111/zsc.12350>
- Nyári, A. & Joseph, L. (2011) Systematic dismantlement of *Lichenostomus* improves the basis for understanding relationships within the honeyeaters (Meliphagidae) and historical development of Australo-Papuan bird communities. *Emu*, 111, 202–211.
<https://doi.org/10.1071/MU10047>
- Ogilvie-Grant, W.R. (1915) Systematic list of the species. *Ibis* 57, Supplement, 2, 1–329.
- Salomonsen, F. (1967) Family Meliphagidae, honeyeaters. In: Paynter, R.A. (Ed.), *A Check-list of Birds of the World. Vol. 12.* Museum of Comparative Zoology, Cambridge, Massachusetts, pp. 338–450.
- Salvadori, T. (1878) Descrizione di trentuna specie nuove di uccelli della sottoregione papuana, e note intorno ad altre poco conosciute. *Annali del Museo civico di storia naturale di Genova*, 12, 317–347.
- Sclater, P.L. (1873) Characters of new species of birds discovered in New Guinea by Signor d'Albertis. *Proceedings of the Zoological Society of London*, 1873, 690–698.
- Sharpe, R.B. (1909) *A Hand-list of the Genera and Species of Birds. Vol. 5.* Trustees of the British Museum, London, 694 pp.
- Sibley, C.G. & Monroe, B.L. (1990) *Distribution and Taxonomy of Birds of the World.* Yale University Press, New Haven, Connecticut, 1111 pp.
- Sparrman, A. (1786) *Museum Carlsonianum. Fasc. 1.* Published by the author, Stockholm, 53 pp.
- Stresemann, E. (1923) Dr. Bürgers' ornithologische Ausbeute im Stromgebiet des Sepik. Ein Beitrag zur Kenntnis der Vogelwelt Neuguineas. *Archiv für Naturgeschichte, Abteilung A*, 7, 1–96.
- Swainson, W. (1837) *Natural History and Classification of Birds. Vol. 2.* Longman, London, 398 pp.
<https://doi.org/10.5962/bhl.title.64988>
- Wolters, H.E. (1979) *Die Vogelarten der Erde. Lief. IV.* Paul Parey, Hamburg, 80 pp.