


A first revision of the *Andrena* of Iraq (Hymenoptera, Andrenidae), with the description of two new species from Iraqi Kurdistan and additional records from surrounding countries

Thomas J. Wood¹, Halgurd R. Ismael², Daniele Baiocchi³, Mudhafar I. Hamad⁴, Tahseen T. Bapir², Marco Selis⁵

1 Naturalis Biodiversity Center, 2333 CR, Leiden, Netherlands

2 Department of Plant Protection, College of Agricultural Engineering Sciences, University of Duhok, Duhok, Kurdistan Region, Iraq

3 Via Matteo Babini 26, 00139, Roma, Italy

4 Khabat Technical Institute, Erbil Polytechnic University, Erbil, Kurdistan Region, Iraq

5 Via dei Tarquini 22, 01100 VT, Viterbo, Italy

Corresponding author: Thomas J. Wood (thomas.wood@naturalis.nl)



Academic editor: Thorleif Dörfel
Received: 1 February 2024
Accepted: 23 April 2024
Published: 1 July 2024

ZooBank: <https://zoobank.org/1093570E-D1C7-4918-BD8F-0BC3D3F486E5>

Citation: Wood TJ, Ismael HR, Baiocchi D, Hamad MI, Bapir TT, Selis M (2024) A first revision of the *Andrena* of Iraq (Hymenoptera, Andrenidae), with the description of two new species from Iraqi Kurdistan and additional records from surrounding countries. ZooKeys 1205: 267–298. <https://doi.org/10.3897/zookeys.1205.120033>

Copyright: © Thomas J. Wood et al.
This is an open access article distributed under terms of the Creative Commons Attribution License ([Attribution 4.0 International – CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

Abstract

Iraq is a large country in the Middle East region that borders both Turkey and Iran, countries known to host two of the largest bee faunas globally, as expected for a group of insects that favour dry to Mediterranean climates. Despite this huge regional species richness, the bee fauna of Iraq is chronically understudied and poorly known, both in relative and absolute terms. This is true for the hyper-speciose bee genus *Andrena*, for which only 17 species have been previously published for Iraq. This work is the first modern contribution to the revision of the *Andrena* fauna of Iraq. Based on new specimen collections in Duhok Governorate (Iraqi Kurdistan) during 2023, a revised total of 59 *Andrena* species for Iraq (42 species recorded for the first time) is presented, including the description of two new species: *Andrena (Aciandrena) duhokensis* Wood, **sp. nov.** and *Andrena (Notandrena) baiocchi* Wood, **sp. nov.** The unknown males of *A. (Micrandrena) elam* Wood, 2022, *A. (Micrandrena) obsidiana* Wood, 2022, and *A. (Notandrena) ayna* Wood, 2023 are described. *Andrena bakrajoensis* Amin & Mawlood, 2019, **syn. nov.** is synonymised with *A. (Holandrena) variabilis* Smith, 1853. Additional records are presented from nearby Middle Eastern countries, particularly Lebanon. These results highlight the fundamentally understudied nature of the Iraqi *Andrena* fauna.

Key words: Middle East, pan trap, solitary bees, taxonomy, understudied fauna

Introduction

Andrena is the second largest genus of bees with approximately 1,700 species following recent revisions (e.g., Pisanty et al. 2022a; Wood and Monfared 2022; Wood 2023, 2024). *Andrena* are known for their explosive radiation and rapid rate of speciation (Bossert et al. 2022), and are likely to

have evolved in the Middle East region ca 25 million years ago (Pisanty et al. 2022b). In line with this expected point of origin, the *Andrena* faunas of several Middle Eastern countries are very large, with ca 220 species known from Israel, 215 species known from Iran, and 388 species known from Turkey (Pisanty et al. 2022a; Wood and Monfared 2022; Wood 2023, 2024; Wood, unpublished data). The *Andrena* fauna of Syria is also large, but has not been comprehensively revised, and many species are present which have not yet had their occurrence formally published, with ~ 154–166 species known but not yet fully demonstrated or confirmed (Wood 2020; Wood et al. 2020; Pisanty et al. 2022a; Wood and Monfared 2022; Wood, unpublished data).

In contrast, the published *Andrena* fauna of the country of Iraq is extremely small, with only 17 species mentioned across various literature sources (Morice 1921a, 1921b; Derwesh 1965; Warncke 1969; Khalaf and Al-Omar 1974; Moalif 1994; Gusenleitner and Schwarz 2002; Scheuchl and Willner 2016; Augul 2018; Amin and Mawlood 2019; Ascher and Pickering 2023), some of which primarily cite previous publications without adding any new information, or which never present detailed specimen records at all. The Iraqi *Andrena* fauna has very rarely received dedicated attention, most frequently with species listed without supporting specimens or mentioned museum depositories, in strong contrast to neighbouring Turkey (e.g., Warncke 1965, 1974, 1975; Scheuchl and Gusenleitner 2009; Scheuchl and Hazir 2012; Hazir et al. 2014; Wood 2023) and to a lesser extent Iran and Syria (e.g., Popov 1967; Khodarahmi Ghahnavieh and Monfared 2019; Wood et al. 2020; Wood 2020; Radchenko et al. 2021; Wood and Monfared 2022).

Given the very high *Andrena* species richness found in nearby countries, it is clearly implausible that the Iraqi *Andrena* fauna is so small. In order to counter this knowledge deficit, we conducted new collections in northern Iraq during spring 2023, and critically reviewed the literature and museum collections in order to produce an updated faunal total.

Materials and methods

Andrena specimens were collected from various localities in Duhok province, Iraq (Kurdistan region) during May 2023 (Fig. 1: locations 1, 2, and 3). Specimens were collected using yellow pan traps which were filled with water and placed adjacent to vegetation (Fig. 2C).

Morphological terminology follows Michener (2007). The following abbreviations are used in the species descriptions: **A** = antennal segments, **S** = metasomal sterna, and **T** = metasomal terga. In diagnoses, the defining characters of a species are given, with those of the indicated comparison species given in parentheses. Subgeneric concepts follow Pisanty et al. (2022b). Specimens were measured from the centre of the clypeus at the front of the head to the apical tip of the metasoma to the nearest 0.5 mm. In the faunal list, taxa are presented alphabetically, first by subgenus and then by species. In the full faunal list, species entries followed by an asterisk (*) indicate the first record for Iraq. Likewise, for the global distributions of species, countries marked with an asterisk (*) indicate the first published record for that country. All specimens were identified by T.J. Wood.




Figure 1. Map of Iraq with novel data sites marked. New data locations: 1. Mount Gara, south of Sarsing, Duhok Governorate; 2. Besereh, Bablo, Duhok Governorate; 3. Mangesh, Duhok Governorate; 4. Mosul, Nineveh Governorate; 5. Hatra, Nineveh Governorate; 6. Baiji, Saladin Governorate. Landscape photographs correspond to localities 1 and 2.



Figure 2. Sampling participants in Duhok Governorate during May 2023 **A** sampling participants from left to right: Mudhafar Hamad, Daniele Baiocchi, Gianluca Magnani, Halgurd Ismael, Pierpaolo Rapuzzi **B** selection of sampling sites **C** yellow pan trap placed adjacent to vegetation **D** pan trap collection and inspection.

Photographs were taken using an Olympus E-M1 Mark II with a 60 mm macro lens. Additional close-ups were taken with the addition of a Mitutoyo M Plan Apo 5X infinity corrected objective lens. Photographs were stacked using Helicon Focus B (HeliconSoft, Ukraine) and plates were prepared in GNU Image Manipulation Program (GIMP) 2.10. Post-processing of some images was made in Photoshop Elements (Adobe Systems, USA) in order to improve lighting to highlight specific characters. The map of Iraq was taken from GISGeography (2024).

Abbreviations of depositories

DUMAI	University of Duhok, Museum of Agriculture College, Duhok, Iraq
MSVI	Personal collection of Marco Selis, Viterbo, Italy
NHMUK	Natural History Museum, London, United Kingdom
OÖLM	Oberösterreiches Landesmuseum, Linz, Austria
RMNH	Naturalis Biodiversity Center, Leiden, the Netherlands
TJWC	Personal collection of Thomas J. Wood, Leiden, the Netherlands

Results

Full Iraqi faunal list

1. *Andrena (Aenandrena) aeneiventris* Morawitz, 1872

Literature records. Gusenleitner and Schwarz (2002: dot map 11); Scheuchl and Willner (2016); Ascher and Pickering (2023).

Remarks. We have not examined any specimens of this species, but the presence of this species in Iraq is highly plausible based on its known global distribution, the map records indicated by Gusenleitner and Schwarz (2002), and the presence of this species in neighbouring Turkey and Iran.

Distribution. West and Central Palearctic (Gusenleitner and Schwarz 2002; Osytshnjuk et al. 2005).

2. *Andrena (Notandrena) aerinifrons* Dours, 1873 *

Material examined. IRAQ: Mosul, edges of a river, 7.iv.1988, 1♂, leg. Olejníček, OÖLM; **LEBANON:** Balbek-Hermel, Sefri, Haouch Snaid, AUB farm, 33.9244°N, 36.0754°E, 1000 m, 6.iv.2023, 1♂, 8♀, leg. T.J. Wood, TJWC.

Remarks. The status of the subspecies *levantina* Hedicke, 1938 (which nominally occurs in the Middle East) is unclear, and may not be merited, although *A. aerinifrons* sensu lato shows high and difficult-to-interpret intraspecific variation in its mitochondrial DNA barcode (Wood, unpublished data).

Distribution. *Andrena aerinifrons* sensu lato is distributed from Iberia and North Africa to the Middle East including Iraq* and Iran (Gusenleitner and Schwarz 2002; Wood and Monfared 2022).

3. *Andrena (Taeniandrena) afzeliella* (Kirby, 1802)

Literature records. Gusenleitner and Schwarz (2002: dot map 345, as *A. ovatula* (Kirby, 1802)); Ascher and Pickering (2023).

Remarks. We have not examined any specimens of this historically confused species (see Praz et al. 2022), but the presence of this species in Iraq is highly plausible based on its known global distribution, the map records indicated by Gusenleitner and Schwarz (2002), and the presence of this species in neighbouring Turkey and Iran where it is abundant.

Distribution. Somewhat unclear due to historical taxonomic confusion, but probably West and Central Palearctic (Praz et al. 2022; Wood and Monfared 2022; Ascher and Pickering 2023).

4. *Andrena (Melandrena) albifacies* Alfken, 1927

Literature records. Warncke (1969); Gusenleitner and Schwarz (2002: dot map 17); Grace (2010); Augul (2018); Wood and Monfared (2022); Ascher and Pickering (2023).

Remarks. We have not examined any specimens of this species, but the presence of this species in Iraq is highly plausible based on its known global distribution, the mention of this species from southern Iraq (Warncke 1969), the map records indicated by Gusenleitner and Schwarz (2002), and the presence of this species in neighbouring Iran.

Distribution. Morocco, Algeria, Tunisia, Libya, Egypt, Israel and West Bank, Jordan, Syria, Iraq, Iran (Warncke 1969; Gusenleitner and Schwarz 2002; Wood and Monfared 2022; Ascher and Pickering 2023).

5. *Andrena (Truncandrena) albopicta* Radoszkowski, 1874

Literature records. Gusenleitner and Schwarz (2002: dot map 18); Grace (2010); Augul (2018); Ascher and Pickering (2023).

Remarks. We have not examined any specimens of this species, but the presence of this species in Iraq is plausible: the species is present in south-eastern Turkey (see specimen records presented by Wood and Monfared 2022), and the dot map presented by Gusenleitner and Schwarz (2002) seems to indicate the presence of this species in northern Iraq. It is therefore tentatively accepted as present.

Distribution. Turkey, Armenia, Iraq, Iran (Gusenleitner and Schwarz 2002; Ascher and Pickering 2023), note that previous records from European Russia were erroneous, see Proshchalykin et al. (2023).

6. *Andrena (Melandrena) albopunctata* (Rossi, 1792)

Literature records. Gusenleitner and Schwarz (2002: dot map 19); Ascher and Pickering (2023).

Remarks. We have not examined any specimens of this species, but the presence of this species in Iraq is highly plausible based on its known global distribution, the map records indicated by Gusenleitner and Schwarz (2002), and the presence of this species in neighbouring Turkey and Iran.

Distribution. West and Central Palearctic (Gusenleitner and Schwarz 2002; Osytsnjuk et al. 2008).

7. *Andrena* (*incertae sedis*) *antilibanotica* Wood, 2020 *

Material examined. IRAQ: Duhok, Mt. Gara [S of Sarsing], 37.0158°N, 43.3506°E, 1912 m, 11.v.2023, 3♂, leg. D. Baiocchi, MSVI.

Distribution. Syria, Turkey, Iraq*, Iran (Wood and Monfared 2022; Wood 2023).

8. *Andrena* (*Chlorandrena*) *astica* Warncke, 1967 *

Material examined. IRAQ: Duhok, Mt. Gara [S of Sarsing], 37.0158°N, 43.3506°E, 1912 m, 11.v.2023, 3♂, leg. D. Baiocchi, MSVI/TJWC; **LEBANON:** Bekaa, Qob Elias valley, 33.7989°N, 35.8192°E, 900 m, 5.iv.2023, 1♂, leg. M. Boustani, TJWC.

Distribution. Greece, Turkey, Cyprus, Israel and West Bank, Lebanon*, Georgia, Iraq*, Iran (Schwenninger 2015; Wood and Monfared 2022).

9. *Andrena* (*Notandrena*) *ayna* Wood, 2023 *

Material examined. IRAQ: Duhok, Mt. Gara [S of Sarsing], 37.0158°N, 43.3506°E, 1912 m, 11.v.2023, 3♂, leg. D. Baiocchi, MSVI/TJWC.

Remarks and diagnosis. Wood (2023) described *A. ayna* from south-eastern Turkey (province of Hakkâri) based on two female specimens. Three male specimens from northern Iraq are now available. They are recognisable as *Notandrena* due to the short and broad head (Fig. 3B; 1.25× wider than long), the apex of the clypeus narrowed and slightly upturned, the at least partially yellow-marked clypeus, the broadened and weakly carinate gena (Fig. 3C; broader than the diameter of the compound eye), and the pronotum with a strong humeral angle. As in the female sex, *A. ayna* is immediately recognisable due to the sculpture of the scutum and scutellum which are almost entirely smooth and shiny over their entirely area, with scattered and sparse punctures (Fig. 3D).

Description. Male. Body length: 6.5–7 mm (Fig. 3A). **Head:** Dark, 1.25× wider than long (Fig. 3B). Clypeus variably yellow-marked, from medial ½ with yellow spot which does not extend to lateral margins to almost entirely dark with small pale dot medio-apically. Clypeus broadly flattened medially, surface densely and shallowly punctate, punctures separated by 0.5–1 puncture diameter, surface dull; apical margin narrow, projecting, anterior margin slightly upturned, weakly emarginate. Process of labrum tiny, trapezoidal, slightly wider than long, surface polished and shining. Gena 1.5× wider than diameter of compound eye, posterior margin weakly carinate, surface covered with curved striations (Fig. 3C); ocelloccipital distance subequal to diameter of lateral ocellus. Head covered with sparse light brownish pubescence, hairs not equalling length of



Figure 3. *Andrena (Notandrena) ayna* Wood, 2023 male **A** habitus, lateral view **B** face, frontal view **C** head, lateral view **D** scutum, dorsal view **E** terga, dorsal view **F** genital capsule, dorsal view.

scape. Antennae basally dark, A4–13 ventrally lightened by presence of greyish-brown scales; A3 exceeding A4, shorter than A4+5.

Mesosoma: Scutum and scutellum polished and shining over almost entire surface, scutum weakly shagreened anteriorly; scutum with scattered irregular punctures, punctures separated by 1–5 puncture diameters, scutellum almost impunctate (Fig. 3D). Pronotum with strongly produced humeral angle, deep vertical furrow with surface shining, remaining lateral face of propodeum

with longitudinal striations. Mesepisternum irregularly microreticulate, weakly shining. Dorsolateral parts of propodeum shagreened, weakly shining, sculpture overlain by network of irregular raised rugosity; propodeal triangle laterally defined by fine straight carinae, internal surface with network of rugae radiating from base, propodeal triangle not strongly differentiated from remaining propodeum. Mesosoma covered with sparse light brown hairs, none equaling length of scape. Legs dark, apical tarsal segments paler orange-brown, pubescence whitish. Hind tarsal claws with inner tooth. Wings hyaline, stigma and venation orange, nervulus antefurcal.

Metasoma: Tergal discs dark, marginal areas broadly lightened hyaline brown-white (Fig. 3E). Terga with sculpture variable, disc of T1 smooth and shining, T2–4 with base weakly shagreened, sculpture disappearing medially, tergal margins without sculpture. T1 with disc sparsely punctate, punctures separated by 2–3 puncture diameters, discs of T2–4 with punctures separated by 1–2 puncture diameters, marginal areas impunctate. Tergal discs with scattered pale hairs, not forming hairbands. T6–7 with light brown hairs overlying pseudopygidial plate. S8 columnar, short, apical margin truncate, ventral surface with sparse short brown hairs. Genital capsule compact, gonocoxae with inner margins almost forming rounded 90° angle, very weakly projecting (Fig. 3F). Gonostyli narrow basally, strongly broadening and flattened apically, inner margin raised, outer surface with short golden-brown hairs. Penis valves occupying ½ space between gonostyli, outer margins slightly thickened, progressively narrowing apically.

Distribution. South-eastern Turkey (Hakkâri) and northern Iraq* (Wood 2023).

10. *Andrena (Notandrena) baiocchii* Wood, sp. nov. *

<https://zoobank.org/6005A148-DD6C-45CE-8284-4DB733F080F1>

Material examined. Holotype: IRAQ: Duhok, Mt. Gara [S of Sarsing], 37.0158°N, 43.3506°E, 1912 m, 11.v.2023, 1♀, leg. D. Baiocchi, RMNH. **Paratypes:** IRAQ: Same information as holotype, 5♀, MSVI/RMNH/TJWC/DUMAI.

Diagnosis. *Andrena baiocchii* can be recognised as part of the subgenus *Notandrena* Pérez, 1890 due to the dorsolateral angle of the pronotum with a strong transverse ridge (= pronotum with a strong humeral angle), dull impunctate terga (Fig. 4F; hence placing it closer to members of the former subgenus *Carandrena* Warncke, 1968), scutum with distinct but weakly shining to dull green-purple metallic reflections (Fig. 4D), hind tibiae which weakly but distinctly broaden medially and apically, resembling a crude isosceles triangle and covered with simple scopal hairs, and lack of any other distinctive characters. This combination of characters places it close to *A. schlettereri* Friese, 1896 (Central Europe to Turkey), *A. purpleomicans* Alfken, 1935 (Turkey), *A. trimarginata* (Radoszkowski, 1886) (= *A. zosterata* Warncke, 1975; Middle East and Central Asia, see below), and *A. aerinifrons* Dours, 1873 sensu lato (Iberia, North Africa, and the Middle East). *Andrena baiocchii* can be separated from all these comparison species by the shape of the head which is almost round, only 1.1× wider than long (Fig. 4B; in comparison species is the head shorter, 1.2–1.4× wider than long) and the clypeus has contrasting surface sculpture, shagreened and weakly shining basally but becoming smooth and shining in its apical half AND with



Figure 4. *Andrena* (*Notandrena*) *baiocchii* sp. nov. female **A** habitus, lateral view **B** face, frontal view **C** clypeus, frontal view detail **D** scutum, dorsal view **E** propodeum, dorsal view **F** terga, dorsal view.

punctures becoming sparse, in the apical 1/3 of the clypeus with punctures separated by 1–3 puncture diameters (Fig. 4C). In comparison species the clypeus usually has a uniform sculpture, typically dull, never smooth and shining in its apical half (with the exception of *A. trimarginata*) and the clypeal punctures are consistently denser. *Andrena trimarginata* can be separated by the overall shape of the head (clearly wider than long) and by the density of clypeal punctures which are separated by a consistent 0.5–1 puncture diameters over the entire surface of the clypeus. Some individual species can also be separated

with additional characters, as *A. baiocchii* has a small body size of 7–8 mm (usually 8–9 mm in *A. aerinifrons*), the propodeal triangle has a network of finely raised rugae covering $\frac{3}{4}$ of its surface (Fig. 4E; surface smooth in *A. aerinifrons*), and the tarsi are dark and the terga are completely impunctate (with the tarsi lightened orange and the terga with shallow but distinct punctures in *A. purpureomicans*). The male is unknown, but is expected to share the relatively elongate head which is unusual within this group of species.

Description. Female. Body length: 7–8 mm (Fig. 4A). **Head:** Dark, 1.1× wider than long (Fig. 4B). Clypeus domed, variably sculptured, in basal $\frac{1}{2}$ with fine granular shagreen, weakly shining, in apical $\frac{1}{2}$ smooth and shining, polished; surface punctate, punctures separated by 1–3 puncture diameters (Fig. 4C). Process of labrum small, rounded trapezoidal, 2× wider than long, apical margin weakly emarginate. Gena marginally exceeding diameter of compound eye; ocelloccipital distance equals $\frac{1}{2}$ diameter of lateral ocellus. Foveae dorsally occupying slightly $< \frac{1}{2}$ space between compound eye and lateral ocellus, slightly narrowing ventrally, separated from inner margin of compound eye by distance subequal to its own diameter; foveae filled with brown hairs. Face, gena, vertex, and scape with sparse light brown hairs, none equalling length of scape. Antennae basally dark, A5–12 ventrally lightened by presence of grey-orange scales; A3 slightly exceeding A4+5, slightly shorter than A4+5+6.

Mesosoma: Scutum and scutellum with dense fine granular microreticulation, weakly shining to dull, laterally and anteriorly with weak but distinct green-purple metallic reflections; surface irregularly and obscurely punctate, punctures separated laterally by 1–3 puncture diameters, medially by 3–5 puncture diameters (Fig. 4D). Pronotum with strong humeral angle. Mesepisternum and dorsolateral parts of propodeum with dense granular microreticulation, dull; propodeal triangle delineated by change in surface sculpture, predominantly covered with finely raised rugae over $\frac{3}{4}$ of its surface (Fig. 4E). Mesosoma with light brown hairs, longest on mesepisternum, not equalling length of scape. Propodeal corbicula incomplete, dorsal fringe composed of long plumose light brown hairs, internal surface with scattered light brown hairs. Legs predominantly dark, apical tarsal segments lightened brownish, pubescence light brown. Flocculus complete but relatively sparse, composed of light brown plumose hairs; femoral and tibial scopae composed of light brown-golden simple hairs. Hind tarsal claws with very small inner tooth. Wings hyaline, stigma bright orange, venation dark orange, nervulus antefurcal.

Metasoma: Terga dark, apical margins weakly but distinctly depressed, partially lightened brown; tergal discs with regular granular microreticulation, weakly shining, essentially impunctate, with very weak and sparse punctures disappearing into background sculpture (Fig. 4F). Terga with scattered short white hairs, not forming apical hairbands. Apical fringe of T5 and hairs flanking pygidial plate orange. Pygidial plate large, broadly rounded triangular, lateral margin weakly raised and impunctate, internal surface densely punctate, punctures separated by < 0.5 puncture diameters.

Male. Unknown.

Etymology. The species is named after Daniele Baiocchi who has collected insects across much of the Mediterranean basin and Middle East, and who led bee collection during the expedition to Duhok Governorate in May 2023.

Distribution. Iraq (Kurdistan region).

11. *Andrena (Plastandrena) bimaculata* (Kirby, 1802)

Literature records. Gusenleitner and Schwarz (2002: dot map 55); Scheuchl and Willner (2016); Ascher and Pickering (2023).

Remarks. We have not examined any specimens of this species, but the presence of this species in Iraq is highly plausible based on its known global distribution, the map records indicated by Gusenleitner and Schwarz (2002), and the presence of this species in neighbouring Turkey and Iran. We note however that the species concept of *A. bimaculata* is unclear, and all Old World members of the subgenus *Plastandrena* require revision using molecular markers. It should therefore be considered in a sensu lato at the present time.

Distribution. In a sensu lato, West and Central Palearctic to Mongolia (Gusenleitner and Schwarz 2002; Ascher and Pickering 2023).

12. *Andrena (Cryptandrena) brumanensis* Friese, 1899 *

Material examined. IRAQ: Duhok, Bessre [Besereh], Bablo, 36.8675°N, 43.1206°E, 1065 m, 5–6.v.2023, 1♀, leg. D. Baiocchi, MSVI; Duhok, E Mangesh, 37.0230°N, 43.1505°E, 1046 m, 8.v.2023, 1♂, leg. D. Baiocchi, MSVI.

Distribution. Southern Europe to Turkey and the Near East, including Iraq* and Iran (Gusenleitner and Schwarz 2002; Wood and Monfared 2022).

13. *Andrena (Truncandrena) canaeae* Strand, 1915 *

Material examined. IRAQ: Duhok, Mt. Gara [S of Sarsing], 37.0158°N, 43.3506°E, 1912 m, 11.v.2023, 1♂, leg. D. Baiocchi, MSVI/TJWC.

Distribution. Greece, Turkey, Cyprus, Syria, Iraq* (Gusenleitner and Schwarz 2002).

14. *Andrena (Micrandrena) cedricola* Wood, 2020 *

Material examined. IRAQ: Duhok, Mt. Gara [S of Sarsing], 37.0158°N, 43.3506°E, 1912 m, 11.v.2023, 2♂, leg. D. Baiocchi, MSVI/TJWC.

Distribution. Israel, Lebanon, Syria, Turkey, Iraq* (Pisanty et al. 2022a).

15. *Andrena (Chlorandrena) cinereophila* Warncke, 1965 *

Material examined. IRAQ: Duhok, Mt. Gara [S of Sarsing], 37.0158°N, 43.3506°E, 1912 m, 11.v.2023, 1♂, leg. D. Baiocchi, MSVI; **LEBANON:** Balbek-Hermel, Sefri, Haouch Snaid, AUB farm, 33.9244°N, 36.0754°E, 1000 m, 6.iv.2023, 11♀, leg. T.J. Wood, TJWC; Beqaa, Anjar, 1 km E, Armenian Cemetery, 33.7372°N, 35.9503°E, 900 m, 7.iv.2023, 1♀, leg. T.J. Wood, TJWC; Beqaa, Beqaa valley, Mansourah, Aammiq wetland preserve, 33.7321°N, 35.7853°E, 850 m, 3.iv.2023, 1♂, leg. T.J. Wood, TJWC; Beqaa, Beqaa valley, Qaraoun dam, 33.5483°N, 35.6851°E, 850 m, 6.iv.2023, 3♂, leg. T.J. Wood, TJWC.

Distribution. East Mediterranean to Central Asia, including Lebanon* and Iraq* (Gusenleitner and Schwarz 2002; Osytsnjuk et al. 2005; Wood and Monfared 2022).

16. *Andrena (Cordandrena) cordialis* Morawitz, 1878

Literature records. Morice (1921b); Derwesh (1965), mentioning “Survey of Iraq Fauna 1915–1919”; Augul (2018).

Material examined. IRAQ: Baiji [Saladin Governorate, 35.0299°N, 43.4489°E], 1–31.iii.1986, 1 ♀, leg. M. Carl, OÖLM; Baiji, 1–30.iv.1986, 2 ♂, leg. M. Carl, OÖLM.

Remarks. *Andrena cordialis* is part of a tricky group of species to identify, with the species *A. cypria* Pittioni, 1950 and *A. torda* Warncke, 1965 described after the 1915–1919 survey was conducted. Morice (1921b) specifically mentioned the species from Basrah in southern Iraq. Based on our examined material, we can confidently confirm both *A. cordialis* and *A. torda* from Iraq, and so the record of Morice (1921b) is considered plausible. We note here that the “Survey of Iraq Fauna 1915–1919” is represented by the publications of Morice (1921a; 1921b), only Morice (1921b) dealing with the *Andrena* fauna.

Distribution. Eastern Europe through Turkey and the Caucasus to Central Asia, including Iraq and Iran (Gusenleitner and Schwarz 2002; Osytsnjuk et al. 2005; Wood and Monfared 2022)

17. *Andrena (Poecilandrena) crassana* Warncke, 1965 *

Material examined. IRAQ: Duhok, Mt. Gara [S of Sarsing], 37.0158°N, 43.3506°E, 1912 m, 11.v.2023, 2 ♂, leg. D. Baiocchi, MSVI; **LEBANON:** Beqaa, Beqaa valley, Qaraoun, 3.5 km W of Madjal Balhis, 33.5377°N, 35.7038°E, 900 m, 4.iv.2023, 1 ♂, leg. T.J. Wood, TJWC.

Remarks. Specimens from Iraq nominally belong to *A. crassana* s. str. And the specimen from Lebanon nominally belongs to ssp. *Inka* Warncke, 1969 (Levant). It is not clear if *inka* merits subspecific status.

Distribution. *Andrena crassana* sensu lato has a distribution of North Macedonia, Greece, Turkey, Cyprus, Israel and West Bank, Lebanon*, Jordan, Syria, Iraq*, Iran (Gusenleitner and Schwarz 2002; Pisanty et al. 2018; Wood and Monfared 2022).

18. *Andrena (Aciandrena) curviociput* Pisanty & Wood, 2022 *

Material examined. IRAQ: Duhok, Mt. Gara [S of Sarsing], 37.0158°N, 43.3506°E, 1912 m, 11.v.2023, 12 ♂, leg. D. Baiocchi, MSVI/TJWC.

Remarks. Known from the Tanin-Tanin pass in south-eastern Turkey (Pisanty et al. 2022a), so its presence in northern Iraq was expected.

Distribution. Israel, Lebanon, Turkey, Iraq* (Pisanty et al. 2022a).

19. *Andrena* (incertae sedis) *discordia* Wood, 2023 *

Material examined. IRAQ: Duhok, Mt. Gara [S of Sarsing], 37.0158°N, 43.3506°E, 1912 m, 11.v.2023, 1♂, leg. D. Baiocchi, MSVI.

Distribution. Turkey, Iraq*, Iran (Wood 2023).

20. *Andrena* (*Aciandrena*) *duhokensis* Wood, sp. nov. *

<https://zoobank.org/3C35CEE1-7063-4C23-8577-1BA21A6CCAFE>

Material examined. Holotype: IRAQ: Duhok, Mt. Gara [S of Sarsing], 37.0158°N, 43.3506°E, 1912 m, 11.v.2023, 1♂, leg. D. Baiocchi, RMNH. **Paratypes: IRAQ:** Same information as holotype, 7♂, MSVI/RMNH/TJWC/DUMAI.

Diagnosis. *Andrena duhokensis* can be recognised as part of the subgenus *Aciandrena* Warncke, 1968 due to the small body size, dark integument with the exception of the pale-marked clypeus (Fig. 5A), finely shagreened propodeal triangle (Fig. 5D; without lateral or basal rugae), more or less impunctate terga (Fig. 5E), strongly antefurcal nervulus, and typical simple genital capsule (Fig. 5F; e.g., without completely reduced gonocoxae as in most members of the subgenus *Graecandrena* Warncke, 1968). Identification of *Aciandrena* species is extremely challenging due to the large number of often cryptic species found in dry environments, their often very local distributions, and lack of good characters in the female sex. Fortunately, males are a little easier to identify through the examination of the genital capsule. *Andrena duhokensis* can be initially recognised due to its almost entirely yellow-marked clypeus (with the exception of two small black marks [sometimes absent] and some narrow black areas at the edges of the clypeus), placing it closest to *A. tenuis* Morawitz, 1877 which was described from the Caucasus. However, *A. duhokensis* can be recognised due to the unique form of the genital capsule which has the gonocoxal teeth with their apexes strongly truncate and asymmetrical so that they diverge from each other (Fig. 5F); in other words, they form somewhat pointed but flattened and diverging teeth. In comparative species of *Aciandrena*, the gonocoxal teeth are either strongly produced and truncate (e.g., *A. diminuta* Wood 2022 from Iran), produced into narrowly pointed or more broadly rounded teeth (e.g., *A. tenuis* from Turkey and the Caucasus, *A. aciculata* Morawitz, 1886 from Europe to the Caucasus), or reduced and truncate but never forming diverging teeth (e.g., *A. judaea* Scheuchl & Pisanty, 2016 from Israel). In this context, *A. duhokensis* can be separated from all similar species.

Description. Female. Unknown.

Male. Body length 4–5 mm (Fig. 5A). **Head:** Dark, 1.25× wider than long (Fig. 5B). Clypeus weakly domed, predominantly yellow or pale yellow marked with exception of two black dots (sometimes absent) and narrow black markings laterally; surface finely shagreened and weakly shining, irregularly punctate, punctures separated by 1–3 puncture diameters. Process of labrum small, rounded trapezoidal, 2× wider than long, apical margin weakly emarginate. Gena subequal to diameter of compound eye; ocelloccipital distance equals diameter of lateral ocellus. Head covered with dirty white hairs, none equalling length of scape. Antennae basally dark, A4–13 ventrally lightened by presence of grey scales; A3 exceeding length of A4, shorter than A4+5; A4 sub-square, shorter than long, A5 quadrate, as long as broad.




Figure 5. *Andrena (Aciandrena) duhokensis* sp. nov. male **A** habitus, lateral view **B** face, frontal view **C** scutum, dorsal view **D** propodeum, dorsal view **E** terga, dorsal view **F** genital capsule, dorsal view.

Mesosoma: Scutum and scutellum with fine granular shagreen, weakly shining, irregularly punctate, punctures separated by 1–4 puncture diameters (Fig. 5C). Mesepisternum and dorsolateral parts of propodeum with regular but large microreticulation, dull to weakly shining; propodeal triangle large, indicated by change in surface sculpture, internal surface with fine granular microreticulation, in some parts with weakly raised network of carinae between microreticulation, surface dull (Fig. 5D). Mesosoma with sparse short whitish hairs. Legs predominantly dark, apical tarsal segments slightly lightened

orange, pubescence white. Hind tarsal claws with strong inner tooth. Wings hyaline, stigma, and venation pale orange, nervulus very strongly antefurcal.

Metasoma: Terga predominantly dark, tergal margins weakly but distinctly depressed, broadly lightened hyaline yellow-brown (Fig. 5E). Tergal discs finely microreticulate to shagreened, weakly shining, surface essentially impunctate, with scattered obscure punctures disappearing into underlying sculpture. Terga with sparse scattered short hairs, not forming hairbands. T6 and T7 with long light brown hairs overlying pseudopygidial plate of T7. S8 narrow, apex slightly broadened like a fish-tail, truncate, ventral surface with dense lateral fan of brown hairs. Genital capsule compact, gonocoxae produced into apical teeth, teeth strongly truncate, apices angled and diverging from each other to form angular anteriorly projecting teeth (Fig. 5F). Gonostyli flattened and spatulate, internal margins raised, forming slight bump on inner margin basally. Penis valves moderate, occupying $\frac{1}{2}$ space between gonostyli basally, strongly tapering apically.

Etymology. The name is derived from the city of Duhok which gives its name to the province in which these specimens were collected.

Distribution. Iraq (Kurdistan region).

21. *Andrena (Micrandrena) elam* Wood, 2022 *

Material examined. IRAQ: Duhok, Mt. Gara [S of Sarsing], 37.0158°N, 43.3506°E, 1912 m, 11.v.2023, 23♂, 1♀, leg. D. Baiocchi, MSVI/TJWC.

Remarks and diagnosis. Wood in Wood and Monfared (2022) described *A. elam* in the female sex from western and southern Iran, and Wood (2023) reported further specimens from southern and south-eastern Turkey. Numerous male specimens from northern Iraq are now available. They can be recognised as *Micrandrena* due to the small body size, dark integument (including clypeus Fig. 6B), entirely rugose propodeal triangle (Fig. 6D), and normal genital capsule. As in the female sex, they can be separated from other *Micrandrena* species by the combination of smooth and shining scutum and scutellum with moderately dense punctures (Fig. 6C), the tergal discs with extremely strong and dense reticulation (Fig. 6E), this reticulation becoming much weaker on the marginal areas, with punctures visible on the discs of T3–4 (though comparatively weaker and less visible than in the female), and by the tergal margins becoming progressively more strongly depressed. Additionally, the genital capsule is distinctive (Fig. 6F), somewhat elongate, with the gonocoxae produced into weak angular teeth, with thickened gonostyli with a weak rounded bump on their inner margins and outer surface lightened and covered with short golden hairs. It resembles the form of two former *Fumandrena* (= *Micrandrena*) species (*A. fabrella* Pérez, 1895 (western Mediterranean) and *A. tomora* Warncke, 1975 (eastern Mediterranean)), but these have the gonostyli apically flattened, not thickened. It is very close to *A. subviridula* Wood, 2022 (northern Iran, see illustrations in Wood and Monfared 2022), but can be separated by the gonocoxae produced into apical points (in *A. subviridula* with the gonocoxae apically truncate), by the weaker bump on the inner margins of the gonostyli (in *A. subviridula* with this bump comparatively more pronounced), and by the more extensively shiny scutum which lacks subtle greasy green-metallic reflections (in *A. subviridula* with the scutum more extensively shagreened anteriorly and laterally).

Description. Male. Body length: 6 mm (Fig. 6A). **Head:** Dark, 1.45× wider than long (Fig. 6B). Clypeus domed, densely punctate, punctures separated by < 0.5–1 puncture diameter, surface with weakly raised latitudinal striations between punctures, surface weakly shining. Process of labrum trapezoidal, 2× wider than long, anterior margin weakly emarginate, surface smooth and shining. Gena equalling diameter of compound eye; ocellocipital distance equaling diameter of lateral ocellus. Head with sparse whitish hairs, none equalling length of scape; clypeus in fresh individuals with moderate “beard” of ventrally projecting pale hairs. Antennae dark, A5–13 ventrally lightened by presence of silver-grey scales; A3 exceeding length of A4, shorter than A4+5.

Mesosoma: Scutum and scutellum polished and shining over almost entire surface, scutum weakly shagreened anteriorly; irregularly punctate, punctures separated by 1–3 puncture diameters (Fig. 6C). Pronotum rounded. Mesepisternum microreticulate, dull. Dorsolateral parts of propodeum microreticulate, dull, sculpture overlain by network of irregular raised rugosity; propodeal triangle laterally defined by irregular carinae, internal surface densely covered with network of evenly-spaced rugae, propodeal triangle thus well-differentiated from dorsolateral parts of propodeum (Fig. 6D). Mesosoma covered with sparse light brown hairs, none equalling length of scape. Legs dark, apical tarsal segments paler dark brown, pubescence whitish. Hind tarsal claws with inner tooth. Wings hyaline, stigma dark brown, venation orange-brown, nervulus interstitial.

Metasoma: Tergal discs dark, marginal areas with apical rim narrowly lightened brown (Fig. 6E). Tergal discs strongly reticulate, reticulation strongest on T1, surface dull, becoming progressively weaker on subsequent terga, discs of T5 and T6 weakly shining; marginal areas shagreened, weakly shining. Tergal discs obscurely punctate, T1 almost impunctate, punctures progressively more visible on subsequent terga, punctures on disc of T4 separated by 1–2 puncture diameters. Tergal discs with scattered pale hairs, not forming hairbands. T6 and T7 with light brown hairs overlying pseudopygidial plate. S8 columnar, apical margin truncate, ventral surface with sparse short brown hairs. Genital capsule somewhat elongate, gonocoxae with apical margins produced into slightly projecting narrow apical teeth (Fig. 6F). Gonostyli robust, thickened, with weak bump on internal margins basally, apically weakly flattened, apical ½ lightened hyaline yellowish. Penis valves narrow, occupying ½ space between gonostyli, uniformly narrowing towards apex.

Distribution. Southern and south-eastern Turkey, northern Iraq*, and western and southern Iran (Wood and Monfared 2022; Wood 2023).

22. *Andrena (Lepidandrena) elisaria* Gusenleitner, 1998 *

Material examined. IRAQ: Duhok, Mt. Gara [S of Sarsing], 37.0158°N, 43.3506°E, 1912 m, 11.v.2023, 1♂, leg. D. Baiocchi, MSVI.

Distribution. Israel, Turkey, Iraq*, Iran (Gusenleitner 1998; Pisanty et al. 2018; Wood and Monfared 2022).

23. *Andrena (incertae sedis) euzona* Pérez, 1895

Literature records. Wood and Monfared (2022).



Figure 6. *Andrena (Micrandrena) elam* Wood, 2022 male **A** habitus, lateral view **B** face, frontal view **C** scutum, dorsal view **D** propodeum, dorsal view **E** terga, dorsal view **F** genital capsule, dorsal view.

Distribution. Morocco, Algeria, Tunisia, Libya, Egypt, Israel and West Bank, Jordan, Syria, Iraq, Iran (Gusenleitner and Schwarz 2002; Wood and Monfared 2022).

24. *Andrena (Plastandrena) eversmanni* Radoszkowski, 1867

Literature records. Wood and Monfared (2022).

Distribution. Turkey, Armenia, Iraq, Iran, Russia (European part), Turkmenistan, Uzbekistan, Kazakhstan, Tajikistan, Pakistan, Mongolia, China (Wood and Monfared 2022; Ascher and Pickering 2023).

25. *Andrena (Hoplendrena) ferox* Smith, 1847 *

Material examined. **IRAQ:** Duhok, Mt. Gara [S of Sarsing], 37.0158°N, 43.3506°E, 1912 m, 11.v.2023, 1♀, leg. D. Baiocchi, MSVI; **LEBANON:** Beqaa, Beqaa valley, Mansourah, Aammiq wetland preserve, 33.7321°N, 35.7853°E, 850 m, 3.iv.2023, 1♀, leg. T.J. Wood, TJWC.

Distribution. West Palaearctic to the Middle East, including Lebanon*, Iraq*, Iran (Gusenleitner and Schwarz 2002; Scheuchl and Willner 2016; Wood and Monfared 2022).

26. *Andrena (Melandrena) flavipes* Panzer, 1799

Literature records. Morice (1921b); Derwesh (1965), mentioning “Survey of Iraq Fauna 1915–1919”; Gusenleitner and Schwarz (2002: dot map 161); Scheuchl and Willner (2016); Augul (2018); Ascher and Pickering (2023).

Remarks. We have not examined any specimens of this species, but the presence of this species in Iraq is extremely plausible based on its known global distribution, its abundance in a wide variety of habitats, the map records indicated by Gusenleitner and Schwarz (2002), and the presence of this species in neighbouring Turkey and Iran.

Distribution. West and Central Palearctic (Gusenleitner and Schwarz 2002; Ascher and Pickering 2023).

27. *Andrena (Bryandrena) florea* Fabricius, 1793 *

Material examined. **IRAQ:** Duhok, Mt. Gara [S of Sarsing], 37.0158°N, 43.3506°E, 1912 m, 11.v.2023, 1♂, leg. D. Baiocchi, MSVI.

Distribution. West Palaearctic, from Morocco to Iraq*, Iran, Turkmenistan, and the Ural mountains (Gusenleitner and Schwarz 2002).

28. *Andrena (Ulandrena) fulvitorsis* Brullé, 1832

Literature records. Gusenleitner and Schwarz (2002: dot map 177); Ascher and Pickering (2023).

Remarks. We have not examined any specimens of this species, but the presence of this species in Iraq is plausible based on its known global distribution, the map records indicated by Gusenleitner and Schwarz (2002), and the presence of this species in neighbouring Turkey.

Distribution. East Mediterranean, from Italy to Crimea, Turkey, the Levant, Iraq (Gusenleitner and Schwarz 2002).

29. *Andrena (incertae sedis) garrula* Warncke, 1965 *

Material examined. IRAQ: Duhok, Mt. Gara [S of Sarsing], 37.0158°N, 43.3506°E, 1912 m, 11.v.2023, 1♀, leg. D. Baiocchi, MSVI.

Remarks. This specimen appears to be *A. garrula* s. str. as opposed to the Levantine ssp. *lomvia* Warncke, 1969; the difference between the two subspecies is difficult to distinguish in the female sex.

Distribution. Bulgaria, Turkey, Israel and West Bank, Lebanon, Jordan, Syria, Iraq*, Iran (Gusenleitner and Schwarz 2002; Wood et al. 2020; Wood and Monfared 2022).

30. *Andrena (Melandrena) grandilabris* Pérez, 1903 *

Material examined. IRAQ: Duhok, Mt. Gara [S of Sarsing], 37.0158°N, 43.3506°E, 1912 m, 11.v.2023, 4♂, leg. D. Baiocchi, MSVI.

Remarks. Turkey, Iraq*, Iran (Wood and Monfared 2022).

31. *Andrena (Euandrena) hermonella* Scheuchl & Pisanty, 2016 *

Material examined. IRAQ: Duhok, Mt. Gara [S of Sarsing], 37.0158°N, 43.3506°E, 1912 m, 11.v.2023, 2♀, leg. D. Baiocchi, MSVI.

Remarks. The original female specimens described by Pisanty et al. (2016) actually belonged to *A. gageae* Wood & Pisanty, 2022 (see Pisanty et al. 2022a). The true female of *A. hermonella* is being described from south-eastern Turkey (Wood, in press). This extends the range from south-eastern Turkey into northern Iraq.

Distribution. Israel, Turkey, Iraq* (Pisanty et al. 2016; Wood, unpublished data).

32. *Andrena (incertae sedis) hosseiniiae* Wood & Monfared, 2022 *

Material examined. IRAQ: Duhok, Mt. Gara [S of Sarsing], 37.0158°N, 43.3506°E, 1912 m, 11.v.2023, 1♀, leg. D. Baiocchi, MSVI.

Remarks. The first record of this species outside of Iran, with all previous records coming from close to Yasuj (Kohgiluyeh and Boyer-Ahmad Province) in southern Iran.

Distribution. Iraq* and Iran (Wood and Monfared 2022).

33. *Andrena (Poecilandrena) hybrida* Warncke, 1975 *

Material examined. IRAQ: Duhok, Mt. Gara [S of Sarsing], 37.0158°N, 43.3506°E, 1912 m, 11.v.2023, 1♀, leg. D. Baiocchi, MSVI.

Remarks. Multiple subspecies are described for *A. hybrida*; their statuses require further study, but this is limited by the scarcity of material. Iraqi material conforms to *A. hybrida* s. str.

Distribution. Ukraine, Russia (European part), Turkey, Iraq*, Iran (Gusenleitner and Schwarz 2002; Wood and Monfared 2022).

34. *Andrena (Graecandrena) hyemala* Warncke, 1973 *

Material examined. IRAQ: Duhok, Mt. Gara [S of Sarsing], 37.0158°N, 43.3506°E, 1912 m, 11.v.2023, 7♂, leg. D. Baiocchi, MSVI; Mosul, edges of a river, 7.iv.1988, 3♀, leg. Olejníček, OÖLM.

Remarks. The specimens presented here appear to be *A. hyemala* s. str. The statuses of the subspecies *repressa* Warncke, 1975 (Levant) and *kushika* Osytshnjuk, 1994 (Central Asia) require further study.

Distribution. *Andrena hyemala* sensu lato is distributed from south-eastern Europe through the Middle East (including Iraq* and Iran) to Central Asia (Gusenleitner and Schwarz 2002; Osytshnjuk et al. 2008; Wood and Monfared 2022).

35. *Andrena (Ulandrena) krieckbaumeri* Schmiedeknecht, 1883 *

Material examined. IRAQ: Duhok, E Mangesh, 37.0230°N, 43.1505°E, 1046 m, 8.v.2023, 1♂, leg. D. Baiocchi, MSVI; **SYRIA:** Apamea, 65 km NW Hama, 270 m, 18.iv.1992, 1♂, leg. K. Warncke, OÖLM; **TURKEY:** Siirt, 5 km E Eruh, 1000 m, 26.v.1983, 1♂, leg. K. Warncke, OÖLM.

Remarks. The finding of *A. krieckbaumeri* in Iraq is notable. One of the most common and abundant *Andrena* species in the southern Balkans, *A. krieckbaumeri* is much less common in Turkey where it is largely restricted to western Turkey (see distribution map of Gusenleitner and Schwarz 2002). Examination of undetermined material in the OÖLM collection produced specimens from northern Syria and a specimen from near to Siirt in eastern Turkey. To our knowledge, this is the most easterly specimen known from Turkey. In this context, the presence of *A. krieckbaumeri* in northern Iraq represents less of an outlier, though it is clear that *A. krieckbaumeri* has a population centre in the Balkan Peninsula.

Distribution. Europe from Italy to Turkey, Syria*, and Iraq* (Gusenleitner and Schwarz 2002).

36. *Andrena (Poecilandrena) laticeps* Morawitz, 1878 *

Material examined. IRAQ: Duhok, Mt. Gara [S of Sarsing], 37.0158°N, 43.3506°E, 1912 m, 11.v.2023, 3♂, leg. D. Baiocchi, MSVI.

Distribution. Turkey, Georgia, Armenia, Iraq*, Iran (Wood and Monfared 2022; Ascher and Pickering 2023).

37. *Andrena (Melandrena) limata* Smith, 1853 *

Material examined. IRAQ: Baiji [Saladin Governorate, 35.0299°N, 43.4489°E], 1–31.iii.1986, 3♀, leg. M. Carl, OÖLM.

Distribution. West and Central Palearctic including Iraq* and Iran (Gusenleitner and Schwarz 2002; Osytshnjuk et al. 2008; Wood and Monfared 2022).

38. *Andrena (Micrandrena) luscinia* Warncke, 1975 *

Material examined. **IRAQ:** Duhok, Mt. Gara [S of Sarsing], 37.0158°N, 43.3506°E, 1912 m, 11.v.2023, 1♀, leg. D. Baiocchi, MSVI; **LEBANON:** Beqaa, Rachaiya, 5 km S, Mount Hermon nature reserve, 33.4586°N, 35.8395°E, 1500 m, 8.iv.2023, 2♀, leg. T.J. Wood, TJWC.

Distribution. Israel, Lebanon*, Turkey, Iraq*, Iran (Pisanty et al. 2018; Wood and Monfared 2022).

39. *Andrena (incertae sedis) monacha* Warncke, 1965 *

Material examined. **IRAQ:** Duhok, Mt. Gara [S of Sarsing], 37.0158°N, 43.3506°E, 1912 m, 11.v.2023, 1♂, leg. D. Baiocchi, MSVI.

Distribution. Greece, Turkey, Cyprus, Lebanon, Syria, Iraq*, Iran (Gusenleitner and Schwarz 2002; Wood et al. 2020; Wood and Monfared 2022).

40. *Andrena (Melandrena) morio* Brullé, 1832

Literature records. Khalaf and Al-Omar (1974); Gusenleitner and Schwarz (2002: dot map 310); Scheuchl and Willner (2016); Augul (2018).

Remarks. We have not examined any specimens of this species, but the presence of this species in Iraq is plausible based on its known global distribution, the map records indicated by Gusenleitner and Schwarz (2002), and the presence of this species in neighbouring Turkey and Iran.

Distribution. West and Central Palaearctic (Gusenleitner and Schwarz 2002; Osytshnjuk et al. 2008; Wood and Monfared 2022).

41. *Andrena (Melandrena) nitidemula* Scheuchl & Hazir, 2012 *

Material examined. **IRAQ:** Duhok, Mt. Gara [S of Sarsing], 37.0158°N, 43.3506°E, 1912 m, 11.v.2023, 1♀, leg. D. Baiocchi, MSVI.

Distribution. Greece, Turkey, Syria, Georgia, Armenia, Iraq*, Iran (Wood and Monfared 2022).

42. *Andrena (Micrandrena) obsidiana* Wood, 2022 *

Material examined. **IRAQ:** Duhok, Mt. Gara [S of Sarsing], 37.0158°N, 43.3506°E, 1912 m, 11.v.2023, 15♂, 4♀, leg. D. Baiocchi, MSVI/TJWC; **TURKEY:** Hakkâri, pass E of Uludere, 6.vi.1977, 1♂, 2♀, leg. K. Warncke, OÖLM/TJWC; Hakkâri, Tanin-Tanin-Pass, 2500 m, 2.vi.1980, 3♂, 1♀, leg. K. Warncke, OÖLM.

Remarks and diagnosis. Wood and Monfared (2022) described *A. obsidiana* in the female sex from southern and south-eastern Turkey and western and southern Iran. Numerous male specimens from northern Iraq are now available,

as well as some additional specimens from south-eastern Turkey that were not available for study at the time of the original description. One of these *A. obsidiana* specimens from Turkey (a female) was separated by Warncke and labelled as "*A. ferulella* spec. nov.", but this name was never published.

Andrena obsidiana can be recognised as *Micrandrena* due to the small body size, dark integument (including clypeus Fig. 7B), entirely rugose propodeal triangle (Fig. 7D), and normal genital capsule (Fig. 7F). Due to the smooth and shining scutum and scutellum with moderately dense punctures (Fig. 7C) and the genital capsule (gonocoxae produced into weak angular teeth, with thickened gonostyli with very small rounded bump on their inner margins, and outer surface lightened and covered with short golden hairs) it is close to *A. elam* and *A. subviridula*, but it can instantly be separated from them by the sculpture of the terga which is almost completely smooth and shining (Fig. 7E), with only superficial shagreenation at the base of the tergal discs. There are also differences in the genital capsule, with that of *A. obsidiana* being comparatively more elongate, with the bump on the inner margin of the gonostyli reduced to a relatively tiny projection.

Description. Male. Body length: 5–6 mm (Fig. 7A). **Head:** Dark, 1.25× wider than long (Fig. 7B). Clypeus domed, densely punctate, punctures separated by < 0.5–1 puncture diameter, surface weakly shining. Process of labrum trapezoidal, 2× wider than long, anterior margin weakly emarginate, surface weakly shining. Gena equalling diameter of compound eye; ocelloccipital distance subequal to diameter of lateral ocellus. Head with sparse whitish hairs, none equalling length of scape; clypeus in fresh individuals with moderate "beard" of ventrally projecting pale hairs. Antennae basally dark, A5–13 ventrally lightened by presence of brownish-grey scales; A3 exceeding length of A4, slightly shorter than A4+5.

Mesosoma: Scutum and scutellum polished and shining over almost entire surface, scutum weakly shagreened anteriorly; irregularly punctate, punctures separated by 1–3 puncture diameters, slightly denser on scutum (Fig. 7C). Pronotum rounded. Mesepisternum microreticulate, dull. Dorsolateral parts of propodeum microreticulate, dull, sculpture overlain by dense network of raised rugosity; propodeal triangle poorly defined laterally, without clear linear carinae, internal surface densely covered with dense network of rugae, propodeal triangle thus poorly differentiated from dorsolateral parts of propodeum (Fig. 7D). Mesosoma covered with sparse whitish to light brown hairs, none equalling length of scape. Legs dark, apical tarsal segments paler dark brown, pubescence whitish. Hind tarsal claws with inner tooth. Wings hyaline, stigma dark brown, venation orange-brown, nervulus weakly to strongly antefurcal.

Metasoma: Tergal discs dark, marginal areas with apical rim obscurely lightened dark brown (Fig. 7E). Tergal discs with weak sculpture, T1 polished and shining, base of remaining terga with fine shagreen, otherwise smooth and shining. Terga deeply punctate, T1 with punctures separated by 1–2 puncture diameters, remaining tergal discs with punctures separated by 1 puncture diameter; punctures only partially extending onto marginal areas, becoming weaker on apical terga, marginal area of T5 almost impunctate. Tergal margins progressively more strongly depressed, weakly on T1, strongly on T5. Tergal discs with scattered pale hairs, not forming hairbands. T6 and T7 with light brown hairs overlying pseudopygidial plate. S8 columnar, apical margin rounded, ventral surface with dense fan of short brown hairs. Genital



Figure 7. *Andrena (Micrandrena) obsidiana* Wood, 2022 male **A** habitus, lateral view **B** face, frontal view **C** scutum, dorsal view **D** propodeum, dorsal view **E** terga, dorsal view **F** genital capsule, dorsal view.

capsule somewhat elongate, gonocoxae with apical margins produced into slightly projecting narrow apical teeth (Fig. 7F). Gonostyli robust, thickened, with weak and obscure bump on internal margins basally, apical $\frac{1}{2}$ lightened hyaline yellowish. Penis valves narrow, occupying $\frac{1}{2}$ space between gonostyli, uniformly narrowing towards apex.

Distribution. Southern and south-eastern Turkey, northern Iraq*, and western and southern Iran (Wood and Monfared 2022).

43. *Andrena (Micrandrena) oediceps* Warncke, 1975 *

Material examined. IRAQ: Duhok, Mt. Gara [S of Sarsing], 37.0158°N, 43.3506°E, 1912 m, 11.v.2023, 2♂, leg. D. Baiocchi, MSVI;

Distribution. Greece, Turkey, Israel and West Bank, Lebanon, Jordan, Syria, Turkey, Iraq*, Iran (Gusenleitner and Schwarz 2002; Pisanty et al. 2018; Wood and Monfared 2022).

44. *Andrena (Pallandrena) pallidicincta* Brullé, 1832 *

Material examined. IRAQ: Duhok, Mt. Gara [S of Sarsing], 37.0158°N, 43.3506°E, 1912 m, 11.v.2023, 11♀, leg. D. Baiocchi, MSVI/TJWC.

Distribution. South-eastern Europe to Turkey, Lebanon, Israel, and Iraq* (Gusenleitner and Schwarz 2002; Wood et al. 2020).

45. *Andrena (Truncandrena) pareklisiae* Mavromoustakis, 1956 *

Material examined. IRAQ: Duhok, Mt. Gara [S of Sarsing], 37.0158°N, 43.3506°E, 1912 m, 11.v.2023, 6♀, leg. D. Baiocchi, MSVI/TJWC.

Distribution. Cyprus, Turkey, Lebanon, Syria, Iraq*, Iran (Wood 2023).

46. *Andrena (Truncandrena) rufomaculata* Friese, 1921 *

Material examined. IRAQ: Duhok, Mt. Gara [S of Sarsing], 37.0158°N, 43.3506°E, 1912 m, 11.v.2023, 1♀, leg. D. Baiocchi, MSVI.

Distribution. Turkey, Israel and West Bank, Lebanon, Jordan, Syria, Iraq*, Iran (Wood et al. 2020; Wood and Monfared 2022).

47. *Andrena (Suandrena) savignyi* Spinola, 1838

Literature records. Moalif (1994); Ascher and Pickering (2023).

Remarks. We have not examined any specimens of this species, but the listing is plausible due to the presence of multiple specimens of *A. savignyi* in western and southern Iran (Wood and Monfared 2022), and the preference of *A. savignyi* for dry desert and semi-desert habitats.

Distribution. West and Central Palaeartic (Wood and Monfared 2022; Ascher and Pickering 2023).

48. *Andrena (Micrandrena) sillata histrionica* Warncke, 1975 *

Material examined. IRAQ: Duhok, Mt. Gara [S of Sarsing], 37.0158°N, 43.3506°E, 1912 m, 11.v.2023, 4♂, 1♀, leg. D. Baiocchi, MSVI.

Distribution. Subspecies *histrionica* is found in Turkey, Iraq*, and Iran only (Wood and Monfared 2022). It may merit species status due to differences in the male genital capsule, but genetic data are required.

49. *Andrena (Euandrena) symphyti* Schmiedeknecht, 1883 *

Material examined. IRAQ: Duhok, Mt. Gara [S of Sarsing], 37.0158°N, 43.3506°E, 1912 m, 11.v.2023, 5♂, 1♀, leg. D. Baiocchi, MSVI.

Distribution. West Palaearctic, including Iraq* and Iran (Gusenleitner and Schwarz 2002; Wood and Monfared 2022).

50. *Andrena (Chlorandrena) tadauchii* Gusenleitner, 1998 *

Material examined. IRAQ: Duhok, E Mangesh, 37.0230°N, 43.1505°E, 1046 m, 8.v.2023, 1♀, leg. D. Baiocchi, MSVI; Duhok, Mt. Gara [S of Sarsing], 37.0158°N, 43.3506°E, 1912 m, 11.v.2023, 14♂, leg. D. Baiocchi, MSVI/TJWC; **LEBANON:** Balbek-Hermel, Sefri, Haouch Snaid, AUB farm, 33.9244°N, 36.0754°E, 1000 m, 6.iv.2023, 2♂, 20♀, leg. T.J. Wood, TJWC; Beqaa, Anjar, 1 km E, Armenian Cemetery, 33.7372°N, 35.9503°E, 900 m, 7.iv.2023, 1♀, leg. T.J. Wood, TJWC; Beqaa, Beqaa valley, Mansourah, Aammiaq wetland preserve, 33.7321°N, 35.7853°E, 850 m, 3.iv.2023, 1♀, leg. T.J. Wood, TJWC; Bekaa, Qob Elias valley, 33.7989°N, 35.8192°E, 900 m, 5.iv.2023, 1♀, leg. T.J. Wood, TJWC; Hrar-Akkar, 34.4572°N, 36.1228°E, 900 m, 17.iv.2021, 1♂, leg. A. Saab, TJWC.

Remarks. Confirmed as present in Lebanon after the unclear listing of Grace (2010).

Distribution. Israel and West Bank, Lebanon*, Syria, Turkey, Iraq* (Pisanty et al. 2022a).

51. *Andrena (Ulandrena) tadorna callida* Warncke, 1974 *

Material examined. IRAQ: Hatra [Nineveh Governorate, 35.5759°N, 42.7254°E], 6.iv.1988, 1♂, leg. Olejníček, OÖLM; **SAUDI ARABIA:** Abha, 2000 m, 31.iii.1980, 1♂, leg. K.M. Guichard, NHMUK; As Nimas [Al Namas], 2450 m, 3–4.iv.1980, 3♂, 3♀, leg. K.M. Guichard, NHMUK; **SYRIA:** 110 km E of Palmyra, 350 m, 21.iv.1992, 2♀, leg. K. Warncke, OÖLM; 30 km W Palmyra, 580 m, 23.iv.1992, 6♀, leg. M. Kraus and K. Warncke, OÖLM; Suweidaono, ENE 80km, 700 m, 27.iii.1988, 2♂, leg. M. Schwarz, TJWC.

Remarks. The status of *A. tadorna callida* is unclear and requires investigation; it may well merit species status due to consistent differences in the shape of the genital capsule (see comments in Gusenleitner and Schwarz 2002).

Distribution. *Andrena tadorna* sensu lato is found in Morocco, Algeria, Tunisia, Libya, Egypt, Israel, Jordan, Syria*, Saudi Arabia*, Iraq* Gusenleitner and Schwarz (2002).

52. *Andrena (Micrandrena) tkalcui* Gusenleitner & Schwarz, 2002 *

Material examined. IRAQ: Duhok, Bessre [Besereh], Bablo, 36.8675°N, 43.1206°E, 1065 m, 5–6.v.2023, 1♀, leg. D. Baiocchi, MSVI; Duhok, Mt. Gara [S of Sarsing], 37.0158°N, 43.3506°E, 1912 m, 11.v.2023, 1♀, leg. D. Baiocchi, MSVI.

Distribution. Israel and West Bank, Jordan, Syria, Turkey, Iraq*, Iran (Gusenleitner and Schwarz 2002; Wood and Monfared 2022).

53. *Andrena (Cordandrena) torda* Warncke, 1965 *

Material examined. IRAQ: Duhok, Mt. Gara [S of Sarsing], 37.0158°N, 43.3506°E, 1912 m, 11.v.2023, 6♂, leg. D. Baiocchi, MSVI/TJWC; **LEBANON:** Beqaa, Beqaa valley, Mansourah, Aammiq wetland preserve, 33.7321°N, 35.7853°E, 850 m, 3.iv.2023, 2♀, leg. T.J. Wood, TJWC; Beqaa, Beqaa valley, Qaraoun, 3.5 km W of Madjal Balhis, 33.5377°N, 35.7038°E, 900 m, 4–5.iv.2023, 2♀, leg. T.J. Wood, TJWC.

Remarks. Records from Lebanon, Jordan, and Syria reported by Wood et al. (2020) were incorrectly identified female specimens of *A. cypria* (see Wood and Monfared 2022), with females of *A. torda* and *A. cypria* being challenging to separate morphologically. The female Lebanese specimens presented here were confirmed as *A. torda* through DNA barcoding (BIN: [BOLD:AES5002](#)) as distinct from *A. cypria* (BIN: [BOLD:AFH0814](#)), this latter BIN containing a bar-coded male specimen of *A. cypria* which can be unambiguously recognised due to the combination of its antennal segment ratios and genital capsule.

Distribution. Greece, Turkey, Cyprus, Israel and West Bank, Lebanon*, Iraq*, Iran (Wood and Monfared 2022).

54. *Andrena (Notandrena) trimarginata* (Radoszkowski, 1886)

Literature records. Wood and Monfared (2022, as *A. zoster* Warncke, 1975).

Material examined. LEBANON: Balbek-Hermel, Sefri, Haouch Snaid, AUB farm, 33.9244°N, 36.0754°E, 1000 m, 6.iv.2023, 4♂, leg. T.J. Wood, TJWC.

Remarks. A taxonomic work will soon recognise *Halictus trimarginatus* Radoszkowski, 1886 (described from Turkmenistan) as the senior name of *A. zoster* (Wood, in press).

Distribution. Israel and West Bank, Syria, Lebanon*, Turkey, Azerbaijan, Iraq, Iran, Turkmenistan, Uzbekistan, Tajikistan (Osytshnjuk et al. 2005 as *A. sub-smaragdina* Osytshnjuk, 1984; Wood and Monfared 2022).

55. *Andrena (Aciandrena) turmalina* Pisanty & Wood, 2022 *

Material examined. IRAQ: Duhok, Bessre [Besereh], Bablo, 36.8675°N, 43.1206°E, 1065 m, 5–6.v.2023, 1♀, leg. D. Baiocchi, MSVI; Duhok, Mt. Gara [S of Sarsing], 37.0158°N, 43.3506°E, 1912 m, 11.v.2023, 4♂, 5♀, leg. D. Baiocchi, MSVI/TJWC; **LEBANON:** Beqaa, Anjar, 1 km E, reforestation area, 33.7311°N, 35.9478°E, 1000 m, 7.iv.2023, 1♀, leg. T.J. Wood, TJWC.

Remarks. As suspected by Pisanty et al. (2022a), *A. turmalina* is also present in Lebanon.

Distribution. Israel, Lebanon*, Turkey, Iraq*, Iran (Pisanty et al. 2022a).

56. *Andrena (Holandrena) variabilis* Smith, 1853

Andrena bakrajoensis Amin & Mawlood, 2019, syn. nov.

Literature records. Amin and Mawlood (2019, as *A. bakrajoensis*)

Remarks. Amin and Mawlood (2019) described *A. bakrajoensis* from Bakrajo in Iraqi Kurdistan. They diagnosed it against *A. (Simandrena) vetula* Lepeletier, 1841, but based on the description and the provided photographs it is clearly a member of the subgenus *Holandrena* Pérez, 1890 due to the predominantly declivous propodeum (almost without a dorsal horizontal area), strong tergal hair-bands, and compact body with rounded head. Based on the long ocelloccipital distance (3× the diameter of a lateral ocellus), it can only be a single species, *A. variabilis*. *Andrena bakrajoensis* is therefore synonymised with *A. variabilis*. The distribution maps of Gusenleitner and Schwarz (2002) do not positively indicate the presence of *A. variabilis* in Iraq, though some dots are present around the Tigris and Euphrates deltas in southern Iraq. However, *A. variabilis* is a widely distributed species in Turkey and Iran (Gusenleitner and Schwarz 2002; Wood and Monfared 2022), and so its presence in Iraq is expected and demonstrated through this synonymy.

57. *Andrena (Planiandrena) veterana* Pisanty, 2022 *

Material examined. IRAQ: Duhok, Mt. Gara [S of Sarsing], 37.0158°N, 43.3506°E, 1912 m, 11.v.2023, 1♀, leg. D. Baiocchi, MSVI; **LEBANON:** Beqaa, Rachaiya, 5 km S, Mount Hermon nature reserve, 33.4586°N, 35.8395°E, 1500 m, 8.iv.2023, 10♀, leg. T.J. Wood, TJWC.

Remarks. These records markedly expand the range of *A. veterana* which was described from Mount Hermon (Pisanty et al. 2022a).

Distribution. Israel, Lebanon*, Iraq* (Pisanty et al. 2022a).

58. *Andrena (Simandrena) vetula* Lepeletier, 1841

Literature records. Morice (1921b); Derwesh (1965), mentioning “Survey of Iraq Fauna 1915–1919”; Gusenleitner and Schwarz (2002: dot map 504); Augul (2018); Ascher and Pickering (2023).

Remarks. We have not examined any specimens of this species, but the presence of this species in Iraq is highly plausible based on its known global distribution, the map records indicated by Gusenleitner and Schwarz (2002), and the presence of this species in neighbouring Turkey and Iran.

Distribution. West and Central Palearctic (Gusenleitner and Schwarz 2002; Wood and Monfared 2022; Ascher and Pickering 2023).

59. *Andrena* (*Poecilandrena*) *viridescens* Viereck, 1916

Literature records. Morice (1921b); Derwesh (1965), mentioning “Survey of Iraq Fauna 1915–1919”, as *A. cyanescens* Nylander, 1852 nec. Haliday; Augul (2018).

Material examined. IRAQ: Duhok, Mt. Gara [S of Sarsing], 37.0158°N, 43.3506°E, 1912 m, 11.v.2023, 5♂, leg. D. Baiocchi, MSVI/TJWC.

Remarks. Members of the subgenus *Poecilandrena* Hedicke, 1933 are challenging to identify (e.g., Pisanty et al. 2018), but we can confirm the record of Morice (1921b) through the examination of newly collected material.

Distribution. Europe to Turkey, Iraq, Iran (Gusenleitner and Schwarz 2002; Wood and Monfared 2022).

Species excluded

Andrena (*Melandrena*) *sigiella* Gusenleitner, 1998

Literature records. Grace (2010); Augul (2018); Ascher and Pickering (2023).

Remarks. We have examined no specimens of this species from outside of the Levant (Israel and West Bank, Jordan, Lebanon, Syria; Wood et al. 2020). Neither Grace (2010) or Augul (2018) report precise specimen records and, pending further investigations, we exclude *A. sigiella* from the list of Iraqi *Andrena* as if truly present it would be a considerable range extension from the Levant.

Andrena (*Suandrena*) *leucocyanea* Pérez, 1895

Literature records. Dylewska (1983).

Remarks. Dylewska (1983: 22) revised members of the subgenus *Suandrena* Warncke, 1968, mentioning *A. leucocyanea* from Iraq from Abu Ghuraib [= Abu Ghraib] and Dewania [= Al Diwanayah]. Given the many taxonomic changes that have occurred in this subgenus since 1983 (e.g., Kratochwil 2021; Pisanty et al. 2022a) and its difficulty of identification, it is unclear what these specimens might actually be. Consequently, we exclude this species from the Iraqi fauna pending inspection of material in light of the currently accepted taxonomic framework for this subgenus.

Discussion

As a measure of its chronic lack of study, the updated faunal list presented here for Iraqi *Andrena* totals some 59 species, a tiny fraction of the 215 and 388 species known from neighbouring Iran and Turkey, respectively (Wood and Monfared 2022; Wood 2023; Wood, unpublished data), and even of the 154–166 species known from neighbouring Syria. Moreover, to the best of our knowledge, the current work represents the first country records for 42 of the 59 Iraqi *Andrena* species reported here. Indeed, the collection around Duhok in May 2023 produced 38 new country records just for *Andrena*, including the two new species for science. Of these 38 new country records, 12 of these species have been described since 2016, highlighting that not

only is Iraq understudied in general, but that it specifically hosts a special *Andrena* fauna containing range-restricted species that has received relatively little attention until recently.

The increase from 17 to 59 *Andrena* species represents a 347% increase in species richness. Should it be applied equally across the entire Iraqi bee fauna, the 101 species listed by Ascher and Pickering (2023) would become 350. Though this is a crude approach, this produced total is plausible and seems reasonable given the huge diversity of bee species known to occur in Turkey (1,786 species as listed by Ascher and Pickering 2023). If anything, it is likely to be an underestimate. Based on the initial determination of other Iraqi bee groups, we will soon report faunal increases of 380% for *Eucera* (Apidae), ~ 666–833% for osmiine bees (Megachilidae), and an estimated 430–530% increase for *Nomada* (Apidae) (M. Selis, unpublished data). When dealing with such a poorly studied but likely species-rich fauna, these massive increases indicate a consistent pattern. We hope that this first focused revision of Iraqi *Andrena* can serve as a model to promote further studies into the Iraqi bee fauna.

Acknowledgements

We thank Esther Ockermüller and Martin Schwarz (OÖLM) for their continued support at the invaluable Linz collection, as well as Joseph Monks (NHMUK). We thank Alireza Monfared, Christian Schmid-Egger, Christophe Praz, and Erwin Scheuchl for helpful comments which meaningfully improved the manuscript.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

No funding was reported.

Author contributions

TJW identified specimens, described new species, and lead writing of the manuscript; HRI and DB coordinated fieldwork, conducted fieldwork, and contributed to writing of the manuscript; MIH and TTB conducted fieldwork; MS coordinated and conducted specimen preparation and identification, and contributed to writing of the manuscript.

Author ORCIDs

Thomas J. Wood  <https://orcid.org/0000-0001-5653-224X>

Marco Selis  <https://orcid.org/0000-0002-8200-9441>

Data availability

All of the data that support the findings of this study are available in the main text.

References

- Amin HM, Mawlood NA (2019) A New Species of The Sand Bees, *Andrena* Fabricius, 1775 (Hymenoptera, Andrenidae) From Kurdistan Region-Iraq. *Journal of University of Garmin* 6(2): 343–347. <https://doi.org/10.24271/garmian.196228>
- Ascher JS, Pickering J (2023) Discover Life bee species guide and world checklist (Hymenoptera: Apoidea: Anthophila). https://www.discoverlife.org/mp/20q?guide=Apoidea_species&flags=HAS
- Augul RS (2018) Study on diversity of bees (Hymenoptera, Apoidea) from different regions of Iraq. *Bulletin of the Iraq Natural History Museum* 15(1): 57–75. <https://doi.org/10.26842/binhm.7.2018.15.1.0057>
- Bossert S, Wood TJ, Patiny S, Michez D, Almeida EAB, Minckley RL, Packer L, Neff JL, Copeland RS, Straka J, Pauly A, Griswold T, Brady SG, Danforth BN, Murray EA (2022) Phylogeny, biogeography and diversification of the mining bee family Andrenidae. *Systematic Entomology* 47(2): 283–302. <https://doi.org/10.1111/syen.12530>
- Derwesh AI (1965) A preliminary list of identified insects and arachnids of Iraq. *Director General Agriculture Research Projections Baghdad. Bulletin* 112: 121–123.
- Dylewska M (1983) *Andrena suerinensis* Friese und verwandte Arten (*suerinensis*- Untergruppe) (Hymenoptera, Apoidea, Andrenidae). *Entomologische Abhandlungen* 47: 15–34.
- GISGeography (2024) Iraq Satellite Map. <https://gisgeography.com/iraq-map/#Satellite-Map> [accessed 27 January 2024]
- Grace A (2010) *Introductory biogeography to bees of the Eastern Mediterranean and Near East*. Bexhill Museum Association, Bexhill, 283 pp.
- Gusenleitner F (1998) Neue westpaläarktische *Andrena*-Arten (Hymenoptera: Apoidea: Andreninae). *Entomofauna* 19: 109–144.
- Gusenleitner F, Schwarz M (2002) Weltweite Checkliste der Bienengattung *Andrena* mit Bemerkungen und Ergänzungen zu paläarktischen Arten (Hymenoptera, Apoidea, Andreninae, *Andrena*). *Entomofauna (Supplement 10)*: 1–1280.
- Hazir C, Keskin N, Scheuchl E (2014) Faunistic, geographical and biological contributions to the bee genus *Andrena* (Hymenoptera, Andrenidae, Andreninae) from Turkey. *Journal of Hymenoptera Research* 38: 59–133. <https://doi.org/10.3897/jhr.38.7288>
- Khalaf AN, Al-Omar MA (1974) A second list of insects of Iraq. *Biological Research Center Publication No. 2.*, Iraq Government Press, Baghdad, 41 pp.
- Khodarahmi Ghahnavieh R, Monfared A (2019) A survey of bees (Hymenoptera: Apoidea) from Isfahan Province, Iran. *Journal of Insect Biodiversity and Systematics* 5(3): 171–201. <https://doi.org/10.52547/jibs.5.3.171>
- Kratochwil A (2021) Taxonomic re-evaluation of *Andrena cyanomicans* Pérez, 1895, *A. fratella* Warncke, 1968, *A. maderensis* Warncke, 1969, *A. mirna* Warncke, 1969, *A. notata* Warncke, 1968, and *A. portosanctana* Warncke, 1969 (Hymenoptera, Anthophila). *Linzer Biologische Beiträge* 53: 631–698.
- Michener CD (2007) *The Bees of the World*. 2nd edn. Johns Hopkins University Press, Baltimore, 953 pp.
- Moalif AS (1994) The bee fauna of Iraq. *Melissa* 7: 5–6.
- Morice FD (1921a) Annotated lists of Aculeate Hymenoptera (except Heterogyna) and Chrysidids recently collected in Mesopotamia and North-west Persia. *Journal of the Bombay Natural History Society* 27: 816–828.
- Morice FD (1921b) Annotated lists of Aculeate Hymenoptera (except Heterogyna) and Chrysidids recently collected in Mesopotamia and North-west Persia. (Continued from p. 828, Vol. XXVII.). *Journal of the Bombay Natural History Society* 28: 192–203.

- Osytsnjuk AZ, Romasenko L, Banaszak J, Cierznik T (2005) Andreninae of the Central and Eastern Palaearctic. Part 1. Polish Entomological Monographs II. Polish Entomological Society, Poznań, Poland, 235 pp.
- Osytsnjuk AZ, Romasenko L, Banaszak J, Motyka E (2008) Andreninae of the Central and Eastern Palearctic. Part 2. Polish Entomological Monographs V. Polish Entomological Society, Poznań, Poland, 233 pp.
- Pisanty G, Scheuchl E, Dorchin N (2016) Eight new species of *Andrena* Fabricius (Hymenoptera: Apoidea: Andrenidae) from Israel – a Mediterranean hotspot for wild bees. *Zootaxa* 4189(3): 485–515. <https://doi.org/10.11646/zootaxa.4189.3.3>
- Pisanty G, Scheuchl E, Dorchin N (2018) Taxonomic review of the subgenus *Andrena* (*Poecilandrena*) (Hymenoptera: Andrenidae) in Israel and the Levant. *Zootaxa* 4374(2): 151–188. <https://doi.org/10.11646/zootaxa.4374.2.1>
- Pisanty G, Scheuchl E, Martin T, Cardinal S, Wood TJ (2022a) Twenty-five new species of mining bees (Hymenoptera: Andrenidae: *Andrena*) from Israel and the Levant. *Zootaxa* 5185(1): 1–109. <https://doi.org/10.11646/zootaxa.5185.1.1>
- Pisanty G, Richter R, Martin T, Dettman J, Cardinal S (2022b) Molecular phylogeny, historical biogeography and revised classification of andrenine bees (Hymenoptera: Andrenidae). *Molecular Phylogenetics and Evolution* 170: 107151. <https://doi.org/10.1016/j.ympev.2021.107151>
- Popov VV (1967) The bees (Hymenoptera, Apoidea) of Iran. *Trudy Zoologicheskogo Instituta, Leningrad* 43: 184–216. [In Russian.]
- Praz C, Genoud D, Vaucher K, Bénon D, Monks J, Wood TJ (2022) Unexpected levels of cryptic diversity in European bees of the genus *Andrena* subgenus *Taeniandrena* (Hymenoptera, Andrenidae): Implications for conservation. *Journal of Hymenoptera Research* 91: 375–428. <https://doi.org/10.3897/jhr.91.82761>
- Proshchalykin MY, Fateryga M, Astafurova YV (2023) Corrections and additions to the catalogue of the bees (Hymenoptera, Anthophila) of Russia. *ZooKeys* 1187: 301–339. <https://doi.org/10.3897/zookeys.1187.113240>
- Radchenko VG, Allahverdi M, Fekrat L (2021) Revision of the mining bee subgenus *Andrena* (*Longandrena*) (Hymenoptera: Apoidea: Andrenidae). *Zootaxa* 5032(4): 489–515. <https://doi.org/10.11646/zootaxa.5032.4.2>
- Scheuchl E, Gusenleitner F (2009) *Andrena* (*Margandrena*) *elsei* nov.sp., eine neue türkische Sandbienenart (Hymenoptera, Apidae, Andreninae). *Linzer Biologische Beiträge* 41: 947–962.
- Scheuchl E, Hazir C (2012) Description of three new *Andrena* species (Hymenoptera: Apoidea: Andrenidae) from Turkey. *The Florida Entomologist* 95(4): 831–838. <https://doi.org/10.1653/024.095.0403>
- Scheuchl E, Willner W (2016) Taschenlexikon der Wildbienen Mitteleuropas – Alle Arten im Porträt. Quelle and Meyer Verlag, Wiebelsheim, 917 pp.
- Schwenninger HR (2015) Revision of the Western Palaearctic species of the *Andrena taraxaci*-group with description of four new species (Hymenoptera: Andrenidae). *Stuttgarter Beiträge zur Naturkunde A, Neue Serie* 8: 251–270.
- Warncke K (1965) Beitrag zur Kenntnis der Bienengattung *Andrena* F. in der Türkei (Hymenoptera, Apoidea). *Mitteilungen der Münchner Entomologischen Gesellschaft* 55: 244–273.
- Warncke K (1969) A contribution on the knowledge of the genus *Andrena* (Apoidea) in Israel. *Israel Journal of Entomology* 4: 377–408.
- Warncke K (1974) Die Sandbienen der Türkei (Hymenoptera, Apoidea, *Andrena*), Teil A. *Mitteilungen der Münchner Entomologischen Gesellschaft* 64: 81–116.

- Warncke K (1975) Die Sandbienen der Türkei (Hymenoptera, Apoidea, *Andrena*), Teil B. Mitteilungen der Münchner Entomologischen Gesellschaft 65: 29–102.
- Wood TJ (2020) Description of a remarkable new *Andrena* species (Hymenoptera: Andrenidae) from Syria. Zoology in the Middle East 66(3): 262–268. <https://doi.org/10.1080/09397140.2020.1782580>
- Wood TJ (2023) New Old World *Andrena* species, with a focus on Turkey (Hymenoptera: Andrenidae). Zootaxa 5266(1): 1–72. <https://doi.org/10.11646/zootaxa.5266.1.1>
- Wood TJ (2024) New Asian *Andrena* species, with notes on the subgenus *Cnemidandrena* (Hymenoptera: Andrenidae). Zootaxa 5404(1): 167–188. <https://doi.org/10.11646/zootaxa.5404.1.11>
- Wood TJ (in press) Further revisions to the Palaearctic *Andrena* fauna (Hymenoptera: Andrenidae). Zootaxa. [in press]
- Wood TJ, Monfared A (2022) A revision of the *Andrena* (Hymenoptera: Andrenidae) fauna of Iran, with the description of 16 new species. European Journal of Taxonomy 843: 1–136. <https://doi.org/10.5852/ejt.2022.843.1947>
- Wood, TJ, Boustani M, Rasmont P (2020) A revision of the *Andrena* (Hymenoptera: Andrenidae) of Lebanon with the description of six new species. Annales de la Société entomologique de France (N.S.) 56: 279–312. <https://doi.org/10.1080/00379271.2020.1794960>