



Ecologica Montenegrina
ISSN 2396-9744 (online) | ISSN 2337-0173 (print)

Ecologica Montenegrina 74: 21-26 (2024)
This journal is available online at: www.biotaxa.org/em
<https://dx.doi.org/10.37828/em.2024.74.3>

Correspondence


First occurrence of the mite *Litarachna duboscqi* Walter, 1925 (Acariformes: Pontarachnidae) in the central Mediterranean Sea

SABRINA LO BRUTTO^{1,2}, ANTONINA BADALUCCO^{1,2} & HARRY SMIT^{3,4}

¹ Department of Earth and Marine Sciences, University of Palermo, via Archirafi 20, 90123, Palermo, Italy;
 <https://orcid.org/0000-0002-9964-904X>,  <https://orcid.org/0000-0002-5003-0692>

² National Biodiversity Future Center (NBFC), Piazza Marina 61, 90133, Palermo, Italy.

³ Naturalis Biodiversity Center, P.O. Box 9517, 2300 RA Leiden, the Netherlands;

 <https://orcid.org/0000-0002-0376-6808>

⁴ Museums Victoria Research Institute, Museums Victoria, GPO Box 666, Melbourne, VIC 3001, Australia.

Received 30 May 2024 | Accepted by V. Pešić: 17 June 2024 | Published online 4 July 2024.

One of the primary goals in assessing marine biodiversity is to measure the degree of species richness in specific geographical areas. We know that biodiversity is declining mainly due to human-induced pressures, but, at the same time, we don't know the extent of abundance and distribution of many species such as the ones ascribable to neglected groups (Schimmenti *et al.* 2016; Lo Brutto & Iacofano 2020). Under the framework of the National Biodiversity Future Center (NBFC), recently funded in Italy (Cena & Labra 2024), a survey on marine biodiversity was conducted aiming to detect rare or overlooked species in Sicily. The present paper reports the water mite identified as *Litarachna duboscqi* collected in March 2024 from nautical ropes in the harbour of Trapani (Sicily). It is the first record for southern Italy and the central Mediterranean Sea.

Sicily is the largest island (more than 25,000 km²) in the Mediterranean Sea with approximately 1,400 km of coastline, bounding the Tyrrhenian Sea to the north, the Ionian Sea to the east and the Strait of Sicily to the south. It represents a strategic natural observatory for monitoring Italian marine biodiversity (Lo Brutto *et al.* 2021; Servello *et al.* 2019; Tiralongo *et al.* 2022). It is a seascape inhabited by a huge variety of animal species which, overall, mirror the Mediterranean biodiversity, from jellyfish to mammals (Lo Brutto *et al.* 2021; Servello *et al.* 2019; Tiralongo *et al.* 2022).

Recent surveys have highlighted its role in detecting species which are considered neglected as they are not targeted in sampling designs or taxonomical studies (e.g. Lo Brutto *et al.* 2024). Marine mites belong to such a category; they have never been widely investigated along the Italian coast and their distribution, ecology and role in ecosystems are almost unknown (Pešić *et al.* 2019; Chatterjee & Durucan, 2022).

The only marine families of mites, Halacaridae Murray, 1877 and Pontarachnidae Koenike, 1910, are exclusively meiobenthic, with body size of less than 2 mm (Durucan 2021). The difference in species richness of the two families, more than 1100 species in Halacaridae (Bartsch 2006) and more than fifty in Pontarachnidae (Chatterjee *et al.* 2019), is maintained in the Italian faunal assemblage (Carriglio 2010). The family Pontarachnidae represents an important and, at the same time, a neglected component of the marine meiofauna (Pešić *et al.* 2019); it has an almost worldwide distribution (Smit,

2003) in tropical and subtropical seas. Species of this family have been reported from a depth of 0.5 m to almost 70 m (Pešić *et al.* 2012), occasionally in freshwater or brackish habitats (Smit 2002).

The family Pontarachnidae comprises only two genera, i.e. *Litarachna* Walter, 1925 and *Pontarachna* Philippi, 1840, with 23 and 30 species worldwide respectively (Chatterjee *et al.* 2019). The two genera can easily be separated by the suture line of the third and fourth coxae, which ends in the posteromedial apodemes in *Pontarachna*, but does not extend to these apodemes in *Litarachna*. *Litarachna duboscqi* is the only species in the Mediterranean Sea with a distinct ventral extension of the fourth segment of the palp.

The few Italian reports of this family concern the works carried out by authors which detected equally the two genera; some examples are *Litarachna communis* Walter, 1925, reported in Ischia (Mari & Morselli 1983); *Pontarachna punctulum* Philippi, 1840, described from samples collected in Trieste (Adriatic Sea; Schaub 1889), and more recently collected in *Posidonia oceanica* meadows along Tuscany coast (Tyrrhenian Sea; Bedini *et al.* 2011); *Pontarachna aenariensis* Mari & Morselli, 1983 was described from samples collected at Ischia island (Tyrrhenian Sea; Mari and Morselli 1983). The last checklist of mites in Italian seas dates to 2010 (Carriglio 2010). Eight species belonging to this family were listed, some of which are now considered invalid. : *Litarachna divergens* Walter, 1925 is considered as *species incerta*; *Litarachna incerta* Walter, 1925 is not valid; *Pontarachna tergestina* Von Schaub, 1889 is considered synonymous of *Pontarachna punctulum* Philippi, 1840 (Chatterjee *et al.* 2019). Thus a focus on their biodiversity assessment needs. In this paper, we report the first record of *Litarachna duboscqi* Walter, 1925 from southern Italy and improve the knowledge on its habitat.

The sampling site was located in the harbour of Trapani (38° 00' 53.7" N 12° 30' 04.4" E), where a rich fouling community mostly represented by sessile suspension feeders thrives attached to artificial substrates of the infrastructures. The samples were collected in March 2024 from a single nautical rope at depths between 0 and 1m. Sorting was carried out under a stereomicroscope to observe and collect small-sized organisms. Two mites were collected alive and then preserved in absolute ethanol. To be able to identify the species, one of the specimens was treated with KOH. This specimen is lodged in Naturalis Biodiversity Center, Leiden (registration number RMNH.ACA.7792).




Figure 1. Specimen of *Litarachna duboscqi* Walter, 1925. Left, dorsal view; right, ventral view.

The specimens were identified as *Litarachna duboscqi* (Figure 1). They were found associated with a rich suspension-feeder fauna. The fouling community was composed of sponges, polychaetes, crustacean tanaids and amphipods, ophiuroids, and tunicates, not identified to species level, and a particularly high richness of molluscs, identified as *Aplysia punctata*, *Columbella rustica*, *Diodora graeca*, *Fissurella nubecula*, *Isognomon* sp., *Jujubinus exasperatus*, *Mytilaster minimus*, *Ocenebra edwardsii*, *Pinctada radiata*, *Tricolia pullus*, *Tritia incrassata* (juvenile).

The species was originally described by Walter (1925) based on material from the French Mediterranean coast and from the Adriatic Sea (Rovigno), and later on reported by Fischetti (1928) from the eastern Mediterranean Sea under the name of *Pontarachna rhodia* (syn. of *Litarachna duboscqi*). Viets (1941) reported *Litarachna duboscqi* from Split, Croatia (Adriatic Sea). According to Chatterjee *et al.* 2019 (and references therein), this species is widely distributed in both the western and eastern Mediterranean Sea and Black Sea (see Figure 2 for details), on various substrates. A single record in the eastern Atlantic region demonstrates that the species can disperse outside the Mediterranean (Zawal & Pešić 2015) and highlights a probable Atlanto-Mediterranean range to investigate.




Figure 2. According to the literature, the geographical range of *Litarachna duboscqi* is herein shown on the map. Black circles are the documented records from the Mediterranean Sea (France, Italy, Croatia, Greece, Turkey), from the Black Sea (Russia, Crimea), and the Atlantic Ocean (Spain, Canary Islands). The red circle is the non-documented Mediterranean record from Italy (Ligurian Sea sector). The green triangle indicates the first southern Italy record from Sicily Island.

Chatterjee *et al.* (2019) gave no records from Italy, although the species is included in Italian checklists (Bernini *et al.* 1996). In the online checklist of the Italian fauna (<https://www.faunaitalia.it/checklist/introduction.html>, viewed May 4, 2024) *Litarachna duboscqi* is included with a reference to Walter (1925). The old record from Rovigno by Walter (1925) was at that time part of Italy but is nowadays part of Croatia. Also Carriglio (2010) included the species in his overview, without giving exact locations. The only record of the present-day Italian territory is from Stammer (1932) who reported the species from Duino in the northern Adriatic Sea.

The Mediterranean littoral environment has been investigated for a long time, especially along the coast of the European Union countries where *L. duboscqi* occurs. A huge literature from these countries regards taxonomic and ecological explorations on a great variety of substrates, and contextually many countries are required to monitor marine ecosystems under the Marine Strategy Framework Directive (Servello *et al.* 2019). Nevertheless, the species has not been detected widely.

Pontarachnidae is a taxon with a globose body shape that can be easily distinguished from the other marine Halacaridae group, however, despite that, it is sporadically detected. Very little is understood about its life cycle and trophic role, and the function of unique morphological features not observed in any other water mite species remains unknown (Smit & Alberti 2009; Kapiris *et al.* 2014; Pešić *et al.* 2019).

Pešić *et al.* (2019) highlighted that pontarachnid mites are overlooked as their bright colour and active behaviour make them one of the groups of marine meiofauna that should be separated directly in the field. The finding of the present samples gave a positive result as the sorting protocol of the sessile communities followed the above suggestion and was performed with the live specimens. It demonstrates that accurate sampling protocols are advisable, particularly in areas, such as Sicily, considered a hotspot for the detection of exotic and rare species or species in the phase of range extension. Further, in light of the local growing pan-oceanic maritime traffic (Lo Brutto *et al.* 2024), neglected species associated with the fouling communities in harbours should be deeply investigated.

Acknowledgments

The authors are indebted to Fabio Crocetta (Stazione Zoologica Anton Dohrn, Naples) for the identification of Mollusca and to Antonio Di Sabatino (L'Aquila) and Reinhard Gerecke (Tübingen) for providing us with a number of references.

Project partially funded under the National Recovery and Resilience Plan (PNRR), Mission 4 Component 2 Investment 1.4 - Call for tender No. 3138 of 16 December 2021, rectified by Decree n.3175 of 18 December 2021 of Italian Ministry of University and Research funded by the European Union – NextGenerationEU. Award Number: Project code CN_00000033, Concession Decree No. 1034 of 17 June 2022 adopted by the Italian Ministry of University and Research, CUP B73C22000790001, Project title “National Biodiversity Future Center - NBFC”.

References

- Bartsch, I. (2006) Halacaroidea (Acari): a guide to marine genera. *Organism Diversity & Evolution*, 6(2), 125.
- Bedini, R., Pertusati, M., Batistini, F. & Piazzini, L. (2011) Spatial and temporal variation of motile macro-invertebrate assemblages associated with *Posidonia oceanica* meadows. *Acta Adriatica*, 52(2), 201–214.
- Bernini, F., Castagnoli, M. & Nanelli, R. (1996) Arachnida Acari. In: Minelli, A., Ruffo, S. & La Posta, S. (eds), Checklist delle specie della Fauna Italiana. Ministero dell’Ambiente, Comitato Scientifico per la Fauna d’Italia.
- Carriglio, D. (2010) Arachnida Acari, Halacaridae – Pontarachnidae. *Biologia Marina Mediterranea*, 17 (suppl. 1), 390–393.
- Cena, H., & Labra, M. (2024) Biodiversity and planetary health: a call for integrated action. *The Lancet*, 403(10440), 1985–1986. [https://doi.org/10.1016/S0140-6736\(24\)00292-7](https://doi.org/10.1016/S0140-6736(24)00292-7)
- Chatterjee, T., Schizas, N.V. & Pešić, V. (2019) A checklist of Pontarachnidae (Acari: Hydrachnidia) and notes on distributional patterns of the species. *Zootaxa* 4619(3), 527–544. <https://doi.org/10.11646/zootaxa.4619.3.6>
- Chatterjee, T. & Durucan, F. (2022) A checklist of halacarid and pontarachnid mites (Acari: Halacaridae and Pontarachnidae) found in seagrass habitats. *Persian Journal of Acarology*, 11(2), 187–223.
- Durucan, F. (2021) New records of marine water mites (Acari: Hydrachnidia, Pontarachnidae) from Turkey. *Persian Journal of Acarology*, 10(2), 145–153.
- Fischetti, E. (1928) Descrizione di un nuovo idracnide marino dell’Egeo “*Pontarachna rhodia*” n. sp. *Archivio Zoologico Italiano*, 12, 365–368.
- Kapiris, K., Apostolidis, C., Baldaconi, R., Basusta, N., Bilecenoglu, M., Bitar, G., Bobori, D.C., Boyaci, Y.O., Dimitriadis, C., Djurovic, M., Dulcic, J., Durucan, F., Gerovasileiou, V., Gokoglu, M., Koutsoubas, D., Lefkaditou, E., Lipej, L., Markovic, O., Mavric, B., Ozvarol, Y., Pesic, V., Petriki, O., Siapatis, A., Sini, M., Tibullo, D. & Tiralongo, F. (2014) New Mediterranean Biodiversity Records (April, 2014). *Mediterranean Marine Science*, 15(1), 198–212. <https://doi.org/10.12681/mms.737>

- Lo Brutto, S., Badalucco, A. & Musco, L. (2024) First occurrence of the family Botryllophilidae Sars GO, 1921 (Copepoda, Cyclopoida) in Sicily, central Mediterranean Sea. *Crustaceana*, 97(3-4), 321–328. <https://doi.org/10.1163/15685403-bja10374>
- Lo Brutto, S., Calascibetta, A., Pavan, G. & Buffa, G. (2021) Cetacean strandings and museum collections: A focus on Sicily Island crossroads for Mediterranean species. *Diversity*, 13(3), 104. <https://doi.org/10.3390/d13030104>
- Lo Brutto, S. & Iacifano, D. (2020) New records of amphipod crustaceans along the Israeli Mediterranean coast, including a rare Mediterranean endemic species, *Maera schieckei* Karaman and Ruffo, 1971. *Biodiversity Data Journal*, 8, e53864. <https://doi.org/10.3897/BDJ.8.e53864>
- Mari, M. & Morselli, I. (1983) Osservazione su due specie di Hydrachnellae (Acari) marine: *Pontarachna aenariensis* n. sp. e *Litarachna communis* Walter. *Atti della Società Toscana di Scienze Naturali: Memorie, Serie B*, 90, 91–98.
- Pešić, V., Chatterjee, T. & Schizas, N.V. (2012) A new species of *Pontarachna* (Acari, Hydrachnidia, Pontarachnidae) from a mesophotic coral ecosystem off Vieques Island, Puerto Rico, Caribbean Sea. *Zootaxa*, 3440, 63–67. <https://doi.org/10.11646/ZOOTAXA.3440.1.3>
- Pešić, V., Durucan, F. & Zawal, A. (2019) Marine mites (Acari: Hydrachnidia) of the Mediterranean Sea: Descriptions of two new species, key for identification and future prospects. *Zootaxa*, 4585(3), 501–516. <https://doi.org/10.11646/zootaxa.4585.3.6>
- Schaub, R. von. (1889) *Über marine Hydrachniden nebst einigen Bemerkungen über Midea* (Bruz.). Kaiserliche Akademie der Wissenschaften in Wien. Mathematisch-Naturwissenschaftliche Klasse. Abt. 1, 98, 163–179.
- Schimmenti, E., Musco, L., Lo Brutto, S., Mikac, B., Nygren, A. & Badalamenti, F. (2016) Mediterranean record of *Eulalia ornata* (Annelida: Phyllodoceidae) corroborating its fidelity link with the *Sabellaria alveolata* reef habitat. *Mediterranean Marine Science*, 17(2), 359–370. <https://doi.org/10.12681/mms.1485>
- Servello, G., Andaloro, F., Azzurro, E., Castriota, L., Catra, M., Chiarore, A., Crocetta, F., D'Alessandro, M., Denitto, F., Frogliola, C., Gravili, C., Langer, M., Lo Brutto, S., Mastrototaro, F., Petrocelli, A., Pipitone, C., Piraino, S., Relini, G., Serio, D., Xentidis, N. & Zenetos, A. (2019) Marine alien species in Italy: A contribution to the implementation of descriptor D2 of the marine strategy framework directive. *Mediterranean Marine Science*, 20(1), 1–48. <https://doi.org/10.12681/mms.18711>
- Smit, H. (2002) Two new species of the water mite family Pontarachnidae (Acari: Hydrachnidia), with a discussion of the taxonomic status of *Pontarachna hinumaensis* Imamura. *Zootaxa*, 22(1), 1–8. <https://doi.org/10.11646/zootaxa.22.1.1>
- Smit, H. (2003) Five new species of the water mite family Pontarachnidae from Western Australia (Acari: Hydrachnidia). *The Marine Flora and Fauna of Dampier, Western Australia*. Western Australian Museum, Perth, pp. 547–562.
- Smit, H. & Alberti, G. (2009) The water mite family Pontarachnidae, with new data on its peculiar morphological structures (Acari: Hydrachnidia). In: Sabelis, M.W. & Bruin, J. (eds.), *Trends in Acarology: Proceedings of the 12th international congress*, Springer, Dordrecht, New York, pp. 71–79.
- Stammer, H.J. (1932) Die Fauna des Timavo. Ein Beitrag zur Kenntnis der Höhlengewässer, des Süß- und Brackwasser im Karst. *Zoologische Jahrbücher, Abteilung für Systematik, Ökologie und Geographie der Tiere*. 63(5-6), 521–656.
- Tiralongo, F., Badalamenti, R., Arizza, V., Prieto, L. & Lo Brutto, S. (2022) The Portuguese Man-of-War has always entered the Mediterranean sea—Strandings, sightings, and museum collections. *Frontiers in Marine Science*, 9, 856979. <https://doi.org/10.3389/fmars.2022.856979>
- Viets, K. (1941) Marine Halacaridae und Hydrachnellae (Acari) von der dalmatinischen Küste bei Split (Jugoslawien). *Godišnjak Oceanografskog Instituta, Split*, 2, 1–9.
- Walter, C. (1925) Marine Hygrobatidae. Revision der Wassermilben-Genera *Pontarachna* Philippi und *Nautarachna* Moniez. *Internationale Revue der Gesamten Hydrobiologie und Hydrographie*, 14(1-2), 1–54.

Zawal, A. & Pešić, V. (2015) The first record of *Litarachna dubosqi* Walter, 1925 (Acari, Pontarachnidae) outside the Mediterranean Sea. *Oceanological and Hydrobiological Studies*, 44(3), 426–429. <https://doi.org/10.1515/ohs-2015-0039>