



Naturalis Repository

## Tropical forests in the Americas are changing too slowly to track climate change

Jesús Aguirre-Gutiérrez, Sandra Díaz, Sami W. Rifai, José Javier Corral-Rivas, María Guadalupe Nava-Miranda +129 more

DOI:

<https://doi.org/10.1126/science.adl5414>

Downloaded from

[Naturalis Repository](#)

### Article 25fa Dutch Copyright Act (DCA) - End User Rights

This publication is distributed under the terms of Article 25fa of the Dutch Copyright Act (Auteurswet) with consent from the author. Dutch law entitles the maker of a short scientific work funded either wholly or partially by Dutch public funds to make that work publicly available following a reasonable period after the work was first published, provided that reference is made to the source of the first publication of the work.

This publication is distributed under the Naturalis Biodiversity Center 'Taverne implementation' programme. In this programme, research output of Naturalis researchers and collection managers that complies with the legal requirements of Article 25fa of the Dutch Copyright Act is distributed online and free of barriers in the Naturalis institutional repository. Research output is distributed six months after its first online publication in the original published version and with proper attribution to the source of the original publication.

You are permitted to download and use the publication for personal purposes. All rights remain with the author(s) and copyrights owner(s) of this work. Any use of the publication other than authorized under this license or copyright law is prohibited.

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the department of Collection Information know, stating your reasons. In case of a legitimate complaint, Collection Information will make the material inaccessible. Please contact us through email: [collectie.informatie@naturalis.nl](mailto:collectie.informatie@naturalis.nl). We will contact you as soon as possible.

## RESEARCH ARTICLE

## FOREST CHANGE

## Tropical forests in the Americas are changing too slowly to track climate change

Jesús Aguirre-Gutiérrez<sup>1,120\*</sup>, Sandra Díaz<sup>2,3</sup>, Sami W. Rifai<sup>4</sup>, Jose Javier Corral-Rivas<sup>5</sup>, Maria Guadalupe Nava-Miranda<sup>6,7</sup>, Roy González-M<sup>8,119</sup>, Ana Belén Hurtado-M<sup>8</sup>, Norma Salinas Revilla<sup>9</sup>, Emilio Vilanova<sup>10</sup>, Everton Almeida<sup>11</sup>, Edmar Almeida de Oliveira<sup>12</sup>, Esteban Alvarez-Davila<sup>13</sup>, Luciana F. Alves<sup>14</sup>, Ana Cristina Segalin de Andrade<sup>15</sup>, Antonio Carlos Lola da Costa<sup>16</sup>, Simone Aparecida Vieira<sup>17</sup>, Luiz Aragão<sup>18,19</sup>, Eric Arets<sup>20</sup>, Gerardo A. Aymard C.<sup>21</sup>, Fabrício Baccaro<sup>22</sup>, Yvonne Vanessa Bakker<sup>23</sup>, Timothy R. Baker<sup>24</sup>, Olaf Bánki<sup>25</sup>, Christopher Baraloto<sup>26</sup>, Plínio Barbosa de Camargo<sup>27</sup>, Erika Berenguer<sup>1,28</sup>, Lilian Blanc<sup>29,30</sup>, Damien Bonal<sup>31</sup>, Frans Bongers<sup>32</sup>, Kauane Maiara Bordin<sup>121</sup>, Roel Brienen<sup>24</sup>, Foster Brown<sup>34</sup>, Nayane Cristina C. S. Prestes<sup>12</sup>, Carolina V. Castilho<sup>35</sup>, Sabina Cerruto Ribeiro<sup>36</sup>, Fernanda Coelho de Souza<sup>37</sup>, James A. Comiskey<sup>38,39</sup>, Fernando Cornejo Valverde<sup>40</sup>, Sandra Cristina Müller<sup>33</sup>, Richarly da Costa Silva<sup>41</sup>, Julio Daniel do Vale<sup>42</sup>, Vitor de Andrade Kamimura<sup>23,43</sup>, Ricardo de Oliveira Perdiz<sup>44,45</sup>, Jhon del Aguila Pasquel<sup>46,47</sup>, Géraldine Derroire<sup>48</sup>, Anthony Di Fiore<sup>49,50</sup>, Mathias Disney<sup>51,52</sup>, William Farfan-Rios<sup>53,54</sup>, Sophie Fauset<sup>55</sup>, Ted R. Feldpausch<sup>56</sup>, Rafael Flora Ramos<sup>23</sup>, Gerardo Flores Llampazo<sup>46</sup>, Valéria Forni Martins<sup>57,58</sup>, Claire Fortunel<sup>59</sup>, Karina Garcia Cabrera<sup>60</sup>, Jorcely Gonçalves Barroso<sup>61</sup>, Bruno Hérault<sup>29,62</sup>, Rafael Herrera<sup>63</sup>, Eurídice N. Honorio Coronado<sup>64</sup>, Isau Huamantupa-Chuquimaco<sup>65,66</sup>, John J. Pipoly<sup>67,68</sup>, Katia Janaina Zanini<sup>33</sup>, Eliana Jiménez<sup>69</sup>, Carlos A. Joly<sup>57</sup>, Michelle Kalamandeen<sup>70</sup>, Joice Klipel<sup>122</sup>, Aurora Levesley<sup>24</sup>, Wilmar Lopez Oviedo<sup>71,72</sup>, William E. Magnusson<sup>73</sup>, Rubens Manoel dos Santos<sup>74</sup>, Beatriz Schwantes Marimon<sup>12</sup>, Ben Hur Marimon-Junior<sup>12</sup>, Simone Matias de Almeida Reis<sup>12,36</sup>, Omar Aurelio Melo Cruz<sup>75</sup>, Abel Monteagudo Mendoza<sup>54,106</sup>, Paulo Morandi<sup>12</sup>, Robert Muscarella<sup>76</sup>, Henrique Nascimento<sup>77</sup>, David A. Neill<sup>78</sup>, Imma Oliveras Menor<sup>1,59</sup>, Walter A. Palacios<sup>79</sup>, Sonia Palacios-Ramos<sup>80</sup>, Nadir Carolina Pallqui Camacho<sup>24,81</sup>, Guido Pardo<sup>82</sup>, R. Toby Pennington<sup>83,84</sup>, Luciana de Oliveira Pereira<sup>56</sup>, Georgia Pickavance<sup>24</sup>, Rayana Caroline Picoletto<sup>33</sup>, Nigel C. A. Pitman<sup>85</sup>, Adriana Prieto<sup>86</sup>, Carlos Quesada<sup>87</sup>, Hirma Ramírez-Angulo<sup>88</sup>, Maxime Réjou-Méchain<sup>89</sup>, Zorayda Restrepo Correa<sup>90</sup>, José Manuel Reyna Huaymacari<sup>46</sup>, Carlos Reynel Rodriguez<sup>91</sup>, Gonzalo Rivas-Torres<sup>50,118</sup>, Anand Roopsind<sup>92</sup>, Agustín Rudas<sup>86</sup>, Beatriz Salgado Negret<sup>93</sup>, Masha T. van der Sande<sup>32</sup>, Flávia Delgado Santana<sup>94</sup>, Flavio Antonio Maës Santos<sup>57</sup>, Rodrigo Scarton Bergamin<sup>95</sup>, Miles R. Silman<sup>60</sup>, Camila Silva<sup>96</sup>, Javier Silva Espejo<sup>97</sup>, Marcos Silveira<sup>36</sup>, Fernanda Cristina Souza<sup>98,123</sup>, Martin J. P. Sullivan<sup>99</sup>, Varun Swamy<sup>100</sup>, Joey Talbot<sup>101</sup>, John J. Terborgh<sup>102</sup>, Peter J. van der Meer<sup>103</sup>, Geertje van der Heijden<sup>104</sup>, Bert van Uff<sup>105</sup>, Rodolfo Vasquez Martinez<sup>106</sup>, Laura Vedovato<sup>19</sup>, Jason Vleminckx<sup>107</sup>, Vincent Antoine Vos<sup>82</sup>, Verginia Wortel<sup>108</sup>, Pieter A. Zuidema<sup>32</sup>, Joeri A. Zwarts<sup>109</sup>, Susan G. W. Laurance<sup>110</sup>, William F. Laurance<sup>110</sup>, Jérôme Chave<sup>111</sup>, James W. Dalling<sup>112,113</sup>, Jos Barlow<sup>28</sup>, Lourens Poorter<sup>32</sup>, Brian J. Enquist<sup>114,115</sup>, Hans ter Steege<sup>116,117</sup>, Oliver L. Phillips<sup>24</sup>, David Galbraith<sup>24</sup>, Yadvinder Malhi<sup>1,120</sup>

Understanding the capacity of forests to adapt to climate change is of pivotal importance for conservation science, yet this is still widely unknown. This knowledge gap is particularly acute in high-biodiversity tropical forests. Here, we examined how tropical forests of the Americas have shifted community trait composition in recent decades as a response to changes in climate. Based on historical trait-climate relationships, we found that, overall, the studied functional traits show shifts of less than 8% of what would be expected given the observed changes in climate. However, the recruit assemblage shows shifts of 21% relative to climate change expectation. The most diverse forests on Earth are changing in functional trait composition but at a rate that is fundamentally insufficient to track climate change.

**F**orest responses to human-driven perturbations such as climate change will largely determine the diversity and function of the terrestrial biosphere through this century and beyond. The Americas host the greatest concentration of tree species in the world (1), including six key biodiversity hotspots (2) and half of Earth's most intact trop-

ical forests (3). In the face of threats from climate change and continuing loss in area and integrity (3–6), it is both critical and urgent to understand the ability of these complex systems to adapt to change and survive.

Within tropical American forests (by which we mean all forests encompassing continental areas from Mexico to Brazil), lowland forests

provide relatively homogeneous climatic conditions over large areas, potentially allowing the existence of common functional adaptations over large spatial extents. By contrast, across mountain forests, climatic conditions tend to change rapidly in space, potentially facilitating the rapid turnover of functional adaptations to local environmental conditions. In Amazonia, changes in precipitation patterns and more frequent droughts have led to an increase in the recruitment of dry-affiliated species (xerophilization) (7). In the Andes, rising temperatures have led to increasing abundances of species tolerant to higher temperature (thermophilization) (8). Across Mesoamerica, it is expected that climate change will cause an expansion of tropical dry forests to higher elevations (>200 m above current average elevation) (9). However, tree species may be unable to shift their distribution fast enough to track their climatic niche given their slow demography (e.g., growth and recruitment), the prevalence of dispersal limitation (10), and different environmental tolerances at different life stages (11). All of these limitations would increase the vulnerability of tree species to climate change across tropical American forests. For instance, in higher latitudes, recent work has shown large range contractions of tree species rather than range expansions or shifts (12). Changes in climate across the tropical Americas are expected to become stronger, with some scenarios projecting temperature increases of up to ~4°C and precipitation reductions close to 20% by 2100 (13–15). This would likely increase the vulnerability of current tree species assemblages because they would face climates that they have not previously experienced (16), potentially selecting for no-analog future plant communities (17).

Functional traits mediate species responses to environmental change, affecting plant performance and species distributions (18–20). These morphological, structural, chemical, and phenological characteristics tend to show consistent relationships with climate and soil conditions (21). Recent work has shown positive relationships between mean annual temperature and leaf area, specific leaf area (SLA), leaf N content, wood density (WD), and leaf thickness (22), depicting plant functional adaptations to local environmental conditions. Other work has detected a negative relationship with elevation for SLA and leaf N content, potentially as an adaptation to cooler environments with lower nutrient availability (22). Therefore, these traits are tightly linked to the capacity of plant species to respond to environmental changes. For instance, having large area (i.e., larger leaves) can increase leaf temperature due to higher solar absorption, whereas smaller leaves dissipate heat more effectively and help to prevent water loss. Plants with lower SLA, i.e., with thicker and tougher leaves,

tend to be more resistant to drought because they can better resist water loss. High WD is tightly related to increased resistance to cavitation, which can increase the plant's capacity to survive droughts. Therefore, a trait-based approach provides a promising framework for predicting the impacts of climate change and resilience across forest ecosystems (19, 23, 24).

It is still unclear how shifts in the abundance and distribution of species translate into changes in the functional trait composition and what functional changes have occurred through the past half century as a response to the onset of a warmer, drier, and more variable climate across the tropical Americas. Moreover, it is unknown whether forest-level functional shifts are more attrib-

utable to differential growth among the surviving trees than to the addition (i.e., recruitment) or removal (i.e., mortality) of trees to the assemblage. It is also uncertain if these functional shifts match the direction of climate change and, if so, whether the rate of functional trait change keeps pace with climate change or lags behind it. Understanding the above will allow the quantification of the present, and likely future, capacity of forest to adapt to a changing climate and also to uncover which functional trait characteristics may confer forests with a higher adaptation capacity to a changing climate.


Here, we address these knowledge gaps by analyzing records from 415 long-term forest plot sites monitored over >40 years (1980 to 2021) by

the RAINFOR and MONAFOR networks and contributors to the ForestPlots.net meta-network. This dataset includes information on the identity, size, recruitment, and mortality of >250,000 individual trees across the tropics from Mexico to southern Brazil. Our effort spans relatively undisturbed forests from the lowland tropics (hereafter forest plots <700 m elevation) to pre-montane and montane zones (>700 m elevation; hereafter referred to as montane) from the Andes to subtropical fringes (Fig. 1 and data S1). These forests are distributed along a wide range of climatic and soil conditions (Fig. 1B) and have experienced strong changes in climate over the past decades (Fig. 1C). We combined this monitoring and analysis of changes in the plant community composition

<sup>1</sup>Environmental Change Institute, School of Geography and the Environment, University of Oxford, Oxford, UK. <sup>2</sup>Consejo Nacional de Investigaciones Científicas y Técnicas, Instituto Multidisciplinario de Biología Vegetal (IMBIV), Córdoba, Argentina. <sup>3</sup>Facultad de Ciencias Exactas, Físicas y Naturales, Universidad Nacional de Córdoba, Córdoba, Argentina. <sup>4</sup>School of Biological Sciences, University of Adelaide, Adelaide, Australia. <sup>5</sup>Facultad de Ciencias Forestales y Ambientales, Universidad Juárez del Estado de Durango, Durango, Mexico. <sup>6</sup>Escuela Politécnica Superior de Ingeniería, Campus Terra. Universidad de Santiago de Compostela, Lugo, España. <sup>7</sup>Colegio de Ciencias y Humanidades. Universidad Juárez del Estado de Durango, Durango, Mexico. <sup>8</sup>Programa Ciencias Básicas de la Biodiversidad, Instituto de Investigación de Recursos Biológicos Alexander von Humboldt, Bogotá, Colombia. <sup>9</sup>Institute for Nature Earth and Energy, Pontificia Universidad Católica del Perú, Lima, Peru. <sup>10</sup>Wildlife Conservation Society, New York, NY, USA. <sup>11</sup>Instituto de Biodiversidade e Florestas da Universidade Federal do Oeste do Pará (UFOPA), Rua Vera Paz, s/n (Unidade Tapajós), Bairro Salé, Santarém, Pará, Brasil. <sup>12</sup>Programa de Pós Graduação em Ecologia e Conservação, Universidade do Estado de Mato Grosso, Nova Xavantina, Brazil. <sup>13</sup>Universidad UNAD-Colombia, Escuela ECAPMA, Bogotá, Colombia. <sup>14</sup>Center for Tropical Research, Institute of the Environment and Sustainability, University of California, Los Angeles, CA, USA. <sup>15</sup>Projeto Dinâmica Biológica de Fragmentos Florestais, Instituto Nacional de Pesquisas da Amazônia - INPA, Av. André Araújo, 2936, Petrópolis, Manaus, Amazonas, Brazil. <sup>16</sup>Universidade Federal do Pará/Instituto de Geociências/Faculdade de Meteorologia/Belém, Pará, Brasil. <sup>17</sup>Center for Environmental Studies and Research, University of Campinas (UNICAMP), Campinas, Brazil. <sup>18</sup>Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, São Paulo, Brazil. <sup>19</sup>University of Exeter, Exeter, UK. <sup>20</sup>Wageningen Research, Wageningen University & Research, Wageningen, Netherlands. <sup>21</sup>UNELLEZ-Guanare, Programa de Ciencias del Agro y el Mar, Herbario Universitario (PORT), Estado Portuguesa, Venezuela. <sup>22</sup>Departamento de Biología, Universidade Federal do Amazonas, Manaus, Amazonas, Brazil. <sup>23</sup>Institute of Biology, University of Campinas (UNICAMP), Campinas, São Paulo, Brazil. <sup>24</sup>School of Geography, University of Leeds, Leeds, UK. <sup>25</sup>Naturalis Biodiversity Center, Leiden, Netherlands. <sup>26</sup>International Center for Tropical Botany (ICTB) Department of Biological Sciences, Florida International University, Miami, FL, USA. <sup>27</sup>Centro de Energía Nuclear na Agricultura, Universidade de São Paulo, São Paulo, Brazil. <sup>28</sup>Lancaster Environment Centre, Lancaster University, Lancaster, UK. <sup>29</sup>CIRAD, UPR Forêts et Sociétés, Montpellier, France. <sup>30</sup>Forêts et Sociétés, Univ Montpellier, CIRAD, Montpellier, France. <sup>31</sup>Université de Lorraine, AgroParisTech, INRAE, UMR Silva, Nancy, France. <sup>32</sup>Forest Ecology and Forest Management Group, Wageningen University & Research, Wageningen, Netherlands. <sup>33</sup>Plant Ecology Lab, Federal University of Rio Grande do Sul, Porto Alegre, Brazil. <sup>34</sup>Woodwell Climate Research Center, Falmouth, MA, USA. <sup>35</sup>Centro de Pesquisa Agroflorestal de Roraima, Embrapa Roraima, Boa Vista, Brazil. <sup>36</sup>Centro de Ciências Biológicas e da Natureza, Universidade Federal do Acre, Campus Universitário, Rio Branco, Brazil. <sup>37</sup>BeZero Carbon, London, UK. <sup>38</sup>National Park Service, Fredericksburg, VA, USA. <sup>39</sup>Smithsonian Institution, Washington, DC, USA. <sup>40</sup>Proyecto Castaña, Madre de Dios, Peru. <sup>41</sup>Instituto Federal de Educação, Ciência e Tecnologia do Acre, Campus Baixada do Sol, Rio Branco, Brazil. <sup>42</sup>Pontificia Universidade Católica do Paraná, Campus Toledo, Toledo, Brazil. <sup>43</sup>Biodiversity and Ecosystem Services, Instituto Tecnológico Vale, Belém, Pará, Brazil. <sup>44</sup>Programa de Pós-Graduação em Botânica, Instituto Nacional de Pesquisas da Amazônia (INPA), Manaus, Amazonas, Brazil. <sup>45</sup>Luz da Floresta, Boa Vista, Roraima, Brazil. <sup>46</sup>Instituto de Investigaciones de la Amazonia Peruana, Iquitos, Peru. <sup>47</sup>Universidad Nacional de la Amazonia Peruana, Iquitos, Peru. <sup>48</sup>Cirad, UMR EcoFoG (AgroParisTech, CNRS, INRAE, Université des Antilles, Université de la Guyane), Campus Agronomique, Kourou, French Guiana. <sup>49</sup>Department of Anthropology, The University of Texas at Austin, Austin, TX, USA. <sup>50</sup>Estación de Biodiversidad Tiputini, Colegio de Ciencias Biológicas y Ambientales, Universidad San Francisco de Quito (USFQ), Quito, Ecuador. <sup>51</sup>Department of Geography, University College London, London, UK. <sup>52</sup>NERC National Centre for Earth Observation (NCEO), London, UK. <sup>53</sup>Biology Department and Sabin Center for Environment and Sustainability, Wake Forest University, Winston-Salem, NC, USA. <sup>54</sup>Herbario Vargas (CUZ), Escuela Profesional de Biología, Universidad Nacional de San Antonio Abad del Cusco, Cusco, Peru. <sup>55</sup>School of Geography, Earth and Environmental Sciences, University of Plymouth, Plymouth, UK. <sup>56</sup>Geography, Faculty of Environment, Science, and Economy, University of Exeter, Exeter, UK. <sup>57</sup>Department of Plant Biology, Institute of Biology, University of Campinas (UNICAMP), Campinas, São Paulo, Brazil. <sup>58</sup>Department of Natural Sciences, Maths, and Education, Centre for Agrarian Sciences, Federal University of São Carlos (UFSCar), Araras, São Paulo, Brazil. <sup>59</sup>AMAP (Botanique et Modélisation de l'Architecture des Plantes et des Végétations), Université de Montpellier, CIRAD, CNRS, INRAE, IRD, Montpellier, France. <sup>60</sup>Biology Department and Sabin Center for Environment and Sustainability, Wake Forest University, Winston-Salem, NC, USA. <sup>61</sup>Federal University of Acre, Cruzeiro do Sul, Brazil. <sup>62</sup>Forêts et Sociétés, Univ Montpellier, CIRAD, Montpellier, France. <sup>63</sup>Instituto Venezolano de Investigaciones Científicas (IVIC), Caracas, Venezuela. <sup>64</sup>Royal Botanic Gardens, Kew, London, Richmond, UK. <sup>65</sup>Herbario Alwyn Gentry (HAG), Universidad Nacional Amazónica de Madre de Dios (UNAMAD), Puerto Maldonado, Madre de Dios, Peru. <sup>66</sup>Centro Ecológico INKAMAZONIA, Valle de Koshiñata, Cusco, Peru. <sup>67</sup>Broward County Parks & Recreation Division, Oakland Park, FL, USA. <sup>68</sup>Department of Biological Sciences, Florida Atlantic University, Boca Raton, FL, USA. <sup>69</sup>Grupo de Investigación en Ecología y Conservación de Fauna y Flora Silvestre, Instituto Amazónico de Investigaciones Imani, Universidad Nacional de Colombia – Sede Amazonia, Amazonas, Colombia, Suramérica. <sup>70</sup>Unique Land Use GmbH, Freiburg im Breisgau, Germany. <sup>71</sup>Smurfit Kappa Colombia, Yumbo, Valle del Cauca, Colombia. <sup>72</sup>Universidad Nacional de Colombia Medellín, Medellín, Antioquia, Colombia. <sup>73</sup>Instituto Nacional de Pesquisas da Amazônia, Manaus, Amazonas, Brazil. <sup>74</sup>Laboratory of Phytogeography and Evolutionary Ecology, Department of Forest Sciences, Federal University of Lavras, Lavras, Minas Gerais, Brazil. <sup>75</sup>Universidad del Tolima, Ibagué, Colombia. <sup>76</sup>Plant Ecology and Evolution, Evolutionary Biology Center, Uppsala University, Uppsala, Sweden. <sup>77</sup>Coordenação de Biodiversidade, Instituto Nacional de Pesquisas da Amazônia, Manaus, Amazonas, Brazil. <sup>78</sup>Universidad Estatal Amazónica, Puyo, Pastaza, Ecuador. <sup>79</sup>Herbario Nacional del Ecuador, Universidad Técnica del Norte, Quito, Ecuador. <sup>80</sup>Universidad Nacional Agraria La Molina, La Molina, Lima, Peru. <sup>81</sup>Universidad Nacional de San Antonio Abad del Cusco, Cusco, Peru. <sup>82</sup>Instituto de Investigaciones Forestales de la Amazonia, Universidad Autónoma del Beni José Ballivián, Riberaltta, Beni, Bolivia. <sup>83</sup>College of Life and Environmental Sciences, University of Exeter, Exeter, Devon, UK. <sup>84</sup>Tropical Diversity Section, Royal Botanic Gardens Edinburgh, Edinburgh, UK. <sup>85</sup>Science & Education, Field Museum of Natural History, Chicago, IL, USA. <sup>86</sup>Instituto de Ciencias Naturales, Universidad Nacional de Colombia, Ciudad Universitaria, Bogotá, Colombia. <sup>87</sup>Coordination of Environmental Dynamics, National Institute for Amazonian Research, Manaus, Amazonas, Brazil. <sup>88</sup>Instituto de Investigaciones para el Desarrollo Forestal (INDEFOR), Universidad de los Andes, Mérida, Venezuela. <sup>89</sup>AMAP, Univ. Montpellier, IRD, CNRS, CIRAD, INRAE, Montpellier, France. <sup>90</sup>Corporación COL-TREE, Medellín, Colombia. <sup>91</sup>Facultad de Ciencias Forestales, Universidad Nacional Agraria La Molina, Lima, Peru. <sup>92</sup>Conservation International, Arlington, VA, USA. <sup>93</sup>Departamento de Biología, Universidad Nacional de Colombia, Bogotá, Colombia. <sup>94</sup>Instituto Nacional de Pesquisas da Amazônia (INPA), Manaus, Amazonas, Brazil. <sup>95</sup>Birmingham Institute of Forest Research (BIFOR), University of Birmingham, Birmingham, UK. <sup>96</sup>Instituto de Pesquisa Ambiental da Amazônia (IPAM), Brasília-DF, Brazil. <sup>97</sup>Departamento de Biología, Universidad de La Serena, La Serena, Chile. <sup>98</sup>Departamento de Ecologia e Conservação, Instituto de Ciências Naturais, Universidade Federal de Lavras, Lavras, Minas Gerais, Brazil. <sup>99</sup>Department of Natural Sciences, Manchester Metropolitan University, Manchester, UK. <sup>100</sup>Center for Energy, Environment & Sustainability, Wake Forest University, Wake Forest, NC, USA. <sup>101</sup>Institute for Transport Studies, University of Leeds, Leeds, UK. <sup>102</sup>Department of Biology, University of Florida, Gainesville, FL, USA. <sup>103</sup>Van Hall Larenstein University of Applied Sciences, Velp, Netherlands. <sup>104</sup>School of Geography, University of Nottingham, University Park, Nottingham, UK. <sup>105</sup>Royal Netherlands Meteorological Institute (KNMI), De Bilt, Netherlands. <sup>106</sup>Jardín Botánico de Missouri, Oxapampa, Peru. <sup>107</sup>Université Libre de Bruxelles, Brussels, Belgium. <sup>108</sup>Department of Forest Management, Centre for Agricultural Research in Suriname, CELOS, Paramaribo, Suriname. <sup>109</sup>Utrecht University, Utrecht, Netherlands. <sup>110</sup>Centre for Tropical Environmental and Sustainability Science, College of Science and Engineering, James Cook University, Cairns, Queensland, Australia. <sup>111</sup>Centre Recherche Biodiversité Environnement, CNRS, Université Paul Sabatier, IRD, INPT, UMR5300, Toulouse, France. <sup>112</sup>Department of Plant Biology, University of Illinois, Urbana-Champaign, IL, USA. <sup>113</sup>Smithsonian Tropical Research Institute, Ancon, Republic of Panama. <sup>114</sup>Department of Ecology and Evolutionary Biology, University of Arizona, Tucson, AZ, USA. <sup>115</sup>Santa Fe Institute, Santa Fe, NM, USA. <sup>116</sup>Tropical Botany, Naturalis Biodiversity Center, Leiden, Netherlands. <sup>117</sup>Quantitative Biodiversity Dynamics, Department of Biology, Utrecht University, Utrecht, Netherlands. <sup>118</sup>Department of Wildlife Ecology and Conservation, University of Florida, Gainesville, FL, USA. <sup>119</sup>Departamento de Ciencias Forestales, Facultad de Ingeniería Forestal, Universidad del Tolima, Tolima, Colombia. <sup>120</sup>Leverhulme Centre for Nature Recovery, University of Oxford, Oxford, UK. <sup>121</sup>Department of Ecology, Universidade Federal do Rio Grande do Sul, Porto Alegre, Brazil. <sup>122</sup>Institute of Ecology, Leuphana University of Lüneburg, Germany. <sup>123</sup>Instituto de Ciências Biológicas, Programa de Pós-Graduação em Ecologia, Universidade Federal do Pará, Pará, Belém, Brazil.

\*Corresponding author. Email: jeaggu@gmail.com





**Fig. 1. Study area showing the distribution and number of vegetation plots sampled across time, principal component analysis, location of sampling plots, and change in climate conditions (A).** (B) Principal component analysis (PC1, PC2, and PC3) depicting the climate and soil chemistry and texture space available in the study area. (B and C) Location of the sampling plots in the environmental space (B) and change in climate conditions (1980 to 1990 versus 2010 to 2020) in the plot network (C). In (B), PC1 is mainly loaded by the MCWD (−0.527) and VPD (−0.515),

PC2 by  $T_{\text{mean}}$  (−0.465) and soil CEC (0.524), and PC3 by soil clay percentage (−0.535) and soil sand percentage (0.486). In (C), the vertical dotted lines indicate zero change. Brown colors depict increases in temperature and drier conditions for MCWD and VPD or increased drought intensity for the SPEI. Blue colors depict an increase in water availability. In MCWD, larger positive values indicate higher water stress. Climate data were derived from the TerraClimate project (48) and soil data from SoilGrids (<https://soilgrids.org/>) (49).

with measurements of 12 plant functional traits that are potentially involved in responses to a changing climate. These include photosynthetic capacity ( $A_{\text{sat}}$ ), leaf chemistry (content of leaf C, N, and P), leaf area, SLA, leaf fresh mass (FM), leaf thickness, abundance of deciduous species (DE), adult maximum height ( $H_{\text{max}}$ ), WD, and seed mass (SM) (table S1). Tree functional trait data were obtained for several plots from local field collections carried out by collaborators (25–27), the Global Ecosystems Monitoring network (GEM) (28), and ForestPlots ([www.ForestPlots.net](http://www.ForestPlots.net)) (29), in addition to databases from the Botanical Information and Ecology Network (BIEN; [bien.nceas.ucsb.edu](http://bien.nceas.ucsb.edu)), the TRY Plant Trait Database ([www.try-db.org](http://www.try-db.org)) (30), and Díaz *et al.* (19, 31).

We first investigated long-term plant trait–environment relationships to understand how climate drives trait distributions in tropical forests of the Americas and if these relationships are consistent across lowland and montane forests. We expected temperature and water availability to be the main drivers of plant trait distributions, with warmer and drier areas facilitating the dominance of more conservative trait syndromes (e.g., smaller and thicker leaves, higher WD, lower  $A_{\text{sat}}$ ) compared with warm and wetter areas (32, 33). Moreover, we expected trait–environment relationships to differ between lowland and montane forests given the different climatic ranges of these forest types.

We next studied how and where lowland and montane tropical American forests have shifted in their functional trait composition


due to changes in the plant community taxonomic composition over the past four decades. We did this by analyzing the annual rate of change ( $\Delta r$ ) of the trait community-weighted mean (CWM) for all forests (lowland and montane together) and for lowland and montane forest separately. Because of the long lifespan of tropical trees (34) and their slow turnover, we performed this analysis at the full community level and separately for the recruiting (“recruit”), mortality (“fatality”), and surviving (hereafter “survivor”) assemblages (Fig. 2). Analyzing changes at the full community level [involving all trees >10 cm diameter at breast height (DBH) alive] allows us to understand how communities are changing in their trait CWM given tree growth, survival, and recruitment together. Analyzing the survivor (change in CWM given by growth) assemblage alone will provide insights into potentially more resistant trait values, whereas analyses for the fatality assemblages will identify potentially less resistant trait values. The recruit community will affect the full community-level trait composition dependent on their basal area. It will also provide information on potentially better-adapted trait values to the current climate that allow them to recruit into the community, as well as indicate the possible composition of future forests.

We further investigated whether observed changes in trait composition have been enough to track climate change to date by comparing observed and expected trait changes based on historical trait–environment relationships [see the materials and methods (35)]. This climate

change tracking analysis was performed for the full community assemblage and for the survivor and recruit assemblages, but not for the fatality assemblage because these individuals will not contribute to future change (Fig. 2).

Given exposure to a drying and warming climate, we could reasonably expect an increased abundance of species exhibiting more drought-tolerance traits (i.e., in the “slow” section of the plant economics spectrum) (36), such as high WD (e.g., to prevent cavitation) (37) and smaller, thicker leaves (e.g., for lower evapotranspiration and reduced radiation exposure) (38). However, it is also possible that increasing drought will drive a shift toward drought-avoidance traits, notably DE (which is often associated with more acquisitive leaves) (32, 39). Seed traits play a pivotal role in the reproduction and dispersal capacity of species (10). Under an unstable, warming, and drying climate, we might expect species with smaller wind-dispersed seeds to increase in abundance (40). This is because wind-dispersed seeds, which are more common in drier and more seasonal biomes, tend to be smaller than animal-dispersed seeds (41). However, other factors, such as wind and fire disturbance and defaunation of frugivorous seed-dispersing mammals and birds, may disrupt the expected trends in seed traits because these drive shifts more strongly at shorter time scales than a changing climate (42). If migration is an important component of species response to climate change, then we would also expect montane forests to show stronger functional responses than lowland forests given their more

## Mechanisms for change in community trait composition



**Fig. 2. Conceptual figure depicting the analyzed mechanisms for change in community trait composition across the study area.** Tree individuals that are alive and have a DBH  $\geq 10$  cm are part of the full community assemblage. Across time, there can be changes in the community trait composition due to growth of the surviving tree individuals (survivor assemblage) given their increase in basal area (top right). Other mechanisms for changing community trait composition across time are the recruitment (recruit assemblage) of new individuals (middle right) and the death (fatality assemblage) of individuals in the community (bottom right).

varied climatic conditions at shorter distances (8, 33) that make it potentially easier to migrate to a favorable climate than in the lowlands (43–46). In montane forests, nutrient availability (e.g., N:P ratios) can vary significantly along altitudinal gradients due to substantial changes in temperature and water availability (47). As a result, we expect strong functional responses to soil nutrient availability across these elevation gradients.

We expect that, given the long lifespan of tropical trees and the rapid pace of recent climate change, forests will show ecological inertia so that changes in functional composition will lag behind changes in climate. We expect the full community and survivor assemblages to show slower change given that their change is largely dependent on tree growth, which is a slow process among tropical forest trees. The recruit and fatality assemblages may show faster and larger community trait responses because they are less dependent on growth and more dependent on local climate conditions.

### Long-term trait-environment relationships


To evaluate long-term (1980 to 2021) trait-climate relationships across tropical American forests,

we used data from 415 forest plots [mean plot size 0.88 ha (minimum: 0.12, maximum: 25) and 5.7 censuses per plot (minimum: 2, maximum: 41)], for which we extracted climate (48) and soil (49) data for their sampling years. Because species' contributions to ecosystem processes likely depend on their relative abundances (50), we calculated the CWM of each plant functional trait (table S1) for each plot based on the relative basal area of the species and their trait value (hereafter “community functional traits”). The trait values were obtained from the sources mentioned above (19, 25–31). We then modeled each community functional trait as a function of the additive effects of relevant and largely uncorrelated climatic drivers of species distributions (fig. S1), i.e., the mean annual values of temperature ( $T_{\text{mean}}$ ), vapor pressure deficit ( $VPD_{\text{mean}}$ ) (51), maximum climatic water deficit ( $MCWD_{\text{mean}}$ ) (52), and the standardized precipitation-evapotranspiration index ( $SPEI_{12}$ ) (53), with each of these interacting with forest type (lowland or montane). Because soil characteristics can affect plant distributions (24), we included cation exchange capacity (CEC), pH, and the percentage of clay and sand for

each plot location in the models [see the materials and methods (35)]. We accounted for differences in the number of censuses, plot size, and census time per vegetation plot and for the potential spatial autocorrelation.

Several community functional traits show consistent relationships with climate across forest type (table S2 and fig. S2), with  $T_{\text{mean}}$  showing some of the strongest effects driving plant trait distributions across lowland and montane forests (Fig. 3). As expected, an increase in  $T_{\text{mean}}$  across space is associated with an increase in community-mean leaf area and SM and a decrease in  $A_{\text{sat}}$ , SLA, and the proportion of DE across lowland and montane forests. Moreover, an increase in water stress ( $MCWD_{\text{mean}}$ ) is associated with decreases in SLA and  $H_{\text{max}}$  for both forest types (table S2 and fig. S2). This represents an increase in the conservative trait strategy linked to more extreme conditions.

The relationship with  $T_{\text{mean}}$  is not, however, consistent across lowland and montane forests for leaf chemistry (C, N, and P content), WD,  $H_{\text{max}}$ , leaf FM, or leaf thickness (Fig. 3). An increase in water stress ( $MCWD_{\text{mean}}$ ) is associated with an increase in  $A_{\text{sat}}$ , leaf N content,



**Fig. 3. Relationship between community-mean plant traits and temperature.** Trait-environment relationships for  $T_{\text{mean}}$  across the vegetation plots. Thick blue lines (for lowland forests) and yellow lines (for montane forests) show the average trait response to the climatic variable, with gray-shaded lines show 700 random draws from the model posterior distribution representing the variability of the expected model fit. Trait-environment relationships for mean maximum MCWD, VPD, and SPEI ( $\text{MCWD}_{\text{mean}}$ ,  $\text{VPD}_{\text{mean}}$ , and  $\text{SPEI}_{\text{mean}}$ , respectively) are shown in fig. S2. For full statistical multivariate model results, see table S2.

leaf area, and WD across lowland forests but decreases in montane forests (table S2 and fig. S2). The increase in these leaf traits in drier forests could be associated with the high photosynthetic rates generally attained by DE over the growing season (54, 55) and the fact that lower  $H_{\text{max}}$  and higher WD tend to correlate with higher resistance to lethally low levels of soil moisture availability (37). However, consistent climatic relationships across both forest types are not apparent for the other traits analyzed (table S2 and fig. S2). One plausible explanation is that this reflects their different position along the climatic gradient (i.e.,  $T_{\text{mean}}$  and precipitation), with lowlands occupying areas with more homogeneous climate conditions across large spatial extents compared with montane forests, which span a large range of climates across smaller spatial extents.

#### Changes in trait composition across time

We next investigated whether and how the functional trait composition of tropical American forests has shifted and how much of this can be explained by the observed changes in climate over the past 40 years. We first calculated the CWM of each plant functional trait for each vegetation census available for full community assemblage and separately for the survivor assemblage (individuals that are alive in two subsequent censuses, e.g., from census

one to census two), recruit assemblage (individuals not present in the previous census and recruited in the subsequent census), and fatality assemblage (individuals alive in previous census but dead in the subsequent census). We defined the recruit assemblage as individuals that passed the threshold of 10 cm DBH between one census and the next. We then calculated their yearly rate of change across time. We tested whether the changes in trait CWM differed from zero across all vegetation plots, with plots separated into lowland and montane forests. We calculated the highest density interval (HDI) containing the 95% most probable effect values and considered it significant when the HDI did not overlap 0. We then investigated whether the observed shifts in trait CWM differed significantly between lowland and montane forests. For shorthand and readability, all mention of mean traits and shifts below refer to CWM trait values.

When considering all plots together for the full community assemblage, we found that seven of the 12 traits analyzed exhibited significant changes in their CWM values (fig. S3; see Fig. 4 for trait changes across assemblages). Only leaf N, FM, SLA, SM, and WD did not show significant shifts across time (table S3 and fig. S4). The survivor assemblage showed the same pattern of community trait changes (table S3 and Fig. 5) as the full community assem-


blage, with the main differences being a significant decrease in leaf FM in the lowlands for the survivor assemblage. Thus, hereafter, we focus on the results from the survivor, recruit, and fatality assemblages. Overall, we found larger variation in trait CWM across space (i.e., with geographical variation in climate) than across time. For the community traits with significant changes for the survivor assemblage, we found an average increase in  $A_{\text{sat}}$  of  $0.0023 \mu\text{mol m}^{-2} \text{s}^{-1} \text{year}^{-1}$  (HDI-low and HDI-high: 0.0007 and 0.0038, respectively), leaf C content  $0.0011\% \text{year}^{-1}$  (0.0004 and 0.0019), leaf P content  $1.6 \times 10^{-5}\% \text{year}^{-1}$  ( $5.7 \times 10^{-6}$  and  $2.7 \times 10^{-5}$ ), abundance of DE  $0.03\% \text{year}^{-1}$  (0.01 and 0.05), and  $H_{\text{max}}$   $0.006 \text{m year}^{-1}$  (0.002 and 0.009), whereas community leaf area decreased on average  $-0.03 \text{cm}^2 \text{year}^{-1}$  ( $-0.06$  and  $-0.007$ ) and leaf thickness decreased  $-0.05 \text{mm year}^{-1}$  ( $-0.08$  and  $-0.02$ ) (table S3 and Fig. 5). In the lowland forests, we detected significant trait changes for six (increasing:  $A_{\text{sat}}$ , leaf C content,  $H_{\text{max}}$ , and abundance of DE; decreasing: leaf area and FM) of the 12 traits analyzed (table S3 and Fig. 5). Montane forests showed significant, but rather small, increases in leaf C and P content, and the abundance of DE (table S3 and Fig. 5).

The recruit assemblage experienced significant changes for seven traits, with six showing decreases, i.e., leaf C content  $-0.014\% \text{year}^{-1}$



**Survivor assemblage****<8% tracking**


↑  $A_{sat}$ , C, P, DE,  $H_{max}$   
 ↓ Area, Fresh mass, Thickness

**Recruit assemblage****<22% tracking**

↑ Fresh mass  
 ↓ C, N, Thickness, DE  
 $H_{max}$ , WD

**Fatality assemblage**

↓ N, Fresh mass  
 Seed mass



**Fig. 4. The analyzed survivor (top panel), recruit (middle panel), and fatality (bottom panel) assemblages in the study.** In each panel, the highlighted vegetation represents the specific assemblage under analysis. Each panel provides a summary of observed changes in community traits and the percentage of climate tracking by each assemblage, with exception of the fatality assemblage, for which climate tracking is not possible.

(HDI-low and HDI-high:  $-0.02$  and  $-0.001$ , respectively, in montane forests), leaf N content  $-0.002\% \text{ year}^{-1}$  ( $-0.004$  and  $-0.0002$ ), leaf thickness  $-0.04 \text{ mm year}^{-1}$  ( $-0.08$  and  $-0.01$ ), DE  $-0.17\% \text{ year}^{-1}$  ( $-0.33$  and  $-0.02$ ),  $H_{max}$   $-0.03 \text{ m year}^{-1}$  ( $-0.07$  and  $-0.003$ ), and WD  $-0.0007 \text{ g cm}^3 \text{ year}^{-1}$ . The leaf FM of recruits increased on average  $0.04 \text{ g year}^{-1}$  ( $0.006$  and

$0.08$ ; table S3 and Fig. 5). For the fatality assemblage, only the CWM of leaf N content  $-0.004\% \text{ year}^{-1}$  ( $-0.007$  and  $-0.001$  in montane forests), leaf FM  $-0.02 \text{ g year}^{-1}$  ( $-0.05$  and  $-0.0003$ ) and SM  $-17.7 \text{ mg year}^{-1}$  ( $-29.9$  and  $-5.7$ ) in lowland forests experienced significant declines (table S3 and Fig. 5).

To help identify the underlying climatic drivers of forest functional change, we used multivariate linear models to estimate the  $\Delta r$  (i.e., from the first to the last census) in the trait values ( $\Delta r$ , trait CWM) as a function of the yearly rate of change in temperature  $T_r$  ( $\Delta T_r$ ), MCWD ( $\Delta \text{MCWD}_r$ ), SPEI ( $\Delta \text{SPEI}_r$ ), and VPD ( $\Delta \text{VPD}_r$ ), each of these interacting with forest type, and accounted for soil characteristics by including in the models the CEC, pH, and clay and sand content (maps in figs. S3 to S8). Our results for the full community assemblage, the survivor assemblage, and the recruit and fatality assemblages (table S4) demonstrate the role of climate, specifically temperature and water availability, as a determinant of trait shifts across the forests and show the differences in response between lowland and montane forests (table S4). Our mapped model predictions (maps in figs. S3 to S8) depict in a spatially explicit way areas where stable CWM trait values (light yellow and light blue) and their increases (darker blue) or decreases (yellow to red) are predicted to have occurred across tropical American forests with some of the strongest CWM trait shifts predicted across forests in Amazonia.

### Can tropical American forest functional composition track climate change?

We next investigated whether the observed community trait changes are sufficient to maintain the expected trait-environment relationships for the full community, survivor, and recruit assemblages based on spatial relationships between traits and climate. We expected recruitment to be more sensitive to climate change because the full community is dominated by the demographic inertia of established adult trees. To quantify the trait changes that would be necessary for forest communities to track predicted climate change, we first quantified the relationship between community traits and environment before most anthropogenic climate changes occurred (1980 to 2005; i.e., as baseline CWM trait-environment relationships). We took our observed trait-climate relationships (built with the 1980 to 2005 period data; table S5) and used them to predict the trait CWM to the 1980 to 2005 climate conditions plus the observed changes in climate across the study sites for the full time period (the past 40 years). This allowed us to predict the CWM trait values that the forests would have if they fully tracked recent climate change, assuming that trait-climate relationships are similar across space and time (table


S6 and table S7). The ratio between the observed and expected changes (for the full and the recruit assemblages) indicates how closely these forest traits are tracking our climate equilibrium predictions based on community changes alone (Fig. 6).

Our results show that for all measured traits of the survivor and full community assemblages, the community trait composition is not changing sufficiently to track climate change, with most changes being rather small and unlikely to represent important impacts on ecosystem functioning. However, the recruit community shows the largest shifts (Figs. 4 and 6; results for all assemblages are in fig. S9). At the region-wide scale for the survivor assemblages, all traits show  $<8\%$  for lowland forests and  $<4\%$  for montane forests of the change required to track climate. For the full community assemblage, all traits show  $<6\%$  of the climate-predicted shifts in the expected direction for lowland forests and  $<7\%$  for montane forests of the expected change (fig. S9 and tables S6 and S7). Several traits show very little change or even modest changes in the opposite direction from those expected (Fig. 6, A and B). We detected larger community trait shifts in the recruit assemblages of an average 21.8% of the change required for lowland forests and 17.5% for montane forests when only traits shifting in the expected direction are considered. When both shifts in the expected direction and those in the opposite direction are considered, the recruit assemblage shows an average shift of 11.4% for lowland forests and  $-0.67\%$  for montane forests (Fig. 6, C and D, and tables S6 and S7). In lowland forests, community mean WD appears to be changing fast enough in the recruit assemblages to track climate change expectation. Overall, we see some evidence of how the recruit forest assemblages of lowland and montane forests are shifting their community traits, often for different sets of community mean trait values, in response to climate change. However, for most traits, even the recruit community does not seem to be changing quickly enough to track climate change. More significant community trait shifts have occurred in lowland forests than in montane forests, which is consistent with a more rapidly drying climate in the former (table S3 and Fig. 5).


### Discussion

Overall, we found that (i) trait-environment relationships are similar for most of the studied traits across lowland and montane tropical American forests; (ii) lowland forests show significant and larger changes in more community traits analyzed than do montane forests; (iii) across the forests and for the full community and survivor assemblages, the abundance of DE is increasing, with accompanying increases in leaf  $A_{sat}$  and decreases


### A Leaf photosynthetic capacity and chemistry



### B Leaf morphology and structure



### C Tree phenology, structure and dispersal



**Significance** ○ Non-significant ● Significant

**Assemblage** ● Survivor ● Recruit ● Fatality

**Fig. 5. Estimated changes in mean community functional trait values across time for tropical American forests.** All traits with their spatial prediction maps are shown in figs. S3 to S8. (A to C) Changes in trait CWM for leaf  $A_{\text{sat}}$  and leaf chemistry traits (A), for leaf morphology and structural traits (B), and tree phenology and structural traits (C). Each panel shows the observed yearly rate of change obtained from sampled vegetation plots from the statistical models in table S3 for all forests together and only for lowland or montane forests for the survivor (blue), recruit (green), and fatality (gray) assemblages. Significant shifts are shown as filled circles and nonsignificant as empty circles. The vertical lines depict the highest density interval (95% HDI), and the horizontal gray dotted line indicates zero change.

in leaf area and leaf thickness, yet the recruit communities in the lowland forests have on average decreased in the abundance of DE, leaf N content, and WD; and (iv) crucially, for the full tree community and survivor assemblages, most of these traits are changing at only a fraction of the rate required to maintain equilibrium with climate. The recruit communities show the best tracking of a changing climate.


The community trait shifts were similar for the survivor and full community assemblages and, although significant in several cases, these

have been rather small over the past 40 years. In general, such community trait changes differ from those of the recruit and fatality assemblages. This is likely because the trait shift responses of the survivor and full community assemblages are dominated by large individuals that continued growing throughout the study period. Another potential explanation is that the survivor and full community assemblages, along with their concurrent functional trait composition, are still able to withstand the observed changes in climate. The survivor and full community assemblages have shifted

toward more deciduous communities with higher  $A_{\text{sat}}$ , leaf chemistry, and  $H_{\text{max}}$ . At the same time, we uncovered a general decrease in leaf thickness for the survivor and recruit assemblages. Temporal increases in VPD have potentially favored increases in the proportion of DE, especially across montane forests, and increases in MCWD partially explain decreases in leaf thickness. Overall, DE tend to have acquisitive leaf traits with higher leaf N and P contents,  $A_{\text{sat}}$ , and photosynthetic N-use efficiency, especially under water stress (56), than do evergreen species (57, 58). The pattern



**Fig. 6. (A to D) Tracking of trait CWM for the survivor [(A) and (B)] and recruit [(C) and (D)] assemblages in lowland [(A) and (C)] and montane [(B) and (D)] forests given the observed changes in climate across the sampling plots.** The x axis shows the ratio of changes in trait CWM based on actual trait CWM changes observed at the plot level through time, versus expected changes in trait CWM based on spatial climate-trait relationships given observed changes in climate. Positive values (black bars) indicate that observed and predicted changes are both positive or both negative and thus are going into the same direction, whereas negative values (gray bars) indicate that observed and predicted changes are going in opposite directions. A ratio of change value of one would indicate perfect tracking. The y axis shows the traits sorted by the change ratio amount (see full statistical details in tables S6 and S7). Values of zero and close to zero represent no or slight trait shifts.



observed across tropical American forests could be attributable to leguminous N-fixing species that dominate in dry forests, which are often deciduous and have higher photosynthetic N-use efficiency (59). This is consistent with a previous report for West African tropical forests, where increasing drought stress co-occurred with an increased abundance of DE, and changes in DE explained changes in other morphological, structural, and leaf chemistry traits (55). The abundance of DE may be limited by soil fertility (60) in areas such as in southeastern Amazonia (more so the Guiana Shield), where short-lived deciduous leaf construction is too costly. Thus, increase in DE is expected to be one adaptation strategy, especially in dry tropical forests with more seasonal precipitation regimes and nutrient-rich soils than wetter tropical forests.

There is a mismatch in trait responses to climate change between the recruit assemblage and both the full community and survivor assemblages. This mismatch is most pronounced with respect to the abundance of DE, leaf C content, and  $H_{max}$ . With increasing temperatures and reduced water availability, we expected an increase in abundance of DE to also be reflected in the recruit assemblage (55). However, the decline in abundance of DE in the recruit assemblage indicates potential shifts in phenological strategies toward more conservative strategies in response to increasing temperatures or altered precipitation patterns.

The recruit assemblages also select for lower leaf C content and species with shorter  $H_{max}$ . This finding suggests a decoupling in trait space between the functional trait characteristics of the mature forests that we see in the present and the possible future functional composition of tropical American forests. The selection for low leaf N content in the recruit and fatality assemblages raises the question of whether and to what extent such recruit assemblages will be able to survive to larger adult sizes (e.g., 56, 61), especially across montane forests, where there is a stronger mismatch. Such a decoupling in trait space between the recruit and survivor assemblages could potentially indicate the slow beginnings of forest-level adjustment to new climatic conditions, which is likely to affect the functioning of tropical forest ecosystems (62). We did not find a significant selection against DE in the fatality assemblage. This suggests that a combination of drought avoidance and drought resistance strategies (38) could both be playing an important role in adaptation to a warming climate across lowland and montane tropical forests.

Other factors may be promoting the observed change in community-mean traits, such as species interactions and defaunation, the latter being a potentially important driver of changes in dispersal traits across time (63). Some wetter regions (e.g., central Amazonia) show slight increases in SM for the full community (fig. S4D), with the fatality assemblage showing

significant declines in individuals with smaller seeds in the lowlands (Fig. 5). However, drier regions (e.g., the southern and eastern fringes of Amazonia) and montane forests show a slight predicted decline in SM (fig. S4D). These changes may be an indicator of defaunation pressure (64), because spatial predictions of decreases in SM broadly match spatial patterns of high defaunation (65), especially in those more accessible areas of Mesoamerica and in both southern and eastern Brazil. They could also be driven by climatic factors because the observed changes are consistent with a shift from endozoochory (animal dispersal) to anemochory (wind dispersal), with the latter exhibiting smaller seeds than those dispersed by animals and being more prevalent in drier biomes (41). Including other relevant traits, such as those related to hydraulics and thermal tolerance, and considering ecological interactions could further bring new evidence of these potential forest adjustments to a changing climate.

The survivor, full community, and recruit assemblages often show more changes in traits in lowland than montane forest. Lowland forests are highly dynamic and harbor a high functional trait diversity that potentially allows for selection from a wider pool of trait values under climate stress. There has been a larger increase in atmospheric VPD in lowland forests than in montane forests caused by more pronounced increases in temperature over the

past 40 years, which could partially explain the shift of a larger number of community functional traits in lowland forests than in montane forests (66). Larger increases in VPD and more severe droughts appear to have modified the community composition of lowland forests more strongly than that of montane forests toward a set of species better adapted to drier and hotter conditions, which could be due to the mortality of more vulnerable species (57). Recent work across sites in the Amazon and the Andes also suggest an important impact of increasing temperatures and declines in water availability on tree trait composition (67). We investigated the impact of macroclimate on the changes in functional trait composition of tropical forests. However, such macroclimate conditions may not directly mirror the microclimatic conditions, such as temperature, found under the forest canopy (68). This is of particular importance when investigating the effects of a changing climate, especially on the recruit assemblages, which tend to occupy the space below the canopies of the older, larger trees. Ultimately, such microclimatic conditions may play an important role in determining the responses of understory plants to a changing climate (45, 69, 70) and therefore the rate of change in community trait composition of the recruit assemblages. Therefore, microclimatic conditions at the plot level may partly explain the differences in trait shifts between the full community and survivor assemblages and the recruit assemblages.

It would mechanistically be expected that increasing drought would cause plant communities to shift to species with higher WD and thicker leaves or that the abundance of DE would increase across time. Such coordinated changes may not readily happen in the community because it is whole phenotypes that are changing, i.e., particular combinations of traits rather than isolated traits. Moreover, coordination of different strategies could allow for alternative adaptations to the same drivers. For example, drier conditions might encourage DE combined with low WD and thin leaves (drought avoidance) or evergreenness combined with high WD and thicker leaves (drought tolerance). The favored combination(s) may depend on forest seasonality patterns and soil nutrients. Furthermore, not all trait combinations may be present in any given regional species pool, even in species-rich biomes, which may limit the shifts in community traits that can occur at any given time as a response to environmental change. Other factors may also contribute to trait shifts or a lack thereof across forest communities. These include soil conditions (71), biotic interactions (e.g., animal-plant interactions) (72), and wind disturbance (73). Our analyses represent community-wide responses mainly based on trait information at the species and

genus level; traits may also express intraspecific plasticity that we are unable to assess here given the scale and multidecadal nature of the study. Some traits may show more or less plasticity than others, and species-intraspecific variation may contribute to adaptation to a changing climate (74, 75). Overall, there is a lack of knowledge and data on the extent to which intraspecific trait variation plays a role in the adaptation of tree communities to a changing climate across the tropics. Here, we analyzed only a set of relevant plant functional traits without adding information on intraspecific trait variation. Further research could focus on understanding responses of tree communities to climate change, including as much as possible information on intraspecific trait variation, and analyzing other relevant traits. These could be hydraulic and thermal tolerance traits, which at the moment are not widely available across tropical American forests.

In conclusion, we found that overall changes in community trait composition are leading to small shifts amounting to only ~10% of the expectation given climate change. These shifts are primarily driven by variation in growth rates of existing trees rather than by recruitment or tree mortality. However, we observed larger changes for the recruit assemblage, directionally tracking climate at an average of 21%, which can potentially contribute to keeping these forests closer to, although still far from, the equilibrium with climate. Trees are long-lived organisms with slow turnover rates compared with the rate of climate change, and this partly explains the differences observed in community trait shifts between the full community and those of the recruit assemblages. There are specific areas where there seems to be a larger lag in forest responses to climate changes, especially in the Maya Forest in Mesoamerica (76) and both the Atlantic Forest and the southern Amazon Forest in Brazil (77), which have become increasingly fragmented over time. Consequently, the impacts of other disturbances across these regions, such as habitat fragmentation and in general a more constrained physical environment, may be affecting the capacity of forests to adjust to new climate conditions (43, 78). Our analysis demonstrates that tree community composition is shifting to track climate change, but that the overwhelming onus would have to be on within-species variability and trait plasticity (79, 80) to adequately do so. However, the changes in climate are likely to be too fast for adaptive phenotypic plasticity to keep track, especially in environments with low climatic heterogeneity (70, 80). Thus, it is overwhelmingly likely that the tree species composition and functional properties of tropical American forests (and probably all tropical forests) are increasingly out of equilibrium with local climate.

Such disequilibrium almost certainly increases vulnerability to a further changing climate.

## Methods summary

### *Understanding trait CWM-climate relationships and the effects of climate change on driving trait CWM changes*

To understand the current trait-climate relationships across forests of the tropical Americas, for each plant trait, we modeled the trait CWM as a function of climatic and soil covariates, with each one of the climatic variables interacting with forest type (lowland or montane) (hereafter referred to as M1). We next analyzed the climatic drivers of shifts in each functional trait given observed changes in climate over the past 40 years for the full community and survivor assemblages, for the recruit community, and for the fatality community. The fatality community is defined as those individuals of a plot that were alive in a previous census but dead in the following census. We calculated the temporal changes in trait CWM at the plot level as the  $\Delta r$  to standardize for a different time between censuses for different plots. We then modeled the  $\Delta r$  CWM trait as a function of  $\Delta r$  of the climatic variables described above, with each of these interacting with forest type, and also included the soil characteristics (hereafter referred to as M2).

### *Understanding shifts in trait CWM*

We used the  $\Delta r$  of the trait CWM of the full, survivor, recruit, and fatality community assemblages to investigate whether the rate of trait changes for the overall forests (lowland and montane together), for the lowland forests alone, and for the montane forest alone was significantly different from 0. We did the same to determine whether there were important differences between the rate of change between lowland and montane forests. To this end, we performed a version of a typical  $t$  test analysis using Bayesian estimation (81, 82). As above, here, we calculated the HDI containing the 95% most probable effect values and considered a result significant when the HDI did not overlap 0.

### *Understanding whether forest community traits are tracking climate changes*

The process outlined below was performed only for the full community assemblage and the survivor and recruit assemblages because the fatality assemblages are not tracking climate. We first built the same type of statistical models as M1 but using only plot and climatic data from between 1980 and 2005, including also the soil variables (hereafter referred to as M1.1). We used the M1.1 trait-environment statistical models and obtained predictions of the trait CWM to a new set of climatic conditions composed of the 1980 to 2005 climate plus the observed

climate yearly rate of change across the study period (hereafter referred to as M2). We then calculated the difference between the trait CWM obtained with the M1.1 and M2 models to obtain the expected trait CWM change. Finally, we compared the expected trait CWM calculated above with the observed  $\Delta r$  CWM trait. This allowed us to understand the expected shift in mean trait values given the 1980 to 2005 trait-climate relationship compared with the observed trait changes across time (i.e., from 1980 to 2021). We tested for significant differences between observed and expected community trait changes using Bayesian estimation (81, 82). We also created map predictions of the 1980 to 2005 M1.1 trait-climate model across tropical American forests by predicting this model to a climate change scenario that was composed of the observed climate (1980 to 2005) plus the yearly rate change observed. We then subtracted the original map predictions (i.e., those made with the M1.1 models without changes in climate conditions) to obtain the expected CWM trait changes at the pixel level (in the map) for across forests in tropical America. Next, we calculated the ratio of the observed, i.e., spatial predictions of the trait changes observed across time (from M2 models) versus the expected and converted to percentage change relative to the 1980 to 2005 condition to determine whether and to what extent the observed trait changes are tracking (values above zero) or not (values of zero) the expected changes given the observed changes in climate or shifting in opposite direction than expected (values below zero).

## REFERENCES AND NOTES

- E. Beech, M. Rivers, S. Oldfield, P. P. Smith, GlobalTreeSearch: The first complete global database of tree species and country distributions. *J. Sustain. For.* **36**, 454–489 (2017). doi: [10.1080/10549811.2017.1310049](https://doi.org/10.1080/10549811.2017.1310049)
- N. Myers, R. A. Mittermeier, C. G. Mittermeier, G. A. B. da Fonseca, J. Kent, Biodiversity hotspots for conservation priorities. *Nature* **403**, 853–858 (2000). doi: [10.1038/35002501](https://doi.org/10.1038/35002501); pmid: [10706275](https://pubmed.ncbi.nlm.nih.gov/10706275/)
- UN Environment Programme (UNEP), Food and Agriculture Organization of the United Nations (FAO), "The state of the world's forests: Forests, biodiversity and people" (UNEP, FAO, 2020); <https://www.unep.org/resources/state-worlds-forests-forests-biodiversity-and-people>.
- D. M. Lapola et al., The drivers and impacts of Amazon forest degradation. *Science* **379**, eabp8622 (2023). doi: [10.1126/science.abp8622](https://doi.org/10.1126/science.abp8622); pmid: [36701452](https://pubmed.ncbi.nlm.nih.gov/36701452/)
- J. S. Albert et al., Human impacts outpace natural processes in the Amazon. *Science* **379**, eabo5003 (2023). doi: [10.1126/science.abo5003](https://doi.org/10.1126/science.abo5003); pmid: [36701466](https://pubmed.ncbi.nlm.nih.gov/36701466/)
- S. R. Loarie et al., The velocity of climate change. *Nature* **462**, 1052–1055 (2009). doi: [10.1038/nature08649](https://doi.org/10.1038/nature08649); pmid: [20033047](https://pubmed.ncbi.nlm.nih.gov/20033047/)
- A. Esquivel-Muelbert et al., Compositional response of Amazon forests to climate change. *Glob. Chang. Biol.* **25**, 39–56 (2019). doi: [10.1111/gcb.14413](https://doi.org/10.1111/gcb.14413); pmid: [30406962](https://pubmed.ncbi.nlm.nih.gov/30406962/)
- B. Fadrique et al., Widespread but heterogeneous responses of Andean forests to climate change. *Nature* **564**, 207–212 (2018). doi: [10.1038/s41586-018-0715-9](https://doi.org/10.1038/s41586-018-0715-9); pmid: [30429613](https://pubmed.ncbi.nlm.nih.gov/30429613/)
- D. A. Prieto-Torres, A. G. Navarro-Sigüenza, D. Santiago-Alarcon, J. R. Rojas-Soto, Response of the endangered tropical dry forests to climate change and the role of Mexican Protected Areas for their conservation. *Glob. Chang. Biol.* **22**, 364–379 (2016). doi: [10.1111/gcb.13090](https://doi.org/10.1111/gcb.13090); pmid: [26367278](https://pubmed.ncbi.nlm.nih.gov/26367278/)
- S. N. Aitken, S. Yeaman, J. A. Holliday, T. Wang, S. Curtis-McLane, Adaptation, migration or extirpation: Climate change outcomes for tree populations. *Evol. Appl.* **1**, 95–111 (2008). doi: [10.1111/j.1752-4571.2007.00013.x](https://doi.org/10.1111/j.1752-4571.2007.00013.x); pmid: [25567494](https://pubmed.ncbi.nlm.nih.gov/25567494/)
- B. V. Smithers, M. P. North, C. I. Millar, A. M. Latimer, Leap frog in slow motion: Divergent responses of tree species and life stages to climatic warming in Great Basin subalpine forests. *Glob. Chang. Biol.* **24**, e442–e457 (2018). doi: [10.1111/gcb.13881](https://doi.org/10.1111/gcb.13881); pmid: [28850759](https://pubmed.ncbi.nlm.nih.gov/28850759/)
- K. Zhu, C. W. Woodall, J. S. Clark, Failure to migrate: Lack of tree range expansion in response to climate change. *Glob. Chang. Biol.* **18**, 1042–1052 (2012). doi: [10.1111/j.1365-2486.2011.02571.x](https://doi.org/10.1111/j.1365-2486.2011.02571.x)
- C. Huntingford et al., Simulated resilience of tropical rainforests to CO<sub>2</sub>-induced climate change. *Nat. Geosci.* **6**, 268–273 (2013). doi: [10.1038/ngeo1741](https://doi.org/10.1038/ngeo1741)
- E. J. Castellanos et al., "Central and South America," in *Climate Change 2022: Impacts, Adaptation, and Vulnerability. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, H.-O. Pörtner et al., Eds. (Cambridge Univ. Press, 2022); pp. 1689–1816; <https://www.ipcc.ch/report/ar6/wg2/chapter/chapter-12/>.
- Intergovernmental Panel on Climate Change, "Climate change and land: An IPCC special report on climate change, desertification, land degradation, sustainable land management, food security, and greenhouse gas fluxes in terrestrial ecosystems – Summary for Policymakers" (IPCC, 2019); <https://wedocs.unep.org/handle/20.500.11822/29261>.
- C. H. Trisos, C. Merow, A. L. Pigot, The projected timing of abrupt ecological disruption from climate change. *Nature* **580**, 496–501 (2020). doi: [10.1038/s41586-020-2189-9](https://doi.org/10.1038/s41586-020-2189-9); pmid: [32322063](https://pubmed.ncbi.nlm.nih.gov/32322063/)
- J. W. Williams, S. T. Jackson, Novel climates, no-analog communities, and ecological surprises. *Front. Ecol. Environ.* **5**, 475–482 (2007). doi: [10.1890/070037](https://doi.org/10.1890/070037)
- C. Violle et al., Let the concept of trait be functional! *Oikos* **116**, 882–892 (2007). doi: [10.1111/j.0030-1299.2007.15559.x](https://doi.org/10.1111/j.0030-1299.2007.15559.x)
- S. Díaz et al., The global spectrum of plant form and function. *Nature* **529**, 167–171 (2016). doi: [10.1038/nature16489](https://doi.org/10.1038/nature16489); pmid: [26700811](https://pubmed.ncbi.nlm.nih.gov/26700811/)
- C. Fortunel, C. E. Paine, P. V. Fine, N. J. Kraft, C. Baraloto, Environmental factors predict community functional composition in Amazonian forests. *J. Ecol.* **102**, 145–155 (2014). doi: [10.1111/1365-2745.12160](https://doi.org/10.1111/1365-2745.12160)
- B. J. Enquist et al., "Scaling from traits to ecosystems: Developing a general trait driver theory via integrating trait-based and metabolic scaling theories," in *Advances in Ecological Research*, S. Pawar, G. Woodward, A. I. Dell, Eds. (Elsevier, vol. 52, 2015); pp. 249–318.
- J. Honeier, T. Seeler, K. Pierick, C. Leuschner, Leaf trait variation in species-rich tropical Andean forests. *Sci. Rep.* **11**, 9993 (2021). doi: [10.1038/s41598-021-89190-8](https://doi.org/10.1038/s41598-021-89190-8); pmid: [33976239](https://pubmed.ncbi.nlm.nih.gov/33976239/)
- N. Madani et al., Future global productivity will be affected by plant trait response to climate. *Sci. Rep.* **8**, 2870 (2018). doi: [10.1038/s41598-018-21172-9](https://doi.org/10.1038/s41598-018-21172-9); pmid: [29434266](https://pubmed.ncbi.nlm.nih.gov/29434266/)
- H. Bruelheide et al., Global trait-environment relationships of plant communities. *Nat. Ecol. Evol.* **2**, 1906–1917 (2018). doi: [10.1038/s41559-018-0699-8](https://doi.org/10.1038/s41559-018-0699-8); pmid: [30455437](https://pubmed.ncbi.nlm.nih.gov/30455437/)
- N. G. Swenson, M. N. Umana, "Data from: Interspecific functional convergence and divergence and intraspecific negative density dependence underlie the seed-to-seedling transition in tropical trees," Dryad (2015); <https://datadryad.org/dataset/doi:10.5061/dryad.j2r53>.
- R. Muscarella, M. Uriarte, Do community-weighted mean functional traits reflect optimal strategies? *Proc. Biol. Sci.* **283**, 20152434 (2016). doi: [10.1098/rspb.2015.2434](https://doi.org/10.1098/rspb.2015.2434); pmid: [27030412](https://pubmed.ncbi.nlm.nih.gov/27030412/)
- M. N. Umaña et al., Interspecific functional convergence and divergence and intraspecific negative density dependence underlie the seed-to-seedling transition in tropical trees. *Am. Nat.* **187**, 99–109 (2016). doi: [10.1086/684174](https://doi.org/10.1086/684174); pmid: [27277406](https://pubmed.ncbi.nlm.nih.gov/27277406/)
- Y. Malhi et al., The Global Ecosystems Monitoring network: Monitoring ecosystem productivity and carbon cycling across the tropics. *Biol. Conserv.* **253**, 108889 (2021). doi: [10.1016/j.biocon.2020.108889](https://doi.org/10.1016/j.biocon.2020.108889)
- ForestPlots.net et al., Taking the pulse of Earth's tropical forests using networks of highly distributed plots. *Biol. Conserv.* **260**, 108849 (2021). doi: [10.1016/j.biocon.2020.108849](https://doi.org/10.1016/j.biocon.2020.108849)
- J. Kattge et al., TRY plant trait database - enhanced coverage and open access. *Glob. Chang. Biol.* **26**, 119–188 (2020). doi: [10.1111/gcb.14904](https://doi.org/10.1111/gcb.14904); pmid: [31891233](https://pubmed.ncbi.nlm.nih.gov/31891233/)
- S. Díaz et al., The global spectrum of plant form and function: Enhanced species-level trait dataset. *Sci. Data* **9**, 755 (2022). doi: [10.1038/s41597-022-01774-9](https://doi.org/10.1038/s41597-022-01774-9); pmid: [36477373](https://pubmed.ncbi.nlm.nih.gov/36477373/)
- S. Fauset et al., Drought-induced shifts in the floristic and functional composition of tropical forests in Ghana. *Ecol. Lett.* **15**, 1120–1129 (2012). doi: [10.1111/j.1461-0248.2012.01834.x](https://doi.org/10.1111/j.1461-0248.2012.01834.x); pmid: [22812661](https://pubmed.ncbi.nlm.nih.gov/22812661/)
- K. J. Feeley, S. J. Davies, R. Perez, S. P. Hubbell, R. B. Foster, Directional changes in the species composition of a tropical forest. *Ecology* **92**, 871–882 (2011). doi: [10.1890/10-0724.1](https://doi.org/10.1890/10-0724.1); pmid: [21661550](https://pubmed.ncbi.nlm.nih.gov/21661550/)
- G. M. Locosselli et al., Global tree-ring analysis reveals rapid decrease in tropical tree longevity with temperature. *Proc. Natl. Acad. Sci. U.S.A.* **117**, 33358–33364 (2020). doi: [10.1073/pnas.2003873117](https://doi.org/10.1073/pnas.2003873117); pmid: [33318167](https://pubmed.ncbi.nlm.nih.gov/33318167/)
- Materials and methods are available as supplementary materials.
- P. B. Reich, The world-wide 'fast-slow' plant economics spectrum: A traits manifesto. *J. Ecol.* **102**, 275–301 (2014). doi: [10.1111/1365-2745.12211](https://doi.org/10.1111/1365-2745.12211)
- X. Liang, Q. Ye, H. Liu, T. J. Brodribb, Wood density predicts mortality threshold for diverse trees. *New Phytol.* **229**, 3053–3057 (2021). doi: [10.1111/nph.17117](https://doi.org/10.1111/nph.17117); pmid: [33251581](https://pubmed.ncbi.nlm.nih.gov/33251581/)
- T. J. Brodribb, J. Powers, H. Cochard, B. Choat, Hanging by a thread? Forests and drought. *Science* **368**, 261–266 (2020). doi: [10.1126/science.aat7631](https://doi.org/10.1126/science.aat7631); pmid: [32299945](https://pubmed.ncbi.nlm.nih.gov/32299945/)
- M. T. Sande et al., Old-growth Neotropical forests are shifting in species and trait composition. *Ecol. Monogr.* **86**, 228–243 (2016). doi: [10.1890/15-1815.1](https://doi.org/10.1890/15-1815.1)
- H. W. Pritchard et al., "Regeneration in recalcitrant-seeded species and risks from climate change" in *Plant Regeneration from Seeds*, C. C. Baskin, J. M. Baskin, Eds. (Academic, 2022); pp. 259–273. doi: [10.1016/B978-0-12-823731-1.00014-7](https://doi.org/10.1016/B978-0-12-823731-1.00014-7)
- F. C. Souza, Características de dispersão de sementes e frutos ao longo dos biomas brasileiros: explorando tendências, predizendo e mapeando correlatos ecológicos (Universidade Federal de Lavras, 2023).
- C. Bello et al., Defaunation affects carbon storage in tropical forests. *Sci. Adv.* **1**, e1501105 (2015). doi: [10.1126/sciadv.1501105](https://doi.org/10.1126/sciadv.1501105); pmid: [26824067](https://pubmed.ncbi.nlm.nih.gov/26824067/)
- J. Lenoir et al., Species better track climate warming in the oceans than on land. *Nat. Ecol. Evol.* **4**, 1044–1059 (2020). doi: [10.1038/s41559-020-1198-2](https://doi.org/10.1038/s41559-020-1198-2); pmid: [32451428](https://pubmed.ncbi.nlm.nih.gov/32451428/)
- R. Bertrand et al., Changes in plant community composition lag behind climate warming in lowland forests. *Nature* **479**, 517–520 (2011). doi: [10.1038/nature10548](https://doi.org/10.1038/nature10548); pmid: [22012261](https://pubmed.ncbi.nlm.nih.gov/22012261/)
- Y. Malhi et al., Introduction: elevation gradients in the tropics: laboratories for ecosystem ecology and global change research. *Glob. Chang. Biol.* **16**, 3171–3175 (2010). doi: [10.1111/j.1365-2486.2010.02323.x](https://doi.org/10.1111/j.1365-2486.2010.02323.x)
- A. S. Jump, C. Mátys, J. Peñuelas, The altitude-for-latitude disparity in the range retractions of woody species. *Trends Ecol. Evol.* **24**, 694–701 (2009). doi: [10.1016/j.tree.2009.06.007](https://doi.org/10.1016/j.tree.2009.06.007); pmid: [19695735](https://pubmed.ncbi.nlm.nih.gov/19695735/)
- J. W. Dalling, K. Heineman, G. González, R. Ostertag, Geographic, environmental and biotic sources of variation in the nutrient relations of tropical montane forests. *J. Trop. Ecol.* **32**, 368–383 (2016). doi: [10.1017/S0266467415000619](https://doi.org/10.1017/S0266467415000619)
- J. T. Abatzoglou, S. Z. Dobrowski, S. A. Parks, K. C. Hegewisch, TerraClimate, a high-resolution global dataset of monthly climate and climatic water balance from 1958–2015. *Sci. Data* **5**, 170191 (2018). doi: [10.1038/sdata.2017.191](https://doi.org/10.1038/sdata.2017.191); pmid: [29313841](https://pubmed.ncbi.nlm.nih.gov/29313841/)
- L. Poggio et al., SoilGrids 2.0: Producing soil information for the globe with quantified spatial uncertainty. *Soil* **7**, 217–240 (2021). doi: [10.5194/soil-7-217-2021](https://doi.org/10.5194/soil-7-217-2021)
- J. P. Grime, Benefits of plant diversity to ecosystems: Immediate, filter and founder effects. *J. Ecol.* **86**, 902–910 (1998). doi: [10.1046/j.1365-2745.1998.00306.x](https://doi.org/10.1046/j.1365-2745.1998.00306.x)
- D. Bauman et al., Tropical tree mortality has increased with rising atmospheric water stress. *Nature* **608**, 528–533 (2022). doi: [10.1038/s41586-022-04737-7](https://doi.org/10.1038/s41586-022-04737-7); pmid: [35585230](https://pubmed.ncbi.nlm.nih.gov/35585230/)
- Y. Malhi et al., Exploring the likelihood and mechanism of a climate-change-induced dieback of the Amazon rainforest. *Proc. Natl. Acad. Sci. U.S.A.* **106**, 20610–20615 (2009). doi: [10.1073/pnas.0804619106](https://doi.org/10.1073/pnas.0804619106); pmid: [19218454](https://pubmed.ncbi.nlm.nih.gov/19218454/)
- S. M. Vicente-Serrano, S. Beguería, J. I. López-Moreno, A multiscale drought index sensitive to global warming: The standardized precipitation evapotranspiration index. *J. Clim.* **23**, 1696–1718 (2010). doi: [10.1175/2009JCLI2909.1](https://doi.org/10.1175/2009JCLI2909.1)
- A. Ishida et al., Photoprotection of evergreen and drought-deciduous tree leaves to overcome the dry season in



- monsoonal tropical dry forests in Thailand. *Tree Physiol.* **34**, 15–28 (2014). doi: [10.1093/treephys/tpt107](https://doi.org/10.1093/treephys/tpt107); pmid: [24336612](https://pubmed.ncbi.nlm.nih.gov/24336612/)
55. J. Aguirre-Gutiérrez et al., Drier tropical forests are susceptible to functional changes in response to a long-term drought. *Ecol. Lett.* **22**, 855–865 (2019). doi: [10.1111/ele.13243](https://doi.org/10.1111/ele.13243); pmid: [30828955](https://pubmed.ncbi.nlm.nih.gov/30828955/)
  56. D. A. Falk et al., Mechanisms of forest resilience. *For. Ecol. Manage.* **512**, 120129 (2022). doi: [10.1016/j.foreco.2022.120129](https://doi.org/10.1016/j.foreco.2022.120129)
  57. I. J. Wright et al., The worldwide leaf economics spectrum. *Nature* **428**, 821–827 (2004). doi: [10.1038/nature02403](https://doi.org/10.1038/nature02403); pmid: [15103368](https://pubmed.ncbi.nlm.nih.gov/15103368/)
  58. G. P. John et al., The anatomical and compositional basis of leaf mass per area. *Ecol. Lett.* **20**, 412–425 (2017). doi: [10.1111/ele.12739](https://doi.org/10.1111/ele.12739); pmid: [28198076](https://pubmed.ncbi.nlm.nih.gov/28198076/)
  59. M. Gei et al., Legume abundance along successional and rainfall gradients in Neotropical forests. *Nat. Ecol. Evol.* **2**, 1104–1111 (2018). doi: [10.1038/s41559-018-0559-6](https://doi.org/10.1038/s41559-018-0559-6); pmid: [29807995](https://pubmed.ncbi.nlm.nih.gov/29807995/)
  60. R. John et al., Soil nutrients influence spatial distributions of tropical tree species. *Proc. Natl. Acad. Sci. U.S.A.* **104**, 864–869 (2007). doi: [10.1073/pnas.0604666104](https://doi.org/10.1073/pnas.0604666104); pmid: [17215353](https://pubmed.ncbi.nlm.nih.gov/17215353/)
  61. L. Li et al., Leaf economics and hydraulic traits are decoupled in five species-rich tropical-subtropical forests. *Ecol. Lett.* **18**, 899–906 (2015). doi: [10.1111/ele.12466](https://doi.org/10.1111/ele.12466); pmid: [26108338](https://pubmed.ncbi.nlm.nih.gov/26108338/)
  62. F. de Bello et al., Functional trait effects on ecosystem stability: Assembling the jigsaw puzzle. *Trends Ecol. Evol.* **36**, 822–836 (2021). doi: [10.1016/j.tree.2021.05.001](https://doi.org/10.1016/j.tree.2021.05.001); pmid: [34088543](https://pubmed.ncbi.nlm.nih.gov/34088543/)
  63. R. W. Vaessen et al., Defaunation changes leaf trait composition of recruit communities in tropical forests in French Guiana. *Ecology* **104**, e3872 (2023). doi: [10.1002/ecy.3872](https://doi.org/10.1002/ecy.3872); pmid: [36121050](https://pubmed.ncbi.nlm.nih.gov/36121050/)
  64. D. de Paula Mateus et al., Defaunation impacts on seed survival and its effect on the biomass of future tropical forests. *Oikos* **127**, 1526–1538 (2018). doi: [10.1111/oik.05084](https://doi.org/10.1111/oik.05084)
  65. A. Benítez-López, L. Santini, A. M. Schipper, M. Busana, M. A. J. Huijbregts, Intact but empty forests? Patterns of hunting-induced mammal defaunation in the tropics. *PLOS Biol.* **17**, e3000247 (2019). doi: [10.1371/journal.pbio.3000247](https://doi.org/10.1371/journal.pbio.3000247); pmid: [31086365](https://pubmed.ncbi.nlm.nih.gov/31086365/)
  66. J. K. Green, J. Berry, P. Ciais, Y. Zhang, P. Gentile, Amazon rainforest photosynthesis increases in response to atmospheric dryness. *Sci. Adv.* **6**, eabb7232 (2020). doi: [10.1126/sciadv.abb7232](https://doi.org/10.1126/sciadv.abb7232); pmid: [33219023](https://pubmed.ncbi.nlm.nih.gov/33219023/)
  67. K. Bai, C. He, X. Wan, D. Jiang, Leaf economics of evergreen and deciduous tree species along an elevational gradient in a subtropical mountain. *AoB Plants* **7**, plv064 (2015). doi: [10.1093/aobpla/plv064](https://doi.org/10.1093/aobpla/plv064); pmid: [26056133](https://pubmed.ncbi.nlm.nih.gov/26056133/)
  68. P. de Frenne et al., Forest microclimates and climate change: Importance, drivers and future research agenda. *Glob. Chang. Biol.* **27**, 2279–2297 (2021). doi: [10.1111/gcb.15569](https://doi.org/10.1111/gcb.15569); pmid: [33725415](https://pubmed.ncbi.nlm.nih.gov/33725415/)
  69. J. M. Alexander et al., Lags in the response of mountain plant communities to climate change. *Glob. Chang. Biol.* **24**, 563–579 (2018). doi: [10.1111/gcb.13976](https://doi.org/10.1111/gcb.13976); pmid: [29112781](https://pubmed.ncbi.nlm.nih.gov/29112781/)
  70. F. Zellweger et al., Forest microclimate dynamics drive plant responses to warming. *Science* **368**, 772–775 (2020). doi: [10.1126/science.aba6880](https://doi.org/10.1126/science.aba6880); pmid: [32409476](https://pubmed.ncbi.nlm.nih.gov/32409476/)
  71. J. Aguirre-Gutiérrez et al., Long-term droughts may drive drier tropical forests towards increased functional, taxonomic and phylogenetic homogeneity. *Nat. Commun.* **11**, 3346 (2020). doi: [10.1038/s41467-020-16973-4](https://doi.org/10.1038/s41467-020-16973-4); pmid: [32620761](https://pubmed.ncbi.nlm.nih.gov/32620761/)
  72. C. J. Gardner, J. E. Bicknell, W. Baldwin-Cantello, M. J. Struwig, Z. G. Davies, Quantifying the impacts of defaunation on natural forest regeneration in a global meta-analysis. *Nat. Commun.* **10**, 4590 (2019). doi: [10.1038/s41467-019-12539-1](https://doi.org/10.1038/s41467-019-12539-1); pmid: [31611554](https://pubmed.ncbi.nlm.nih.gov/31611554/)
  73. S. W. Rifai et al., Landscape-scale consequences of differential tree mortality from catastrophic wind disturbance in the Amazon. *Ecol. Appl.* **26**, 2225–2237 (2016). doi: [10.1002/eap.1368](https://doi.org/10.1002/eap.1368); pmid: [27755720](https://pubmed.ncbi.nlm.nih.gov/27755720/)
  74. A. C. Westerband, J. L. Funk, K. E. Barton, Intraspecific trait variation in plants: A renewed focus on its role in ecological processes. *Ann. Bot.* **127**, 397–410 (2021). doi: [10.1093/aob/mcab011](https://doi.org/10.1093/aob/mcab011); pmid: [33507251](https://pubmed.ncbi.nlm.nih.gov/33507251/)
  75. S. Patiño et al., Branch xylem density variations across the Amazon Basin. *Biogeosciences* **6**, 545–568 (2009). doi: [10.5194/bg-6-545-2009](https://doi.org/10.5194/bg-6-545-2009)
  76. S. Bhagwat, “The history of deforestation and forest fragmentation: A global perspective” in *Global Forest Fragmentation*, C. J. Kettle, L. P. Koh, Eds. (CABI, 2014); pp. 5–19.
  77. S. Pütz et al., Long-term carbon loss in fragmented Neotropical forests. *Nat. Commun.* **5**, 5037 (2014). doi: [10.1038/ncomms6037](https://doi.org/10.1038/ncomms6037); pmid: [25289858](https://pubmed.ncbi.nlm.nih.gov/25289858/)
  78. N. M. Haddad et al., Habitat fragmentation and its lasting impact on Earth’s ecosystems. *Sci. Adv.* **1**, e1500052 (2015). doi: [10.1126/sciadv.1500052](https://doi.org/10.1126/sciadv.1500052); pmid: [26601154](https://pubmed.ncbi.nlm.nih.gov/26601154/)
  79. A. B. Nicotra et al., Plant phenotypic plasticity in a changing climate. *Trends Plant Sci.* **15**, 684–692 (2010). doi: [10.1016/j.tplants.2010.09.008](https://doi.org/10.1016/j.tplants.2010.09.008); pmid: [20970368](https://pubmed.ncbi.nlm.nih.gov/20970368/)
  80. M. Van Kleunen, M. Fischer, Constraints on the evolution of adaptive phenotypic plasticity in plants. *New Phytol.* **166**, 49–60 (2005). doi: [10.1111/j.1469-8137.2004.01296.x](https://doi.org/10.1111/j.1469-8137.2004.01296.x); pmid: [15760350](https://pubmed.ncbi.nlm.nih.gov/15760350/)
  81. J. K. Kruschke, *Doing Bayesian Data Analysis: A Tutorial with R, JAGS, and Stan* (Academic, 2014).
  82. J. K. Kruschke, Bayesian estimation supersedes the t test. *J. Exp. Psychol. Gen.* **142**, 573–603 (2013). doi: [10.1037/a0029146](https://doi.org/10.1037/a0029146); pmid: [22774788](https://pubmed.ncbi.nlm.nih.gov/22774788/)
  83. S. Begueria, S. M. Vicente-Serrano, F. Reig, B. Latorre, Standardized precipitation evapotranspiration index (SPEI) revisited: Parameter fitting, evapotranspiration models, tools, datasets and drought monitoring. *Int. J. Climatol.* **34**, 3001–3023 (2014). doi: [10.1002/joc.3887](https://doi.org/10.1002/joc.3887)
  84. J. Aguirre-Gutiérrez, Code for: Tropical forests in the Americas are changing too slowly to track climate change, Zenodo (2024); <https://doi.org/10.5281/zenodo.14192690>.
  85. J. Aguirre-Gutiérrez, Data for: Tropical forests in the Americas are changing too slowly to track climate change, Zenodo (2024); <https://doi.org/10.5281/zenodo.13996841>.

## ACKNOWLEDGMENTS

We thank the following networks for access for long-term plot data and plant trait data: RAINFOR (Amazon Forest Inventory Network, <https://rainfor.org/en/>), GEM (Global Ecosystems Monitoring network), and MONAFORE ([forestplots.net](https://forestplots.net/)), data request 109 and 184). We also thank the Herbarium of the Centro Interdisciplinario de Investigación para el Desarrollo Integral Regional campus Durango (CIIDIR), the botanical experts in the CIIDIR and the local support of Ejidos and Comunidades in Mexico. We also thank the following individuals for contributing data and for playing key roles at different stages from data collection and curation to delivery: A. Vicentini, A. Araújo-Murakami, A. Parada Gutierrez, A. S. Lima, A. Torres-Lezama, A. Dourdan, B. Burban, C. Mendoza, D. Catchpole, D. Galiano Cabrera, D. H. Neill Asanza, D. Morton, E. Hase, E. Chavez, F. Ramirez Arevalo, G. Damasco, G. Lopez-Gonzalez, G. Bañares de Dios, H. Buttgenbach Verde, I. Mendoza Polo, J. Richardson, J. Singh, J. Lleigue, J. L. Camargo, J. Stropp, J. Engel, J. Serrano, K. Melgaco, L. Ferreira, L. G. Canesi Ferreira, L. F. Duque, L. Arroyo, M. Peña-Claros, M. Steinger, M. Salgado Vital, M. Carmozina Araújo, M. Toledo, M. Corrales Medina, M. Tobler, M. Baisie, N. Silva, P. Petronelli, P. Alvarez Loayza, P. Núñez Vargas, P. van der Hout, P. Naisso, R. Oliveira de Araújo, R. Thomas, R. Boot, R. Guillén Villarroel, R. Zagt, S. Murakami, T. Killeen, V. Chama Moscoso, V. Bezard, W. Betian, W. Castro, Y. Soto Shareva, Y. T. Huilca Aedo. This manuscript is an output of ForestPlots.net Research Projects #109 and #184 “Plant functional trait changes across space and time in South American forests”, as part of the NERC-FAPESP funded project ARBOLES (NE/S011811/1). ForestPlots.net is a meta-network and cyber-initiative developed at the University of Leeds to unite permanent plot records and support tropical forest scientists. The ForestPlots.net Collaboration and Data Request Committee for the Americas (T.R.B., E.N.H.C, A.L., O.L.P., B.S.M.) facilitated this project and associated data management. The development of RAINFOR, ForestPlots.net and data curation have been funded by several grants including NE/B503384/1, NE/N012542/1 - ‘BIO-RED’, NE/X018903/1 - ‘AMSINK’, ERC Advanced Grant 291585 - ‘T-FORCES’, NE/F005806/1 - ‘AMAZONICA’, NE/N004655/1 - ‘TREMOR’, NERC New Investigators Awards, the Gordon and Betty Moore Foundation (‘RAINFOR’, ‘MonANPeru’), ERC Starter Grant 758873 - ‘TreeMort’, and from EU Framework 5, 6, and 7. Finally, we thank these contributors who are deceased: N. Davila Cardozo, T. Erwin, A. Gentry, S. Patiño, Antonio Pena Cruz, E. Armas, and J.-P. Veillon. **Funding:** J.A.-G. was funded by the Natural Environment Research Council under an independent research fellowship (NERC grant NE/T011084/1), NERC Pushing the Frontiers (NE/Z504191/1), and the Oxford University John Fell Fund (grant 10667). Global traits collection and traits

analyses under GEM were funded by an ERC Advanced Investigator Award (GEM-TRAIT grant 321131) to Y.M. under the European Union’s Seventh Framework Programme (grant FP7/2007–2013) with additional support from NERC grant NE/D014174/1 and NE/J022616/1 for traits work in Peru and NERC grant ECOFOR (NE/K016385/1) for traits work in Santarem. Plot inventories in Peru were supported by funding from the US National Science Foundation Long-Term Research in Environmental Biology program (LTREB grant DEB 1754647) and the Gordon and Betty Moore Foundation Andes–Amazon Program. Plot inventories in Nova Xavantina (Brazil) were supported by the National Council for Scientific and Technological Development (CNPq) PQ1 grants 311027/2019-9 and 303492/2022-8 to B.H.M. and B.S.M., the Long-Term Ecological Research Program (PELD), processes 441244/2016–5 and 441572/2020-0, and the Foundation of Research Support of Mato Grosso (FAPEMAT), Project ReFor, processes 589267/2016 and PELD/FAPEMAT 0346321/2021. S.D. acknowledges support from CONICET, Universidad Nacional de Córdoba and Red Federal de Alto Impacto CONATURAR (grant 2023-102072649-APN-MCT) Argentina and the Oxford Martin School. C.A.J. acknowledges support from the Brazilian National Research Council/CNPq (PELD process 403710/2012–0), NERC, and the State of São Paulo Research Foundation/FAPESP as part of the projects Functional Gradient, PELD/BIOA and ECOFOR (processes 2003/12595-7, 2012/51509-8 and 2012/51872-5, within the BIOA/FAPESP Program–The Biodiversity Virtual Institute: COTEC/IF 002.766/2013 and 010.631/2013 permits. B.S.M. was supported by the CNPq/PELD projects 441244/2016-5 and 441572/2020-0 and CAPES project 136277/2017-0. M.S. acknowledges funding for Andes Biodiversity and Ecosystem Research Group (ABERG) plot network from the US National Science Foundation (NSF) Long-Term Research in Environmental Biology (LTREB grant 1754647), the Gordon and Betty Moore Foundation’s Andes to Amazon Initiative, and RAINFOR. E.B., J.B., and Y.M. acknowledge support from NERC under projects NE/K016431/1 and NE/S01084X/1. Y.M. is supported by the Frank Jackson Foundation and the Leverhulme Trust. The MONAFORE network in Mexico has been funded by several projects highlighting those by the National Forestry Commission (CONAFOR) and the Council of Science and Technology of the State of Durango (COCYTED); F.C.S. acknowledges the Coordination for the Improvement of Higher Education Personnel – Brazil (CAPES) (Finance Code 001), and the Brazilian National Research Council/CNPq (PDJ Process No. 152806/2024-5). B.J.E. was supported by NSF awards (grant nos 2225078 and 2225076). **Author contributions:** Conceptualization: J.A.-G., S.D., S.R., Y.M.; Data gathering: all authors; Data management: J.A.-G., A.L., O.L.P., G.P.; Methodology: J.A.-G., S.D., Y.M., S.R.; Supervision (general): J.A.-G.; Supervision (forest and traits networks): J.C.C.-R., Y.M., O.P.; Writing – original draft: J.A.-G.; Writing – review & editing: all authors. **Competing interests:** The authors declare no competing interests. **Data and materials availability:** The vegetation census and plant functional traits data that support the findings of this study are available from (28), [www.ForestPlots.net](https://www.ForestPlots.net) (29), and their other original sources. Given data sovereignty from the original data owners, raw data on vegetation censuses across time are not publicly available but can be requested by contacting all researchers through the ForestPlots (30) data request protocol described in [forestplots.net/en/join-forestplots/working-with-data](https://forestplots.net/en/join-forestplots/working-with-data). Raw climate data can be accessed through the TerraClimate database at <https://www.climateatlas.org/terraclimate.html> (48). The SPEI data can be obtained from the SPEI database at <https://spei.csic.es/> (83). The computer code used to reproduce the main findings in this manuscript (84) and the plot-level processed data (85) are archived in the Zenodo repository at <https://zenodo.org/>. **License information:** Copyright © 2025 the authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original US government works. <https://www.science.org/about/science-licenses-journal-article-reuse>

## SUPPLEMENTARY MATERIALS

[science.org/doi/10.1126/science.adf5414](https://science.org/doi/10.1126/science.adf5414)

Materials and Methods

Figs. S1 to S11

Tables S1 to S7

References (86–95)

Submitted 26 October 2023; resubmitted 25 March 2024

Accepted 8 January 2025

10.1126/science.adf5414