



Northernmost record of *Crypthelia medioatlantica* (Hydrozoa: Stylasteridae) in Icelandic deep waters

Kaveh Samimi-Namin^{1,2,3} · Narissa Bax^{4,5} · Guðmundur Guðmundsson⁶ · Íris Sampaio⁷ · Stephen D. Cairns⁸

Received: 16 July 2025 / Revised: 24 November 2025 / Accepted: 25 November 2025
© The Author(s) 2026

Abstract

The stylasterid coral *Crypthelia medioatlantica* Zibrowius & Cairns, 1992, previously reported only from the Mid-Atlantic Ridge (~23–37° N), is newly documented from Icelandic waters based on material collected by the BIOICE programme. Six dry, subfossil skeletal fragments were recovered at 1539 m depth west of Iceland in 1996, extending the known range of the species by ~28° latitude into sub-Arctic environments and representing its northernmost record. Detailed morphological analysis of coenosteal texture and cyclosystem structure confirms the identification. This discovery adds a fifth stylasterid coral species to the high-latitude North Atlantic fauna and demonstrates that *C. medioatlantica* has occurred in deep sub-Arctic benthic communities at least historically. It also highlights overlooked stylasterid diversity in under-explored regions and the value of historical collections. Combining re-examination of archived material with targeted, molecularly suitable collections will be essential to clarify the biogeographic history and conservation status of *C. medioatlantica* in the North Atlantic and to determine whether the Icelandic fragments represent an extant but unsampled population or the remnants of a past local population.

Keywords Historical collections · Taxonomy · Marine biodiversity · BIOICE programme · Biogeography · Deep-sea fauna

Communicated by S. Brix

This article is a contribution to the Topical Collection *Diversity and distribution of Cold-Water Corals in Arctic and North Atlantic waters*

Kaveh Samimi-Namin and Narissa Bax share co-first authorship.

✉ Kaveh Samimi-Namin
kaveh.samimi@naturalis.nl

¹ Marine Evolution and Ecology Group, Naturalis Biodiversity Center, P.O. Box 9517, 2300 RA Leiden, The Netherlands

² Department of Biology, University of Oxford, Oxford OX1 3SZ, UK

³ Natural History Museum, Cromwell Road, London SW7 5BD, UK

⁴ Institute of Natural Resources, Climate Research Centre, Nuuk, Greenland

⁵ Centre for Marine Socioecology, Institute for Marine and Antarctic Studies, University of Tasmania, Hobart, Australia

Introduction

The family Stylasteridae Gray, 1847 (Hydrozoa: Anthoathecata) is commonly referred to as lace corals and includes predominantly cold- and deep-water taxa (Cairns 2015). Within this family, the genus *Crypthelia* Milne Edwards & Haime, 1849 is characterized by small, delicate, often uniplanar colonies bearing cyclosystems, each partially covered by a fixed

⁶ Nature Research Institute of Iceland, Urriðahóltsstraeti 6–8, Gardabaer, Iceland

⁷ CIIMAR/CIMAR LA, Interdisciplinary Centre of Marine and Environmental Research, University of Porto, Terminal de Cruzeiros Do Porto de Leixões, Matosinhos 4450-208, Portugal

⁸ Department of Invertebrate Zoology, National Museum of Natural History, Smithsonian Institution, Washington D.C., USA




Fig. 1 Bathymetric map of the North Atlantic showing the new Icelandic deep-water occurrence of *Crypthelia medioatlantica* (red marker) and the three previously known Mid-Atlantic Ridge localities (yellow markers) from Zibrowius and Cairns (1992). Bathyme-

try from the GEBCO 2023 Grid (General Bathymetric Chart of the Oceans). The Icelandic locality (BIOICE station 2918) represents the northernmost known record of the species

lid. *Crypthelia* currently comprises 56 extant species (Cairns 2024; Schuchert et al. 2025), distributed throughout the Atlantic and Indo-Pacific. Because many species are known from relatively few localities, each new record provides important information on depth range, environmental tolerances, and the contribution of *Crypthelia* colonies to deep-sea habitat complexity alongside other framework-forming corals. Although *Crypthelia* species occur across broad geographic and bathymetric scales, records from high northern latitudes of the Atlantic remain rare, leaving gaps in our understanding of their true distribution and ecological role.

The North Atlantic margin of Iceland and the adjacent Reykjanes Ridge represent a dynamic and complex deep-sea environment, shaped by volcanic topography, sediment plains, and a mosaic of hard substrates such as basalt outcrops and glacial dropstones (Astthorsson et al. 2007). Strong mixing of cold North Atlantic Deep Water and Arctic-derived currents creates stable, oxygen-rich bathyal conditions that support a diverse assemblage of benthic fauna, including deep-water corals. This diversity occurs despite the region's sub-Arctic latitude (Astthorsson et al. 2007; Read 2000; Hansen and Østerhus 2000; Burgos et al. 2020).

Within this region, the BIOICE (Benthic Invertebrates of Icelandic Waters) programme conducted a comprehensive


national inventory of benthic diversity between 1991 and 2004, with sampling depths from 20 to 3000 m (Steingrímsson et al. 2020; Guðmundsson 2023). BIOICE was coordinated primarily by the Marine Research Institute and the Icelandic Institute of Natural History and has produced extensive collections of deep-sea invertebrates from around Iceland.

Here, we report the first record of *Crypthelia medioatlantica* Zibrowius & Cairns, 1992 from west Iceland, based on BIOICE material (Fig. 1). Previously, the species was known only from a few localities on the Mid-Atlantic Ridge between ~23° and 37° N (Zibrowius and Cairns 1992). This new record at ~65°N represents the northernmost known occurrence of the species, highlights hidden stylasterid diversity in the North Atlantic, and demonstrates the value of re-examining historical deep-sea collections for refining stylasterid biogeography.

Materials and methods

The specimens were collected in 1996 using a detritus sledge ("Sneli") west of Iceland during the BIOICE programme and are deposited in the collections of the Nature Research Institute of Iceland (NRII). The sampling site (station 2918) is

Fig. 2 Fragments of *Crypthelia medioatlantica* Zibrowius & Cairns, 1992 (NRII cat. no. 43546); **a** branch segment with cyclo systems, most with broken lids; **b** male cyclo system with ampullae; **c** magnified view of cyclo system with lid and male ampullae



located in the Denmark Strait (Fig. 1), close to a major route of southward-flowing deep-water currents from the Nordic Seas (Jochumsen et al. 2016, 2017), which provide cold, dense waters typical of stylasterid habitats. At this station, the detritus sledge sampled a depth of 1539 m on 26 August 1996; associated environmental data (bottom temperature 3.2 °C, salinity 34.9‰) were recorded during sampling. The material consists of six small, dry, dead branch fragments lacking soft tissue. The fragments show subfossil preservation (surface patination, localized smoothing) (Fig. 2).

Morphological examination

Morphological features were examined under a stereomicroscope, focusing on cyclo system size, coenosteal texture, and ampullar structure. Measurements of branch diameter,

cyclo system diameter, and dactylo pore counts were taken using digital calipers from all well-preserved cyclo systems on the six fragments (see Material examined and Diagnosis for sample sizes). Scanning electron microscopy (SEM) was performed at the Naturalis Biodiversity Center (Leiden, The Netherlands) using a JEOL JSM-IT510 SEM operated at 10 kV. Terminology follows Cairns (1983) for the description of stylasterid corals.

Results

Systematic account

Hydrozoa Owen, 1843

Anthoathecata Cornelius, 1992

Stylasteridae Gray, 1847

Crypthelia Milne Edwards & Haime, 1849

Crypthelia medioatlantica Zibrowius & Cairns, 1992

Figures 2 and 3

Crypthelia medioatlantica Zibrowius & Cairns, 1992: 110 (table), 112–114, figs. 36J–M, 38A–H.

Material examined BIOICE station 2918, west of Iceland, 65.21583° N, 29.24867° W, 1539 m depth, detritus sledge (“Snelli”), 26 August 1996, six dry, subfossil fragments, Nature Research Institute of Iceland (NRII), cat. no. 43546.

Taphonomic assessment Macroscopic inspection and SEM indicate subfossil preservation (surface patination; localized smoothing of the coenosteum), but diagnostic micromorphology is intact, allowing confident identification to *C. medioatlantica*.

Diagnosis Subfossil fragments of branches, delicate and small; complete colonies unknown but likely uniplanar. Largest fragment ~8 mm long bearing four cyclosystems; subterminal branches slender, 0.45 mm in diameter (Fig. 2a). Coenosteum exclusively linear-imbricate, with slightly convex strips 70–85 µm wide (Fig. 3a–c), densely covered by narrow platelets 4–14 µm wide; nematopores absent. Cyclosystems circular, 1.2–1.4 mm in diameter, each with 14–19 dactylopores (mean 16.1, mode 15). Upper gastropore chamber ~0.60 mm in diameter, with ring constriction ~0.35 mm; lower chamber ~0.65 mm in diameter and ~0.10 mm deep. Cyclosystem lid digitiform, inclined upwards, sometimes absent, covering 0–20% of the cyclosystem; ~0.30 mm wide. Dactylotomes and pseudosepta ~0.10 mm wide; pseudosepta distinctly concave (Fig. 3d). Male ampullae discrete, hemispherical, ~0.6 mm in diameter, 1–4 arranged around proximal cyclosystem wall (type C1 *sensu* Cairns 2015: Table 9) (Fig. 2b, c); surface covered with a reticulate system of carinae (~60 µm high), extending over the lid. Each ampulla bears an apical depression (~0.18 mm wide) with a central raised papilla (~0.10 mm diameter) and an apical efferent pore 40 µm in diameter. Measurements are based on all well-preserved cyclosystems on the six fragments ($n=9$ cyclosystems).

Remarks *Crypthelia medioatlantica* was previously known from only three records included in its original description, all from the Mid-Atlantic Ridge between 23° 31' N and 36° 50' N (ca. 23–37° N; the latter off the Azores), at depths ranging from 861 to 2644 m (Zibrowius and Cairns 1992) (Fig. 1). This places it among the deepest stylasterid species known to date, comparable to *Crypthelia affinis* Moseley, 1879, which was collected at 2790 m (Moseley 1879). The present record from west of Iceland represents the northernmost known occurrence of *C. medioatlantica* and extends its known latitudinal range by approximately 28° into sub-Arctic waters. The Icelandic record indicates that the species has occurred in sub-Arctic deep-sea environments, but because the material consists solely of dead skeletal fragments, the current status of local populations and the timing of their occurrence remain uncertain.

Female ampullae are unknown but are inferred to be of B-type *sensu* Cairns (2015: Table 9), based on known reproductive dimorphism in congeners.

The subfossil state indicates that the fragments may pre-date collection; establishing their age would require direct dating (U–Th or radiocarbon), so this record documents past presence rather than demonstrating a living population at the time of sampling.

Discussion

Biogeographical and ecological significance

Stylasterid corals are frequently under-represented in benthic biodiversity surveys because their small and fragile colonies, including those of the genus *Crypthelia*, are easily damaged or lost during trawl-based sampling. In the North Atlantic, only a few stylasterid species have confirmed high-latitude records: *Stylaster norvegicus* (Gunnerus, 1768), *Pliobothrus symmetricus* Pourtalès, 1868, *Stylaster gemmascens* (Esper, 1794), and *Stylaster erubescens* Pourtalès, 1868. The present record of *C. medioatlantica* from Icelandic waters adds a fifth species to this group, as revealed by historical BIOICE material, and significantly extends the known northern range of the species.

The new Icelandic locality, together with previous records from the Mid-Atlantic Ridge, suggests a broader ecological tolerance for *C. medioatlantica* and raises important questions about population connectivity and evolutionary history. Like many stylasterids, *Crypthelia* is inferred to be a brooder (Cairns 1986, 2011), implying limited larval dispersal and rare colonization events. In the absence of live material or genetic data, however, we cannot distinguish whether the Icelandic record reflects a long-lived local population that persists at depth, a now-extinct relict population, or a rare long-distance dispersal event from more southerly source populations; the available data are too limited to favor any of these scenarios.

The geographic position of BIOICE station 2918 in the Denmark Strait region suggests a plausible dispersal corridor from the Mid-Atlantic Ridge into sub-Arctic bathyal habitats. Southward-flowing overflow waters and deep boundary currents from the Nordic Seas could facilitate rare long-distance transport along ridge and slope hard substrates, consistent with the species' deep distribution (Jochumsen et al. 2016, 2017). The overlap between the species' bathymetric range (861–2644 m) and these overflow/boundary-current pathways in the Denmark Strait makes deep-current mediated dispersal plausible, potentially aided by ridge and slope “stepping-stone” habitats (Jochumsen et al. 2017; Hansen and Østerhus 2000; Etter




Fig. 3 *Crypthelia medioatlantica* Zibrowius & Cairns, 1992 (NRII cat. no. 43546); **a–c** coenosteal texture; **d** concave pseudosepta; **e, f** stereo pair of coenosteal texture; **g, h**, stereo pair of cyclosystem partially surrounded by male ampullae with visible efferent pores; **i, j**

cyclosystem partially surrounded by male ampullae (lid broken in **i**); **k, l** longitudinal section of cyclosystem showing gastropore chambers (stereo pair) (scale bars: **a–f**=0.1 mm; **g–l**=0.5 mm)

and Bower 2015; Breusing et al. 2016), and mesoscale or topographically steered circulation events (Spall and Price 1998; Koszalka et al. 2017).

Given its depth range, *C. medioatlantica* is likely adapted to environments that remained relatively thermally stable during Pleistocene glacial–interglacial cycles (Rex et al. 2006; Thistle 2003). Such stability may have facilitated

the persistence of cold-adapted stylasterids while glaciation reshaped shallow-water habitats, but this hypothesis remains to be tested. However, Schuchert (2000) reported a sharp decline in hydrozoan diversity below 200 m in Icelandic waters during the BIOICE survey, with no stylasterids documented at all, emphasizing that this depth zone remains severely under-sampled for stylasterids.

Conservation implications

Stylasterid corals are notable for their fragility, slow growth, and limited dispersal capacity (Zibrowius and Cairns 1992; Cairns 1992, 2011). Stylasterids are listed under CITES Appendix II and are recognized as Vulnerable Marine Ecosystem (VME) indicator taxa (Burgos et al. 2020). The Icelandic record of *C. medioatlantica*, although based on a few subfossil fragments, contributes to the baseline mapping of stylasterid diversity in sub-Arctic North Atlantic waters and highlights the importance of continued deep-sea biodiversity assessments in under-explored regions such as Iceland's deep benthic habitats. Resolving remnant vs. extant status will require targeted modern sampling (ROV/post-trawl fine sorting) and, where feasible, direct dating of stylasterid skeletons (U–Th or radiocarbon) to constrain residence time. Stylasterids are easily overlooked in bulk trawl samples due to their small size, and the absence of soft tissue in dead fragments further complicates identification. Although the negative impacts of trawling on corals are increasingly recognized, conservation measures remain species- and region-specific. Records of deep-water corals, including stylasterids, have contributed to the scientific basis for recent Icelandic fisheries closures, such as the designation of the Coral Conservation Area at Papagrúnn slope (Stjórnartíðindi 2023). To improve detection and documentation of cryptic or rarely encountered taxa, targeted sampling strategies, such as fine-scale post-trawl sorting, detailed specimen imaging, and early involvement of taxonomic specialists, are crucial.

In addition to trawling, emerging anthropogenic threats, such as deep-sea mining, oil and gas exploration, and the impacts of climate change, pose new risks to vulnerable coral communities in the North Atlantic (Roberts et al. 2006; Cordes et al. 2016; Levin et al. 2016; Mengerink et al. 2014). Although the specific effects on stylasterid populations are poorly documented, the expansion of industrial activity into deep-sea environments highlights the need for enhanced baseline knowledge and more targeted conservation actions.

The value of historical collections and future directions

The re-examination of archived BIOICE material revealing *C. medioatlantica* highlights the enduring value of historical collections in biodiversity research. As Schuchert (2000) emphasized, the richness of the BIOICE collection continues to reveal previously undocumented hydrozoan diversity, even among well-studied groups. Recent initiatives, such as the 2023 Cold-Water Coral Workshop in Iceland (Sigwart et al. 2025), have further promoted the re-examination of museum

holdings and fostered international taxonomic collaboration. Expanding such efforts will be vital for improving future deep-sea assessments and refining our understanding of coral diversity and distribution in the region.

At the same time, this study illustrates the limitations of relying solely on preserved skeletal material, which in this case lacks soft tissue and therefore precludes molecular analyses that could resolve population connectivity, phylogeographic structure, and evolutionary history. Fresh collections of *C. medioatlantica*, ideally including material from Iceland and from intermediate localities along the Mid-Atlantic Ridge, are needed to test whether the Icelandic record represents an isolated occurrence, a now-extinct relict population, or part of a broader sub-Arctic population system. Complementary life-history research, including studies on age, growth, and reproductive strategies, will also be critical for assessing the vulnerability and resilience of *C. medioatlantica* to pressures such as ocean acidification. Targeted integrative surveys that combine detailed morphology, molecular data, and environmental information at sites like BIOICE station 2918 will therefore be key to resolving the biogeographic context of *C. medioatlantica* and informing conservation planning for stylasterid corals in sub-Arctic regions.

Acknowledgements We thank the crew of the research vessel, the BIOICE programme, and the Nature Research Institute of Iceland (NRII) for providing access to the collection. We are grateful to S. Brix, S. Korfhage, and J. Sigwart (Senckenberg) for organizing the cold-water coral workshop at the Nature Center Sandgerði, Iceland, in September 2023, held as part of the SOSA, IceAGE, and iAtlantic programmes, and for supporting the first author's participation. We also thank K. Jeskulke and A. Fischer (DZMB, Hamburg) for their technical support and data management during the workshop. We thank H. Zibrowius for early discussions on the manuscript. N. Bax acknowledges the 56th European Marine Biology Symposium for the plenary funding that enabled her attendance at the workshop. S. Brix and two anonymous reviewers are thanked for constructive comments that helped improve the manuscript.

Declarations

Conflict of interest The authors declare no competing interests.

Data availability All data supporting the findings of this study are included in the article. The examined material (NRII catalogue no. 43546) and associated metadata are deposited in the Nature Research Institute of Iceland.

Author contribution KS-N conceptualized the study, conducted observations, curated samples, performed imaging, and wrote the first draft. NB conceptualized the study, conducted observations, and contributed to writing and reviewing the manuscript. GG curated samples and data and contributed to writing and reviewing. IS contributed to writing and reviewing. SDC provided advisory support and contributed to writing and reviewing. All authors read and approved the final version of the manuscript.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing,

adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Asthorsson OS, Gislason A, Jonsson S (2007) Climate variability and the Icelandic marine ecosystem. *Deep-Sea Res II* 54:2456–2477. <https://doi.org/10.1016/j.dsr.2007.07.030>
- Breusing C, Biastoch A, Drews A, Metaxas A, Jollivet D, Vrijenhoek RC, Bayer T, Melzner F, Schilhabel MB, Rosenstiel P, Reusch TBH (2016) Biophysical and population genetic models predict the presence of “phantom” stepping stones connecting Mid-Atlantic Ridge vent ecosystems. *Curr Biol* 26:2257–2267. <https://doi.org/10.1016/j.cub.2016.06.062>
- Burgos JM, Buhl-Mortensen L, Buhl-Mortensen P, Ólafsdóttir SH, Steingrund P, Ragnarsson SÁ, Skagseth Ø (2020) Predicting the distribution of indicator taxa of Vulnerable Marine Ecosystems in the Arctic and sub-Arctic waters of the Nordic Seas. *Front Mar Sci* 7:131. <https://doi.org/10.3389/fmars.2020.00131>
- Cairns SD (1983) A generic revision of the Stylasterina (Coelenterata: Hydrozoa). Part I. Description of the genera. *Bull Mar Sci* 33(2):427–508
- Cairns SD (1986) A revision of the Northwest Atlantic stylasterids (Cnidaria: Hydrozoa: Stylasteridae). *Smithson Contrib Zool* 418:1–131
- Cairns SD (1992) Worldwide distribution of the Stylasteridae (Cnidaria: Hydrozoa). *Sci Mar* 56(2–3):125–130
- Cairns SD (2011) Global diversity of the Stylasteridae (Cnidaria: Hydrozoa: Athecata). *PLoS ONE* 6(7):e21670. <https://doi.org/10.1371/journal.pone.0021670>
- Cairns SD (2015) Stylasteridae (Cnidaria: Hydrozoa: Anthoathecata) of the New Caledonian Region. *Mém Mus Natl Hist Nat* 207:1–361
- Cairns SD (2024) Stylasteridae (Cnidaria: Hydrozoa: Anthoathecata) from the insular southwest Indian Ocean. *Mém Mus Natl Hist Nat* 218:223–329
- Cordes EE, Jones DOB, Schlacher TA, Amon DJ, Bernardino AF, Brooke S, Carney R, DeLeo DM, Dunlop KM, Escobar-Briones EG, Gates AR, Génio L, Gobin J, Henry L-A, Herrera S, Hoyt S, Joye M, Kark S, Mestre NC, Metaxas A, Pfeifer S, Sink K, Sweetman AK, Witte U (2016) Environmental impacts of the deep-water oil and gas industry: a review to guide management strategies. *Front Environ Sci* 4:58. <https://doi.org/10.3389/fenvs.2016.00058>
- Etter RJ, Bower AS (2015) Dispersal and population connectivity in the deep North Atlantic estimated from physical transport processes. *Deep-Sea Res I* 104:159–172. <https://doi.org/10.1016/j.dsr.2015.06.009>
- Guðmundsson G (2023) A taxonomic baseline to monitor retreating Arctic biota: the marine invertebrate collection of the Icelandic Institute of Natural History. *Collections* 19(3):353–365. <https://doi.org/10.1177/15501906221147358>
- Hansen B, Østerhus S (2000) North Atlantic-Nordic Seas exchanges. *Prog Oceanogr* 45:109–208. [https://doi.org/10.1016/S0079-6611\(99\)00052-X](https://doi.org/10.1016/S0079-6611(99)00052-X)
- Jochumsen K, Schnurr SM, Quadfasel D (2016) Bottom temperature and salinity distribution and its variability around Iceland. *Deep Sea Res Part I Oceanogr Res Pap* 111:79–90. <https://doi.org/10.1016/j.dsr.2016.02.009>
- Jochumsen K, Moritz M, Nunes N, Quadfasel D, Larsen KMH, Hansen B, Valdimarsson H, Jonsson S (2017) Revised transport estimates of the Denmark Strait overflow in the Irminger Basin. *Ocean Model* 112:90–98. <https://doi.org/10.1016/j.ocemod.2017.03.001>
- Koszalka IM, Haine TWN, Magaldi MG (2017) Mesoscale mixing of the Denmark Strait Overflow in the Irminger Basin. *Ocean Model* 112:90–98. <https://doi.org/10.1016/j.ocemod.2017.03.001>
- Levin LA, Mengerink K, Gjerde KM, Rowden AA, van Dover CL, Clark MR, Ramirez-Llodra E, Currie B, Smith CR, Sato KN, Gallo N, Sweetman AK, Lily H, Armstrong CW, Bridger J (2016) Defining “serious harm” to the marine environment in the context of deep-seabed mining. *Mar Policy* 74:245–259. <https://doi.org/10.1016/j.marpol.2016.09.032>
- Mengerink KJ, Van Dover CL, Ardron J, Baker M, Clark MR, Escobar-Briones E, Gjerde KM, Levin LA, Ramirez-Llodra E, Lara-Lopez A, Squires D et al (2014) A call for deep-ocean stewardship. *Science* 344(6185):696–698. <https://doi.org/10.1126/science.1251458>
- Moseley HN (1879) On the structure of the Stylasteridae, a family of the hydroid stony corals. *Philos Trans R Soc Lond B Biol Sci* 169:425–503
- Read JF (2000) Water masses and circulation in the region of the Iceland-Faroe Front. *ICES J Mar Sci* 57:297–306
- Rex MA, Etter RJ, Morris JS, Crouse J, McClain CR, Johnson NA, Stuart CT, Thies R, Avery R (2006) Global bathymetric patterns of species richness in the deep-sea benthos. *Deep Sea Res Part I Oceanogr Res Pap* 53(5):693–705
- Roberts JM, Wheeler AJ, Freiwald A (2006) Reefs of the deep: the biology and geology of cold-water coral ecosystems. *Science* 312(5773):543–547. <https://doi.org/10.1126/science.1119861>
- Schuchert P (2000) Hydrozoa (Cnidaria) of Iceland collected by the BIOICE programme. *Sarsia* 85:411–438. <https://doi.org/10.1080/00364827.2000.10414592>
- Schuchert P, Choong H, Galea H, Hoeksema B, Lindsay D, Manko M, Pica D (2025) World Hydrozoa Database. *Cryptothelia* Milne Edwards & Haime, 1849. <https://www.marinespecies.org/hydrozoa/aphia.php?p=taxdetails&id=117241>. Accessed via World Register of Marine Species on 23 Nov 2025
- Sigwart JD, Allcock AL, Arantes RCM, Barnhill KA, Bax N, Beneti JS, Brix S, Guðmundsson G, McFadden CS, Korfhage SA, Linardich C, Morrissey D, de Oliveira Nascimento P, de Moura Neves B, Ólafsdóttir SH, Ragnarsson S, Samimi-Namin K, Sampaio Í, de Wilt ME (2025) The first IUCN Red List of cold-water corals highlights global declines. *Mar Biodivers* 55:51. <https://doi.org/10.1007/s12526-025-01533-0>
- Spall MA, Price JF (1998) Mesoscale variability in Denmark Strait: the PV outflow hypothesis. *J. Phys Oceanogr* 28(8):1598–1623. [https://doi.org/10.1175/1520-0485\(1998\)028%3c1598:MVIDST%3e2.0.CO;2](https://doi.org/10.1175/1520-0485(1998)028%3c1598:MVIDST%3e2.0.CO;2)
- Steingrímsson SA, Guðmundsson G, Helgason GV (2020) The BIOICE station and sample list: revised compilation, March 2020. Zenodo. <https://doi.org/10.5281/zenodo.3728257>
- Stjórnartíðindi (2023) <https://www.stjornartidindi.is/Advert.aspx?RecordID=df88f6db-5dc5-4e03-80c0-7ca095b16b20>. Accessed 15 Sept 2023
- Thistle D (2003) The deep-sea floor: an overview. In: Tyler PA (ed) *Ecosystems of the deep oceans*. Elsevier, Amsterdam, pp 5–38
- Zibrowius H, Cairns SD (1992) Revision of the northeast Atlantic and Mediterranean Stylasteridae (Cnidaria: Hydrozoa). *Mém Mus Natl Hist Nat A Zool* 153:1–136