



## Naturalis Repository

# The distribution of the crested and marbled newt species (Amphibia: Salamandridae: Triturus) – an addition to the New Atlas of Amphibians and Reptiles of Europe

Ben Wielstra, Neftalí Sillero, Judit Vörös, and Jan W. Arntzen

Downloaded from:

<http://dx.doi.org/10.1163/15685381-00002960>

### Article 25fa Dutch Copyright Act (DCA) - End User Rights

This publication is distributed under the terms of Article 25fa of the Dutch Copyright Act (Auteurswet) with consent from the author. Dutch law entitles the maker of a short scientific work funded either wholly or partially by Dutch public funds to make that work publicly available following a reasonable period after the work was first published, provided that reference is made to the source of the first publication of the work.

This publication is distributed under the Naturalis Biodiversity Center 'Taverne implementation' programme. In this programme, research output of Naturalis researchers and collection managers that complies with the legal requirements of Article 25fa of the Dutch Copyright Act is distributed online and free of barriers in the Naturalis institutional repository. Research output is distributed six months after its first online publication in the original published version and with proper attribution to the source of the original publication.

You are permitted to download and use the publication for personal purposes. All rights remain with the author(s) and copyrights owner(s) of this work. Any use of the publication other than authorized under this license or copyright law is prohibited.

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the department of Collection Information know, stating your reasons. In case of a legitimate complaint, Collection Information will make the material inaccessible. Please contact us through email: [collectie.informatie@naturalis.nl](mailto:collectie.informatie@naturalis.nl). We will contact you as soon as possible.

# The distribution of the crested and marbled newt species (Amphibia: Salamandridae: *Triturus*) – an addition to the New Atlas of Amphibians and Reptiles of Europe

Ben Wielstra<sup>1,2,\*</sup>, Neftalí Sillero<sup>3</sup>, Judit Vörös<sup>4</sup>, Jan W. Arntzen<sup>2</sup>

**Abstract.** In the recently published New Atlas of Amphibians and Reptiles of Europe (Sillero et al., 2014a), the distribution of the newt genus *Triturus* was not resolved at the level of the species. The main reason for this was the lack of high quality distribution data from in and around the parapatric contact zones between species, where interspecific hybridization occurs. We are working extensively on *Triturus* and the (particularly genetic) data we have accumulated allow us to map the individual *Triturus* species at the appropriate scale. We here provide a database composed of distribution data for the individual species, at generally high resolution, particularly from in and around contact zones. Based on this database we produce maps at the 50 × 50 km UTM grid resolution as used in the new atlas and highlight those grid cells in which more than one *Triturus* species occurs.

*Keywords:* contact zone, herpetofauna, hybrid zone, *Triturus ivanbureschi*.

Recently, the New Atlas of Amphibians and Reptiles of Europe was published (Sillero et al., 2014a). We refer to this publication as ‘the new atlas’ throughout this paper. One of the aims of the new atlas was to identify gaps in the currently available knowledge on species distributions. Such knowledge gaps were particularly prevalent where, in the light of taxonomic progress, the recognition of new species has resulted in poorly understood range boundaries. One of the genera highlighted as troublesome to map was *Triturus*. Only the two main groups encompassing this genus, namely the marbled and crested newts, were represented in the new atlas, rather than the individual species that constitute these two groups (Supplemen-

tary Atlas S1 Distribution Amphibians in Sillero et al., 2014a). For the two species of marbled newts, species identification of larvae for numerous records was uncertain, especially for Portugal (footnote 4 to table 3 in Sillero et al., 2014a). For the six species of crested newts (referred to as a ‘complex’, although in our opinion the term ‘superspecies’ as used by Wallis and Arntzen, 1989 is more appropriate), species were not unambiguously distinguished in the available databases (footnote 20 to table 3 in Sillero et al., 2014a). We here supplement the new atlas by mapping the different European *Triturus* species.

The taxonomy of *Triturus* has been regularly revised (Wolterstorff, 1923; Bucci-Innocenti et al., 1983; Garcia-Paris et al., 1993; Arntzen et al., 2007; Espregueira Themudo and Arntzen, 2007a; Arntzen and Wielstra, 2010; Wielstra et al., 2013c). As a consequence, the number of recognized species and, indeed, parapatric contact zones has increased. Although the geographical distribution of *Triturus* species is well understood in a broad sense (Arntzen, 2003; Herrero et al., 2003a, 2003b), documentation of the contact zones between species constitutes an ongoing effort (Arntzen and Espregueira The-

1 - Department of Animal and Plant Sciences, University of Sheffield, S10 2TN Sheffield, UK

2 - Naturalis Biodiversity Center, P.O. Box 9517, 2300 RA Leiden, The Netherlands

3 - Centro de Investigação em Ciências Geo-Espaciais, Observatório Astronómico Prof. Manuel de Barros, Alameda do Monte da Virgem, 4430-146 Vila Nova de Gaia, Portugal

4 - Department of Zoology, Hungarian Natural History Museum, 1088 Budapest, Hungary

\*Corresponding author;

e-mail: b.wielstra@sheffield.ac.uk


mudo, 2008; Wielstra et al., 2013b). Where species meet, hybridization occurs to a smaller or greater extent (Arntzen, 2003; Arntzen et al., in press). This means that *Triturus* populations can in general be assigned to species unambiguously, except those that are positioned in or near to contact zones. Considering our interest in interspecific hybridization, our research effort has particularly focussed on documenting the position of *Triturus* hybrid zones (Espregueira Themudo and Arntzen, 2007b; Wielstra and Arntzen, 2012; Arntzen et al., in press).

Although morphological criteria provide a reasonable estimation of species identity for *Triturus*, the diagnostic value of these criteria breaks down near hybrid zones (Arntzen and Wallis, 1999; Arntzen, 2003; Arntzen et al., in press). Furthermore, for the group of species previously treated as *T. karelinii*, no species specific morphological character states are as yet known (Ivanović et al., 2013). Genetic data, on the other hand, provide an objective measure of species identity. For species assignment and hybrid zone analysis we have previously used mitochondrial DNA restriction fragment length polymorphism (Wallis and Arntzen, 1989; Arntzen and Wallis, 1991), allozymes (Arntzen and Wallis, 1991; Espregueira Themudo and Arntzen, 2007a; Vörös and Arntzen, 2010; Arntzen et al., in press), microsatellites (Jehle et al., 2005; Arntzen et al., 2009, 2010) and sequence data for mitochondrial DNA (Wielstra et al., 2010, 2013b; Wielstra and Arntzen, 2011) and up to five nuclear DNA markers (Espregueira Themudo et al., 2009, 2012; Wielstra et al., 2013a). We have recently developed an Ion Torrent next-generation sequencing protocol targeting 52 nuclear DNA markers (Wielstra et al., 2014; Wielstra and Arntzen, unpublished) and in the context of our ongoing hybrid zone research we are genotyping a large number of *Triturus* newts held in the collection of Naturalis Biodiversity Center (e.g. Wielstra and Arntzen, 2014a).

With the above marker systems we have greatly increased our understanding of the *Tritu-*

*rus* distribution in and around the contact zones. Localities from the Naturalis Biodiversity Center collection form the core of our database of *Triturus* distribution data (579 data points). Additional distribution data were taken from the literature (e.g. Tarkhnishvili and Gokheshvili, 1999; Tzankov and Stoyanov, 2008; Litvinchuk and Borkin, 2009; Canestrelli et al., 2012; Mikulíček et al., 2012; 1469 data points), from an atlas (Loureiro et al., 2010; 375 data points), from natural history collections (71 data points) and from the national database for Hungary (not yet included in the new atlas upon its publication; 60 data points) or were communicated to us by colleagues (88 data points). Furthermore, digitized atlas, survey and museum data were taken from the Global Biodiversity Information Facility (accessible via [www.gbif.org](http://www.gbif.org); 2526 data points). A map showing the *Triturus* distribution data is provided in fig. 1. The underlying database (5168 data points in total), including a complete list of references, is available online as Supplementary Material S1. We updated an earlier version of this database (Wielstra et al., 2012, 2013b), with the most notable changes being that we incorporated field data up to and including 2014, digitized museum data for Hungary and recently published distribution data for Romania (Cogălniceanu et al., 2013).

We converted the data following Sillero et al. (2014a) to map the individual *Triturus* species at a 50 × 50 km UTM grid resolution. Not considered for these maps were distribution data falling outside of the area covered by the new atlas or from around contact zones for which either no genetic data could be consulted or for which genetic admixture was detected (but for the sake of completeness these distribution data are included in fig. 1 and Supplementary Material S1). Because our database particularly focusses on the contact zones between the *Triturus* species, it is less complete away from contact zones, particularly in the core of the range of *T. cristatus* (although our database does manage to fill a considerable




**Figure 1.** The *Triturus* distribution database presented in this paper, plotted on a map. Raw data and references are provided as Supplementary Material S1. Data points are colour coded according to species identity and we distinguish one more candidate species. Genetically admixed populations or populations for which genetic data is required to establish identity are noted as such. Ranges are mostly parapatric but those of the marbled newt *T. marmoratus* and the crested newt *T. cristatus* overlap in western France. This figure is published in colour in the online version.

number of additional grid cells in Central Europe and on the Balkan Peninsula compared to the new atlas). Fortunately, in the new atlas a large number of additional  $50 \times 50$  km UTM grid cells containing *Triturus* records positioned away from the contact zones were presented. Guided by the database in the Supplementary Material S1, these grid cells were allocated to species when this could be done unambiguously, i.e. when they were positioned away from contact zones, and incorporated on the species maps. An overview of the maps is provided in fig. 2 and detailed maps can be found as online Supplementary fig. S1. Shape files are available under Supplementary Material S1. The new distribution data for the individual *Triturus* species can now be integrated in ‘NA2RE’, the web-based spatial data infrastructure underpinning the new atlas (Sillero et al., 2014b; <http://na2re.ismai.pt>).

The new atlas used the name *T. arntzeni* for the Balkan representative of the group of species traditionally referred to as *T. karelinii* (hereafter *T. karelinii sensu lato*). We have recently shown that the name *T. arntzeni* is not

valid, as newts from the type locality are genetically admixed, containing genetic material of (mostly) *T. macedonicus* and *T. karelinii sensu lato* (Wielstra et al., 2013c; Wielstra and Arntzen, 2014a) and we have introduced the name *T. ivanbureschi* instead (Wielstra et al., 2013c). A distinct genetic group currently subsumed in *T. ivanbureschi*, distributed in Asiatic Turkey and hence outside of the region covered by the new atlas, may represent an as yet undescribed crested newt species (Wielstra et al., 2013a, 2013c).

Mapping of interspecific *Triturus* hybrids was beyond the purpose of the new atlas, but we provide some general insight on the distribution of hybrids. F1 hybrids of the marbled newt *T. marmoratus* and the crested newt *T. cristatus* can be expected to occur throughout the area where the two parent species are in sympatry in western France. These hybrids represent an ‘evolutionary dead end’ with the frequency of backcrossing being negligible (Wolterstorff, 1904; Arntzen and Wallis, 1991; Arntzen et al., 2009). Backcrossing is possible with hybrids between marbled newt species or hy-



**Figure 2.** Distribution maps for European *Triturus* newts. Maps are in the format of the New Atlas of Amphibians and Reptiles of Europe, at a  $50 \times 50$  km UTM grid resolution and restricted to Europe. The composite map in the centre shows the number of *Triturus* species per grid cell (red grid cells reflecting the presence of a single and blue ones the presence of two species) and the surrounding maps (arranged in line with geography) show the individual species. This figure is published in colour in the online version. Higher resolution versions of the individual maps can be found online in Supplementary fig. S1.

brids between crested newt species, as is reflected by asymmetric mtDNA introgression (Espregueira Themudo and Arntzen, 2007b; Espregueira Themudo et al., 2012; Wielstra and Arntzen, 2012, 2014b). However, the narrowness of the hybrid zones suggests the prevalence of reproductive isolating mechanisms, limiting gene flow and allowing species to maintain their genetic integrity (Arntzen, 2003; Arntzen et al., in press). Grid cells where multiple *Triturus*

species co-occur and where interspecific hybridization could be expected are presented in fig. 2 and Supplementary fig. S1.

We here improved on the New Atlas of Amphibians and Reptiles of Europe (Sillero et al., 2014a) by providing distribution data for the individual species comprising the marbled and crested newts, particularly from in and around the parapatric contact zones between species (fig. 2, fig. S1). As stated in the new atlas,

“a precise knowledge of the spatial distribution of taxa is essential for decision-making processes in land management and biodiversity conservation, both for present and under future global change scenarios”. Additionally, *Triturus* provides a suitable system for hybrid zone study and for this aim a clear distribution picture for the different *Triturus* species is essential. The new atlas highlighted additional taxa for which detailed studies of the contact zones between constituent species, such as provided here for *Triturus*, are warranted. We hope this paper would stimulate researchers working on such taxa to publish up-to-date distribution maps.

**Acknowledgements.** We are grateful to all our colleagues who contributed *Triturus* distribution data over the years. Mathieu Denoël provided helpful comments on an earlier version of this manuscript. B.W. is a Newton International Fellow. J.V. is supported by the Hungarian Scientific Research Fund OTKA K84071.

## References

- Arntzen, J.W., Wallis, G.P. (1991): Restricted gene flow in a moving hybrid zone of the newts *Triturus cristatus* and *T. marmoratus* in western France. *Evolution* **45**: 805-826.
- Arntzen, J.W., Wallis, G.P. (1999): Geographic variation and taxonomy of crested newts (*Triturus cristatus* superspecies): morphological and mitochondrial data. *Contrib. Zool.* **68**: 181-203.
- Arntzen, J.W. (2003): *Triturus cristatus* superspecies – Kammolch-Artenkreis (*Triturus cristatus* Laurenti, 1768) – Nördlicher Kammolch, *Triturus carnifex* (Laurenti, 1768) – Italienischer Kammolch, *Triturus dobrogicus* (Kiritzescu, 1903) – Donau-Kammolch, *Triturus karelinii* (Strauch, 1870) – Südlicher Kammolch). In: *Handbuch der Reptilien und Amphibien Europas*. Schwanzlurche IIA, p. 421-514. Grossenbacher, K., Thiesmeier, B., Eds, Aula-Verlag, Wiebelsheim.
- Arntzen, J.W., Espregueira Themudo, G., Wielstra, B. (2007): The phylogeny of crested newts (*Triturus cristatus* superspecies): nuclear and mitochondrial genetic characters suggest a hard polytomy, in line with the paleogeography of the centre of origin. *Contrib. Zool.* **76**: 261-278.
- Arntzen, J.W., Espregueira Themudo, G. (2008): Environmental parameters that determine species geographical range limits as a matter of time and space. *J. Biogeogr.* **35**: 1177-1186.
- Arntzen, J.W., Jehle, R., Bardakci, F., Burke, T., Wallis, G.P. (2009): Asymmetric viability of reciprocal-cross hybrids between crested and marbled newts (*Triturus cristatus* and *T. marmoratus*). *Evolution* **63**: 1191-1202.
- Arntzen, J.W., Burke, T., Jehle, R. (2010): Estimating the propagule size of a cryptogenic crested newt population. *Anim. Conserv.* **13**: 74-81.
- Arntzen, J.W., Wielstra, B. (2010): Where to draw the line? A nuclear genetic perspective on proposed range boundaries of the crested newts *Triturus karelinii* and *T. arntzeni*. *Amphibia-Reptilia* **31**: 311-322.
- Arntzen, J.W., Wielstra, B., Wallis, G.P. (in press): The modality of nine *Triturus* newt hybrid zones, assessed with nuclear, mitochondrial and morphological data. *Biol. J. Linn. Soc.*, DOI:10.1111/bij.12358.
- Bucci-Innocenti, S., Raggiamenti, M., Mancino, G. (1983): Investigations of karyology and hybrids in *Triturus boscai* and *T. vittatus*, with a reinterpretation of the species groups within *Triturus* (Caudata: Salamandridae). *Copeia* **1983**: 662-672.
- Canestrelli, D., Salvi, D., Maura, M., Bologna, M.A., Nascetti, G. (2012): One species, three Pleistocene evolutionary histories: phylogeography of the Italian crested newt, *Triturus carnifex*. *PLoS ONE* **7**: e41754.
- Cogălniceanu, D., Székely, P., Samoilă, C., Ruben, I., Tudor, M., Plăiașu, R., Stănescu, F., Rozyłowicz, L. (2013): Diversity and distribution of amphibians in Romania. *ZooKeys* **296**: 35-57.
- Espregueira Themudo, G., Arntzen, J.W. (2007a): Molecular identification of marbled newts and a justification of species status for *Triturus marmoratus* and *T. pygmaeus*. *Herpetol. J.* **17**: 24-30.
- Espregueira Themudo, G., Arntzen, J.W. (2007b): Newts under siege: range expansion of *Triturus pygmaeus* isolates populations of its sister species. *Divers. Distrib.* **13**: 580-586.
- Espregueira Themudo, G., Wielstra, B., Arntzen, J.W. (2009): Multiple nuclear and mitochondrial genes resolve the branching order of a rapid radiation of crested newts (*Triturus*, Salamandridae). *Mol. Phylogenet. Evol.* **52**: 321-328.
- Espregueira Themudo, G., Nieman, A., Arntzen, J.W. (2012): Is dispersal guided by the environment? A comparison of interspecific gene flow estimates among differentiated regions of a newt hybrid zone. *Mol. Ecol.* **21**: 5324-5335.
- Garcia-Paris, M., Herrero, P., Martin, C., Dorda, J., Esteban, M., Arano, B. (1993): Morphological characterization, cytogenetic analysis, and geographical distribution of the Pygmy marbled newt, *Triturus marmoratus pygmaeus* (Wolterstorff, 1905) (Caudata, Salamandridae). *Bijdr. Dierkd.* **63**: 3-14.
- Herrero, P., Montori, A., Arano, B. (2003a): *Triturus marmoratus* (Latreille, 1800) – Nördlicher Marmormolch. In: *Handbuch der Reptilien und Amphibien Europas*. Schwanzlurche IIA, p. 515-541. Grossenbacher, K., Thiesmeier, B., Eds, Aula-Verlag, Wiebelsheim.
- Herrero, P., Montori, A., Arano, B. (2003b): *Triturus pygmaeus* (Wolterstorff, 1905) – Südlicher Marmormolch. In: *Handbuch der Reptilien und Amphibien Europas*. Schwanzlurche IIA, p. 543-553. Grossenbacher, K., Thiesmeier, B., Eds, Aula-Verlag, Wiebelsheim.

- Ivanović, A., Üzüm, N., Wielstra, B., Olgun, K., Litvinchuk, S.N., Kalezić, M.L., Arntzen, J.W. (2013): Is mitochondrial DNA divergence of Near Eastern crested newts (*Triturus karelinii* group) reflected by differentiation of skull shape? *Zool. Anz.* **252**: 269-277.
- Jehle, R., Wilson, G.A., Arntzen, J.W., Burke, T. (2005): Contemporary gene flow and the spatio-temporal genetic structure of subdivided newt populations (*Triturus cristatus*, *T. marmoratus*). *J. Evol. Biol.* **18**: 619-628.
- Litvinchuk, S.N., Borkin, L.J. (2009): Evolution, Systematics and Distribution of Crested Newts (*Triturus cristatus* Complex) in Russia and Adjacent Countries. Evropejsky Dom Press, St. Petersburg.
- Loureiro, A., Ferrand, N., Carretero, M.A., Paulo, O.S. (2010): Atlas dos Anfíbios e Répteis de Portugal Lisboa, Instituto da Conservação da Natureza e da Biodiversidade.
- Mikulíček, P., Horák, A., Zavadil, V., Kautman, J., Piálek, J. (2012): Hybridization between three crested newt species (*Triturus cristatus* superspecies) in the Czech Republic and Slovakia: Comparison of nuclear markers and mitochondrial DNA. *Folia Zool.* **61**: 202-218.
- Sillero, N., Campos, J., Bonardi, A., Corti, C., Creemers, R., Crochet, P.-A., Crnobrnja Isailović, J., Denoël, M., Ficetola, G.F., Gonçalves, J., Kuzmin, S., Lymberakis, P., de Pous, P., Rodríguez, A., Sindaco, R., Speybroeck, J., Toxopeus, B., Vieites, D.R., Vences, M. (2014a): Updated distribution and biogeography of amphibians and reptiles of Europe. *Amphibia-Reptilia* **35**: 1-31.
- Sillero, N., Oliveira, M.A., Sousa, P., Sousa, F., Gonçalves-Seco, L. (2014b): Distributed database system of the New Atlas of Amphibians and Reptiles in Europe: the NA2RE project. *Amphibia-Reptilia* **35**: 33-39.
- Tarkhishvili, D.N., Gokheshvili, R.K. (1999): The Amphibians of the Caucasus. Pensoft, Sofia.
- Tzankov, N., Stoyanov, A. (2008): *Triturus cristatus* (Laurenti, 1768): a new species for Bulgaria from its southernmost known localities. *Salamandra* **44**: 153-161.
- Vörös, J., Arntzen, J.W. (2010): Weak population structuring in the Danube crested newt, *Triturus dobrogicus*, inferred from allozymes. *Amphibia-Reptilia* **31**: 339-346.
- Wallis, G.P., Arntzen, J.W. (1989): Mitochondrial-DNA variation in the crested newt superspecies: limited cytoplasmic gene flow among species. *Evolution* **43**: 88-104.
- Wielstra, B., Espregueira Themudo, G., Güclü, Ö., Olgun, K., Poyarkov, N.A., Arntzen, J.W. (2010): Cryptic crested newt diversity at the Eurasian transition: the mitochondrial DNA phylogeography of Near Eastern *Triturus* newts. *Mol. Phylogenet. Evol.* **56**: 888-896.
- Wielstra, B., Arntzen, J.W. (2011): Unraveling the rapid radiation of crested newts (*Triturus cristatus* superspecies) using complete mitogenomic sequences. *BMC Evol. Biol.* **11**: 162.
- Wielstra, B., Arntzen, J.W. (2012): Postglacial species displacement in *Triturus* newts deduced from asymmetrically introgressed mitochondrial DNA and ecological niche models. *BMC Evol. Biol.* **12**: 161.
- Wielstra, B., Beukema, W., Arntzen, J.W., Skidmore, A.K., Toxopeus, A.G., Raes, N. (2012): Corresponding mitochondrial DNA and niche divergence for crested newt candidate species. *PLoS ONE* **7**: e46671.
- Wielstra, B., Baird, A.B., Arntzen, J.W. (2013a): A multimer phylogeography of crested newts (*Triturus cristatus* superspecies) reveals cryptic species. *Mol. Phylogenet. Evol.* **67**: 167-175.
- Wielstra, B., Crnobrnja-Isailović, J., Litvinchuk, S.N., Reijnen, B.T., Skidmore, A.K., Sotiropoulos, K., Toxopeus, A.G., Tzankov, N., Vukov, T., Arntzen, J.W. (2013b): Tracing glacial refugia of *Triturus* newts based on mitochondrial DNA phylogeography and species distribution modeling. *Front. Zool.* **10**: 13.
- Wielstra, B., Litvinchuk, S., Naumov, B., Tzankov, N., Arntzen, J.W. (2013c): A revised taxonomy of crested newts in the *Triturus karelinii* group (Amphibia: Caudata: Salamandridae), with the description of a new species. *Zootaxa* **3682**: 441-453.
- Wielstra, B., Arntzen, J.W. (2014a): Kicking *Triturus arntzeni* when it's down: large-scale nuclear genetic data confirm that newts from the type locality are genetically admixed. *Zootaxa* **3802**: 381-388.
- Wielstra, B., Arntzen, J.W. (2014b): Exploring the effect of asymmetric mitochondrial DNA introgression on estimating niche divergence in morphologically cryptic species. *PLoS ONE* **9**: e95504.
- Wielstra, B., Duijm, E., Lagler, P., Lammers, Y., Meilink, W.R.M., Ziermann, J.M., Arntzen, J.W. (2014): Parallel tagged amplicon sequencing of transcriptome-based genetic markers for *Triturus* newts with the Ion Torrent next-generation sequencing platform. *Mol. Ecol. Resour.* **14**: 1080-1089.
- Wolterstorff, W. (1904): *Triton Blasii* de l'Isle, ein Kreuzungsprodukt zwischen *Triton marmoratus* und *Tr. cristatus*. *Zool. Anz.* **28**: 82-86.
- Wolterstorff, W. (1923): Übersicht den Unterarten und Formen des *Triton cristatus* Laur. *Blätter für Aquarien- und Terrarienkunde* **34**: 120-126.

Submitted: May 14, 2014. Final revision received: August 4, 2014. Accepted: August 11, 2014.

Associate Editor: Francesco Ficetola.