



Naturalis Repository

## The effect of area and isolation on insular dwarf proboscideans

Alexandra A. E. van der Geer, Gerrit D. van den Bergh, George A. Lyras, Unggul W. Prasetyo, Rokus Awe Due, Erick Setiyabudi, Hara Drinia

Downloaded from:

<https://doi.org/10.1111/jbi.12743>

### Article 25fa Dutch Copyright Act (DCA) - End User Rights

This publication is distributed under the terms of Article 25fa of the Dutch Copyright Act (Auteurswet) with consent from the author. Dutch law entitles the maker of a short scientific work funded either wholly or partially by Dutch public funds to make that work publicly available following a reasonable period after the work was first published, provided that reference is made to the source of the first publication of the work.

This publication is distributed under the Naturalis Biodiversity Center 'Taverne implementation' programme. In this programme, research output of Naturalis researchers and collection managers that complies with the legal requirements of Article 25fa of the Dutch Copyright Act is distributed online and free of barriers in the Naturalis institutional repository. Research output is distributed six months after its first online publication in the original published version and with proper attribution to the source of the original publication.

You are permitted to download and use the publication for personal purposes. All rights remain with the author(s) and copyrights owner(s) of this work. Any use of the publication other than authorized under this license or copyright law is prohibited.

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the department of Collection Information know, stating your reasons. In case of a legitimate complaint, Collection Information will make the material inaccessible. Please contact us through email: [collectie.informatie@naturalis.nl](mailto:collectie.informatie@naturalis.nl). We will contact you as soon as possible.



ORIGINAL  
ARTICLE



# The effect of area and isolation on insular dwarf proboscideans

Alexandra A. E. van der Geer<sup>1,2\*</sup>, Gerrit D. van den Bergh<sup>1,3</sup>, George A. Lyras<sup>2</sup>, Unggul W. Prasetyo<sup>3,4</sup>, Rokus Awe Due<sup>5†</sup>, Erick Setiyabudi<sup>4</sup> and Hara Drinia<sup>2</sup>

<sup>1</sup>Department of Geology, Netherlands Biodiversity Center Naturalis, Leiden, The Netherlands, <sup>2</sup>National and Kapodistrian University of Athens, Zografou, Greece, <sup>3</sup>Centre for Archaeological Science, School of Earth & Environmental Sciences, University of Wollongong, Wollongong Australia, <sup>4</sup>Geology Museum, Bandung Indonesia, <sup>5</sup>National Research Center for Archaeology, Jakarta, Indonesia

## ABSTRACT

**Aim** We investigated the hypothesis that insular body size of fossil elephants is directly related to isolation and surface area of the focal islands.

**Location** Palaeo-islands worldwide.

**Methods** We assembled data on the geographical characteristics (area and isolation) of islands and body size evolution of palaeo-insular species for 22 insular species of fossil elephants across 17 islands.

**Results** Our results support the generality of the island rule in the sense that all but one of the elephants experienced dwarfism on islands. The smallest islands generally harbour the smallest elephants. We found no support for the hypothesis that body size of elephants declines with island isolation. Body size is weakly and positively correlated with island area for proboscideans as a whole, but more strongly correlated for Stegodontidae when considered separately. Average body size decrease is much higher when competitors are present.

**Main conclusions** Body size in insular elephants is not significantly correlated with the isolation of an island. Surface area, however, is a significant predictor of body size. The correlation is positive but relatively weak; *c.* 23% of the variation is explained by surface area. Body size variation seems most strongly influenced by ecological interactions with competitors, possibly followed by time in isolation. Elephants exhibited far more extreme cases of dwarfism than extant insular mammals, which is consistent with the substantially more extended period of deep geological time that the selective pressures could act on these insular populations.

## Keywords

*Elephas*, fossil record, insularity, island biogeography, island rule, *Palaeoloxodon*, Pleistocene, *Stegodon*

\*Correspondence: Alexandra A. E. van der Geer, Netherlands Biodiversity Center Naturalis, Leiden, The Netherlands.

E-mail: alexandra.vandergeer@naturalis.nl

†The fifth author passed away while the article was in an advanced stage.

## INTRODUCTION

Vertebrates on islands generally follow a body size trend from gigantism in small species to dwarfism in large species, a trend coined ‘the Island Rule’ by Van Valen (1973) and subsequent authors. The many deviations from this pattern have given rise to considerable criticism regarding generality and causality of the rule: see, for example, Meiri *et al.* (2004, 2006, 2008a) for mammalian carnivores (but see Lyras *et al.*, 2010), Meiri (2007) for lizards, Meiri *et al.* (2011) for birds and lizards (though they found weak support for mammals), and Itescu *et al.* (2014) for insular tortoises worldwide. A

large study has investigated the generality of the island rule (Lomolino *et al.*, 2013), which, by analysing both extant and extinct mammals (including very large mammals such as elephants in the dataset), includes the important factor deep geological time under which species evolved in isolation. Context (ecological interactions, isolation, area, climate etc.) appears to be of crucial importance and differences herein can explain differences in observed body size evolution between populations of the same species (Lomolino *et al.*, 2012). The various contextual factors do not contribute equally to the eventual body mass outcome, for example, the presence of ecologically relevant competitors appears to have

a higher impact than ancestral body mass and climatic variability (Sondaar, 1977; Lomolino, 1985; Raia & Meiri, 2006; Palombo, 2007, 2009; Lomolino *et al.*, 2012; van der Geer *et al.*, 2013; *inter alia* Itescu *et al.*, 2014). For example, insular elephants have been shown to have maintained a larger body mass in the presence of deer and hippo, likely because the presence of smaller bodied intraguild competitors prevented elephants from reaching a similar size class (Palombo, 2007).

Taxa such as murids and deer are well represented in both the fossil as well as the extant insular record. However, very large-bodied taxa such as elephants and hippos are extremely limited in the extant record despite being common, not to say typical, elements of island faunas worldwide during the Pleistocene. Elephants provide some of the most spectacular cases of body size evolution in palaeo-insular mammals (Fig. 1). For example, the elephant (*Palaeoloxodon falconeri*) from Spinagallo Cave (Siracuse, Sicily; Middle Pleistocene) dwarfed to just 2% of the size (body mass) of its mainland ancestor (*Palaeoloxodon antiquus*) (Ambrosetti, 1968; Palombo & Ferretti, 2005; Lomolino *et al.*, 2013).

Here, we expand the dataset for insular proboscideans as given in Lomolino *et al.* (2013) and further evaluate insular size trends of these very large insular mammals which are lacking in basically extant insular faunas. The dataset is limited to North America and Eurasia as no fossil proboscideans have been discovered from islands of Africa, South America or Antarctica.



**Figure 1** Reconstruction of four insular dwarf proboscideans with their respective mainland ancestors. Mainland proboscideans: 1, *Palaeoloxodon antiquus*; 2, *Mammuthus columbi*; 3, *Stegodon zdanskyi*. Insular proboscideans: 4, *Palaeoloxodon 'mnaidriensis'*; 5, *Palaeoloxodon falconeri*; 6, *Mammuthus exilis*; 7, *Stegodon aurorae*. Based on skeletons at Museo di Paleontologia, University of Rome, Italy (1), American Museum of Natural History, New York (2), Taylor Made Fossils, U.S. (3), Museo di Paleontologia e Geologia G.G. Gemmellaro, Palermo, Italy (4), Forschungsinstitut und Naturmuseum Senckenberg, Frankfurt, Germany (5), Santa Barbara Museum of Natural History, Santa Barbara, U.S. (6), Taga Town Museum, Honshu, Japan (7). Photos 1–2, 4–7 George Lyras, photo 3 courtesy of TaylorMadeFossils.com, reproduced here with permission.

In the conceptual model as proposed by Lomolino *et al.* (2012), the principal selection force for (very) large mammals is ecological release. Ecological release, however, is more important on smaller islands (where there are fewer competitors/predators), which would indirectly introduce an effect of area on elephant body size. The number of ecologically relevant competitors/predators is also related to isolation, as the chance of successful dispersal is inversely related to isolation. This means that both area and isolation are expected to have an indirect effect on the degree of dwarfism of elephants through ecological release.

The effect of factors associated with thermoregulation (e.g. island latitude, temperature and seasonality) is potentially relevant, but will not be tested here as the variation in climatic variables, especially lowest winter temperature and rainfall, across the focal palaeo-islands and the different geological periods (including glacials, interglacials and spanning the entire Pleistocene) cannot be estimated reliably.

The testable predictions of the ecological model of body size evolution (Lomolino *et al.*, 2012) for palaeo-insular proboscideans are as follows.

1. Proboscideans should exhibit dwarfism on islands, following the general 'Island Rule' prediction that large mammals become smaller.
2. Body size in proboscideans should be positively correlated with island area but negatively correlated with isolation. The rationale here is that both area and isolation are correlated with the presence and number of ecologically relevant competitors and predators: the first as a function of carrying capacity, the second as a function of dispersal chances.
3. The degree of dwarfism should be highest in the absence of ecologically interacting taxa. The presence of large to medium-sized competing herbivores and/or of predators is expected to prevent dwarfism to avoid competition and/or escape predation respectively.

## MATERIALS AND METHODS

### Estimation of body size and relative size index ( $S_i$ )

As is detailed within Table 1, the dataset comprises 22 insular proboscidean species. Because this study includes fossilized individuals, preservation is inconsistent. Therefore, and of necessity, different methods for estimating body size and relative size index were applied. The calculated relative size index ( $S_i$  or size ratio) for each pair of insular proboscidean and its mainland ancestral species was obtained by dividing the estimated body mass of the focal species by that of the ancestral species (=SR in Meiri *et al.*, 2008a). Body size (or mass) of insular fossil species and their mainland ancestors was either estimated using taxon-specific equations for estimating body masses developed by Christiansen (2004) for living elephants, or taken from published sources (see Appendix S1 in Supporting Information for details and references). The alternative method of calculating ratios based on linear measurements and applying the cubic law instead of


**Table 1** Variation in body mass indices ( $S_i$  as a proportion of ancestral body size) of insular proboscideans in alphabetical order of species name. Body mass estimations were based on postcranial elements (humerus, femur, tibia) using the equations of Christiansen (2004) for *Mammuthus*, *Palaeoloxodon* and for *Stegodon* species for which these elements were available. For *Stegodon* species of which only (fragments of) adult mandibles were available, body mass estimates were based on a regression between log-transformed postcranial-based average body mass estimates and average mandibular width measurements of adult specimens (M18 of van den Bergh, 1999; only mandibles were used in which the last molar was in function), with body mass as the 'dependent' variable. This method was developed based on *Stegodon* species for which both variables were available based on averages of more than 1 specimen (see Appendix S1). Indices based on the lower third molar ( $M_3$ ) are calculated as the ratio of the cubed linear dimensions, following Lomolino (2005). Although this method likely overestimates body size of large-sized insular forms, it appears reliable in small-sized phylogenetic dwarfs (for a discussion on teeth and body size reduction in phylogenetic dwarfs, see Lister, 1996 and see Appendix S1). The species indicated with an asterisk is an older chronospecies preceding the smaller form. Sources for  $S_i$ : 1, van den Bergh, 1999; 2, Lomolino *et al.*, 2013; 3, this study, based on M, mandible; P, postcranial; or T, third molar (see Appendix S1 for specimens, method and details); 4, van der Geer *et al.*, 2014. The 'large-sized *Elephas*' from Sulawesi (Tanrunga Formation) is likely *P. namadicus*. No complete long bones or third molars were found, but according to van den Bergh (1999) the size of a molar fragment falls within the size range of *P. namadicus*, hence its body size index is tentatively taken here to be 1.0. This may be an overestimation in case size decrease had just begun. For calculation of area and isolation at sea level 60 m below present, see Appendix S1. Geological age: EP, early Pleistocene; MP, middle Pleistocene; LP, late Pleistocene; H, Holocene.

Palaeo-island	Species	Mainland ancestor	$S_i$	Source	Area (km <sup>2</sup> )/isolation (km)	Geological age
Crete	<i>Mammuthus creticus</i>	<i>M. meridionalis</i>	0.04	2	(4175 or 8350)/90	EP
Santa Rosae	<i>Mammuthus exilis</i>	<i>M. columbi</i>	0.17	2	2200/10	LP
Sardinia	<i>Mammuthus lamarmorai</i>	<i>M. meridionalis</i>	0.08	2	26,343/50	LP
Wrangel	<i>Mammuthus primigenius</i>	<i>M. primigenius</i>	0.47	2	7600/140	H
Sicily	<i>Palaeoloxodon 'mnaidriensis'</i>	<i>P. antiquus</i>	0.17	2	23,000/(1 or 4)	Late MP-early LP
Crete	<i>Palaeoloxodon creutzburgi</i>	<i>P. antiquus</i>	0.38	2	9460/90	LP
Kassos	<i>Palaeoloxodon 'creutzburgi'</i>	<i>P. antiquus</i>	0.30	3 (T)	600/50	LP
Cyprus	<i>Palaeoloxodon cypriotes</i>	<i>P. antiquus</i>	0.02	4	9234/69	LP
Sicily	<i>Palaeoloxodon falconeri</i>	<i>P. antiquus</i>	0.02	2	(2200/200) or (11,600/100)	Early MP
Naxos	<i>Palaeoloxodon lomolinoi</i>	<i>P. antiquus</i>	0.08	4	1226/30	LP
Rhodos	<i>Palaeoloxodon</i> sp.	<i>P. antiquus</i>	0.19	4	1660/15	LP?
Sulawesi	<i>Palaeoloxodon</i> sp. ( 'large-sized <i>Elephas</i> ' )	<i>P. namadicus</i>	c. 1.00	1	174,600/50	MP
Tilos	<i>Palaeoloxodon tiliensis</i>	<i>P. antiquus</i>	0.09	2	116/15	LP
Japan	<i>Stegodon aurorae</i>	<i>S. zdanskyi</i>	0.25	3 (P)	263,237/20	EP
Flores	<i>Stegodon florensis florensis</i> *	<i>S. ganesa</i>	0.51	3 (P)	21,910/(45 or 225)	MP
Flores (Liang Bua)	<i>Stegodon florensis insularis</i>	<i>S. ganesa</i>	0.17	3 (P)	21,910/(45 or 225)	LP
Luzon	<i>Stegodon luzonensis</i>	<i>S. ganesa</i>	0.39	3 (T)	(157,000 or 59,250)/(45 or 65)	LP
Sulawesi	<i>Stegodon sompoensis</i>	<i>S. ganesa</i>	0.32	3 (M)	(174,600 or 24,020)/50	EP
Flores (So'a Basin)	<i>Stegodon sondaari</i>	<i>S. elephantoides</i>	0.15	3 (P)	21,910/(45 or 225)	EP
Sulawesi	<i>Stegodon</i> sp. B	<i>S. ganesa</i>	0.57	3 (T)	174,600/50	MP
Sumba	<i>Stegodon sumbaensis</i>	<i>S. ganesa</i>	0.08	3 (M)	3250/119	MP-LP
Timor	<i>Stegodon timorensis</i>	<i>S. ganesa</i>	0.23	3 (M)	10,560/78	MP
Java (Trinil H.K.)	<i>Stegodon trigonocephalus</i>	<i>S. ganesa</i>	0.65	3 (P)	138,794/(1 or 4)	Late EP

indirectly via body mass has been demonstrated to work equally well (Meiri *et al.*, 2006, 2008a), but we preferred the indirect method enabling inclusion of data from the literature and thereby increasing our sample size. Parameters that were used in this study to estimate body mass were the lengths of appendicular bones (humerus, femur, tibia) that provided the most reliable regression results in the study by Christiansen (2004). For each species, we calculated body masses based on the average lengths for each element (or one or two of these elements, if not all three were available, again in order to increase our  $S_i$  sample size). We then took the average of the mass estimates based on different appendicular bones, assuming dependency between the different elements by absence of evidence that the 'bauplan' of insular elephants had changed drastically. We found that estimations of  $S_i$  based on a single element versus based on all three

elements produced similar results (see Appendix S1 following Table S3). We also based our body weight estimates on length measurements only (as opposed to minimum circumferences of long bone diaphyses) because it has been observed that stegodonts had comparatively sturdy long bones as compared to modern elephants (Hooijer, 1955).

For species for which no suitable postcranial elements were available,  $S_i$  was calculated as the ratio of the cubed linear dimensions of the lower third molar following Lomolino (2005). The problem with only using dental elements, however, is that on geological time-scales, molar dimensions have been found to lag behind in the initial stages of dwarfism (Lister, 1989, 1996). This also applies to palaeoloxodontine dwarf elephants, where decrease in humerus length (as a proxy for body size) and molar width (as a proxy for dental size) have initially different scaling rates (see Appendix S1).



**Figure 2** Simplified map showing the geographical position of the islands with insular Proboscidea used in this study. 1, California Channel Islands; 2, Sardinia; 3, Sicily; 4, Islands of the Aegean (Crete, Karpathos, Palaeo-Cyclades, Rhodos); 5, Cyprus; 6, Java; 7, Flores; 8, Sumba; 9, Timor; 10, Sulawesi; 11, Philippines; 12, mainland Japan and the Ryukyu Islands; 13, Wrangel. The coastline is based on a map from d-maps.com.

For *Stegodon* species of which no suitable postcranial was available, but adult mandibles instead, a new regression was established (see Appendix S1 Table S3). This regression is derived from femur/tibia/humerus lengths against the mandibular width of other *Stegodon* species, for which both long bones and (fragments of) mandibles of full-grown individuals could be measured (see Appendix S1).

The resulting dataset comprises 22 island–mainland pairs of fossil proboscideans and their calculated size indices ( $S_i$  values) on 17 islands worldwide (Fig. 2).

### Estimation of island area and isolation

Island area and isolation were estimated with GLOBAL MAPPER 15 based on published data on palaeogeography, geodynamics, sea level changes, extension of tectonically undisturbed marine deposits and offshore bathymetry. For most of the islands in the Wallacea region, we used the map of Hanebuth *et al.* (2011) representing the palaeogeography at 12.7 kyr BP, when the sea level was 60 m below present. This can be considered as the ‘mean’ sea level that prevailed for the longest periods of time during the Quaternary (calculated from Pleistocene sea levels as given in Bintjana *et al.*, 2005). A sea level of –60 m also applies to the Mediterranean, following Lambeck & Chappell (2001) for the late Pleistocene. We assume here that body mass evolution in proboscideans takes a considerable time, and thus likely depends more on mean surface area rather than on the relatively short-lived extreme values. We infer this from the observation that Holocene endemics did not reach the same drastic body size decrease as Pleistocene endemics did (Lomolino *et al.*, 2013). Initially, dwarfism may evolve rapidly (in a few thousand years), but is then followed by a second stage of slower, but ultimately much more pronounced size change (Lister, 1989, 1996; van der Geer *et al.*,

2013). We are aware that –60 m may be somewhat arbitrary, but since we apply this level to all islands, the results are biased in the same direction, and remain the same when comparing islands mutually.

Sumba and Timor are exceptional cases, because their late Pleistocene uplift far exceeded the eustatic sea level fluctuations. The strong recent tectonic uplift of Sumba has a rate of 0.2–0.5 m kyr<sup>-1</sup>, as evidenced by dated raised coral terraces (Bard *et al.*, 1996). We have taken the 500-m contour level (the highest coral terrace is 475 m above present-day sea level) to calculate the palaeo-surface area. Timor has also been strongly uplifted, at least 700 m during the last 200,000 kyr (de Smet *et al.*, 1990), and therefore we have taken the 500-m contour as a rough estimate of the middle Pleistocene coastline of Timor. Flores in contrast, appears not to have been uplifted in such a dramatic way as Timor and Sumba were. The recent uplift of Flores is much less substantial, as indicated by the flat-lying terrestrial sediments of early and middle Pleistocene age now occurring at an elevation of c. 350 m above present sea level (Brumm *et al.*, 2010), unlike the highly uplifted Pleistocene marine coralline sediments of similar ages on Timor–Sumba. We, therefore, have estimated the area and isolation for Flores in a similar way as for most other islands, at –60-m sea level.

The following islands have alternative values for either area or isolation (for details, see Appendix S1). Crete in the early Pleistocene may have still consisted of two islands as it did at the end of the Pliocene (van Hinsbergen & Meulenkamp, 2006), or alternatively, was already merged due to the uplift into a single island as suggested by the presence of *Kritimys* in both parts. The uncertainty is caused by the uncertain age of the Cretan dwarf mammoth (Herridge & Lister, 2012). Flores (Pleistocene) was isolated from the mainland via Komodo, Sumbawa and Lombok or alternatively, via Sulawesi and the Philippines, based on ocean currents and the fossil record (Morwood & Jungers, 2009; Dennell *et al.*, 2014). Luzon (late Pleistocene) was part of the palaeo-island Greater Luzon (Heaney *et al.*, 2005) or alternatively, its northern part (where the fossils come from) constituted a separate island during much of the Pleistocene, separated from the southern part by a NW–SE trending fault zone (Defant & Ragland, 1988) at present bordered to the south by the low-lying Central Valley. Luzon (late Pleistocene) was isolated from the mainland via Mindoro and Palawan or alternatively, via Mindoro, Palawan and Borneo, assuming that Palawan was not connected to Borneo. Sicily (early Middle Pleistocene) still consisted of two palaeo-islands, or alternatively of one larger, merged island (see palaeo-maps in Bonfiglio *et al.*, 2002; partially based on endemism of the fauna, implying a potential circularity), which influences the estimated values for both area and isolation. Furthermore, a connection between Sicily and Malta is unproven (Herridge, 2010), and if a connecting ridge was ever exposed, it likely was merely a series of stepping stones (van der Geer *et al.*, 2010) as exposure of a continuous land bridge would require a drop in sea level of at least 120–150 m. Because of these uncertainties we have not included

the Maltese material attributed to *P. falconeri* into our analysis. South-western Sulawesi during the early Pleistocene still constituted a separate palaeo-island or alternatively, may have been already merged with the other parts of Sulawesi at some stage during the middle or late Pleistocene (van den Bergh, 1999).

### Taxonomic and phylogenetic framework

Here, we use the generic name *Palaeoloxodon* instead of *Elephas* for the mediterranean insular and mainland straight-tusked elephants (following Shoshani *et al.*, 2007; Todd, 2010). *Palaeoloxodon* 'mnaidriensis', referred to as *Palaeoloxodon* nov. sp. in Herridge (2010), still awaits revision. The 'large-sized *Elephas*' from Sulawesi (*Palaeoloxodon* sp., Tarrung Formation) is here considered conspecific with, or at least identical in size to *P. namadicus* (following van den Bergh, 1999).

Ancestral taxa for each insular species are based on the literature (Roth, 1982; van den Bergh, 1999; Palombo, 2001; van der Geer *et al.*, 2010; Herridge, 2010; Herridge & Lister, 2012). For mainland *Palaeoloxodon*, we use specimens from Italy and Greece, that are considered ancestral to the insular populations. In case an insular species was derived from a mainland taxon that must have migrated via an intermediate island (Flores, Timor, Sumba), we calculated body size index relative to the mainland ancestor, and not relative to the dwarfed taxon from the intermediate island. Most likely, the ancestral species from the intermediate island did not disperse to the island farther away following drastic reduction in body size on the intermediate island, but more likely this happened relatively soon after colonization when it was still large-bodied. This inference is considered the most likely scenario, because smaller body size for swimming elephants would signify reduced dispersal probability due to more restricted fasting endurance and reduced buoyancy (pneumatic bone tissue is lost in dwarf elephants) (Sondaar, 1977).

The *S. trigonocephalus* sample (Java) includes various subspecies of different sizes. Here, we only use the Trinil H.K. material, which forms a homogenous sample from a single layer estimated to 0.9 Ma (van den Bergh, 1999; but 0.54–0.43 Ma in Joordens *et al.*, 2014), when Java constituted an island, based on the endemic character of the fauna from this site (van den Bergh, 1999). The alternative scenario is that Java was connected through a filtered corridor with the mainland (Larick *et al.*, 2000; see for an extensive discussion of pros and cons, Meijaard, 2004). Evidence for at least a partial isolation is provided by the presence of the following endemic taxa at Trinil H.K.: *Axis lydekkeri*, *Bibos palaesondai-cus*, *Bubalus palaeokerabau*, *Duboisia santeng*, *Mececyon trinilensis* (= *Cuon* in Louys, 2014; here, we follow Lyras *et al.*, 2010, awaiting a revision of the Javanese canids), and *Sus brachygnathus*.

*Palaeoloxodon leonardi* (Sicily) and *P. sp.* (Delos) are excluded because their stratigraphic positions remain unre-

solved. This also applies to the significantly larger elephant species (*P. xylophagou*) of Cyprus (Athassiou *et al.*, 2014, 2015). The indeterminate Proboscidea and *E. or P. beyeri* from Luzon (de Vos & Bautista, 2003) are omitted because their ancestry, chronology and context are insufficiently known.

### Ecological assemblages

The number of competitors and predators most likely to have directly interacted with the focal insular species are based on published fauna lists (van den Bergh, 1999; van der Geer *et al.*, 2010) for local faunal assemblages containing the focal species. Judgement on the likelihood of competition or predation were based on the consultation of references on the diet and habitats of those species in order to determine which ones were likely to be significant competitors or predators of the focal insular species. Reptiles were not considered as potential interacting taxa because of their low energetic demands.

Late middle to late Pleistocene Sicily, however, poses a problem. The dwarf elephant belongs to two successive faunal units or complexes (*P. mnaidriensis* FC and San Teodoro FC) (Bonfiglio *et al.*, 2000). The elephant sample we use here is from Puntali Cave, dated either late middle Pleistocene (c. 180 ka) or late Pleistocene, comparable to that of San Teodoro (c. 32 ka) (Ferretti, 2008). We here tentatively include faunal elements from both FC's, with the exception of the lion, which we omit as it is restricted to San Ciro cave and Canita. Elements not retrieved from Puntali Cave itself are indicated with a question mark.

### Statistical analyses

We first tested for the normality of our data (body mass index, area, isolation) by using the Shapiro–Wilk test (Shapiro & Wilk, 1965). Then we used two-tailed Pearson's test to check for correlations between the log-transformed variables (significance level  $\alpha = 0.05$ ). For significantly correlated variables, a simple linear regression analysis (parametric) was used to test for the goodness-of-fit ( $r^2$ ; proportion of variability in the response) of the correlation. We used an independent samples' *t*-test to compare the insular body size evolution of the various genera. Statistical analyses were performed using IBM SPSS STATISTICS FOR WINDOWS 21. A 1-step ANOVA was used to test the effect of the presence of competitors on body size evolution.

For the development of a regression to estimate body size with mandibular width measurements, a reduced major axis regression on log-transformed (base 10) data was used.

## RESULTS

As illustrated in Table 1 and consistent with Prediction 1, body size variation among insular proboscideans matches previously reported patterns for extant mammals and other

vertebrates and was entirely consistent with the island rule. That is, all proboscidean species except one had a smaller body size than the respective mainland ancestral species ( $n = 23$ ; mean  $S_i = 0.28$ ,  $SD = 0.24$ ). The most pronounced case of size reduction is exhibited by *P. falconeri* (Sicily;  $S_i = 0.02$ ) and *P. cypriotes* (Cyprus;  $S_i = 0.02$ ) for *Palaeoloxodon*, *M. creticus* (Crete;  $S_i = 0.04$ ) for *Mammuthus* and *S. sumbaensis* (Sumba;  $S_i = 0.08$ ) for *Stegodon*. No size reduction is observed in the large *Elephas* species from Sulawesi. Limited size reduction is seen in *S. trigonocephalus* (Java, Trinil) ( $S_i = 0.65$ ). On a whole, *Stegodon* species ( $n = 10$ ) show a lesser body size reduction (or higher  $S_i$ ) than *Palaeoloxodon* ( $n = 8$ ), with a respective mean  $S_i = 0.33$  ( $SD = 0.19$ ) and  $S_i = 0.16$  ( $SD = 0.13$ ). This difference is significant [ $t(16) = -2.20$ ,  $P = 0.04$ ]. The number of *Mammuthus* species ( $n = 4$ ) is too low to be included in the  $t$ -test.

The log-transformed (base 10) variables of surface area, alternative surface area and untransformed minimal isolation were normally distributed, but maximal isolation is not (for SPSS output, see Appendix S1, after Table S4). However, the normality of maximal isolation is only minimally violated, and we proceed with it as if normally distributed.

Neither minimal nor maximal isolation is statistically correlated with insular body size index (Pearson  $R = -0.09$  and  $-0.05$ , respectively, with  $P = 0.69$  and  $P = 0.84$ ). Surprisingly, some islands within visual range (12 km) have very small elephants ( $S_i \leq 0.20$ ). There is no trend ( $P = 0.08$ ) for increased dwarfism on islands further away than 48 km (the maximum reported distance covered by swimming for Asian elephants in Johnson, 1980).

Log surface and log surface alternative are very strongly correlated with each other (Pearson  $R = 0.96$ ,  $P < 0.01$ ) and with  $S_i$  (Pearson  $R = 0.48$ ,  $P < 0.05$  and  $0.53$ ,  $P < 0.05$ , respectively). Based on this result, we proceeded with the first only, because this is considered the most reliably estimated value for any given island. The correlation between island area and body size index is positive but weak, with area explaining only 23% of the total variance ( $r^2 = 0.23$ ,  $P = 0.02$ ). The residuals show no correlation with latitude ( $r^2 = 0.02$ ,  $P = 0.52$ ). The residuals for *Stegodon* only, however, show a single outlier: *S. aurorae* from Japan. The difference from the predicted value is  $-0.26$  compared to  $-0.17$  to  $0.09$  for the other species. *Stegodon aurorae* occurred at a much higher latitude as compared to all other *Stegodon* species. Considering the three genera separately, the correlation is stronger and more significant for *Stegodon* ( $r^2 = 0.83$ ,  $P < 0.01$ ; with the tentative exclusion of the Japanese stegodont based on latitude) than for all taxa taken together.

The effect of competition (Table 2) on body size variation in proboscideans is limited (on average,  $S_i = 0.41$  with competitors against  $S_i = 0.17$  without competitors) but significant ( $P < 0.01$ ). Within the group lacking competitors, there is one less size-reduced species (*S. florensis florensis*,  $S_i = 0.51$ ). This species is a 700 kyr older chronosubspecies of the much smaller form (*S. florensis insularis*,  $S_i = 0.17$ ).

The effect of predation remains unclear. Given the lack of large terrestrial predators on the focal islands (except for three or five islands, depending on whether hominins can be considered as effective predators; Table 2), we were unable to assess the potential effect of predation on body size variation in these very large mammals. However, between species that occur on the same island but during a different time period, the species with predators show a smaller size reduction than the species without predators. This is shown by *P. mnaidriensis* versus *P. falconeri* and perhaps by *S. florensis* versus *S. sondaari*. Admittedly, a different geological period likely implies different vegetation, area, isolation etc. as well, so the interpretation of this observation is tentative.

## DISCUSSION

### Temporal variation in body size of proboscideans

Fossil proboscideans exhibit much more extreme cases of dwarfism than living species ( $< 5\%$  the body mass of their ancestors; Fig. 1). One plausible explanation is that these differences in degrees of body size evolution between extant mammals and those studied here is simply a function of the much more extended geological period over which the evolutionary change could be analysed in the fossil record, rather than the comparatively short time span of  $c. 15$  kyr since the last sea-level rise during which size reduction occurred in most extant mammals occurring on continental islands, as was also suggested by Lomolino *et al.* (2013).

### Factors associated with body size variation in proboscideans

Body size reduction in proboscideans appears to be most pronounced for the species occurring on the most isolated palaeo-island: *P. falconeri* of south-eastern Sicily, whether its isolation is 100 or 200 km. Yet, when considering all species together, no significant correlation was found between isolation and the degree of body size reduction. A possible explanation for the lack of a correlation may be that isolation *per se* is not a good estimator for accessibility. For example, despite the great distance, Sulawesi appears to have been comparatively 'easy' to colonize, if we take into account the comparatively large number of terrestrial immigrants that were able to cross Makassar Strait: apart from murine rodents, also squirrels, bovids, suids, macaques and hominins reached Sulawesi at some stage prior to the late Pleistocene (de Bruyn *et al.*, 2014; van den Bergh *et al.*, 2016). These are all clades that are not typical island taxa. It is likely that prevailing oceanic currents are a major factor in determining island accessibility (Ali & Huber, 2010), but these are difficult to incorporate into a generalized isolation index. Similarly, despite the relatively small distance between Flores and the mainland (one or two crossings with visibility of the target island) the strong Indonesian through-flow currents

**Table 2** Presence of ecologically relevant competitors and predators for the insular species mentioned in Table 1. Predators < c. 15 kg (e.g. otters, foxes) are not considered potentially relevant for proboscideans and omitted here. EP, early Pleistocene; MP, middle Pleistocene; LP, late Pleistocene; P, Pleistocene (no further stratigraphical evidence); H, Holocene.

Species	Palaeo-island	Geological time	Competitors	Predators
<i>E. sp.</i>	Sulawesi	MP	Elephant ( <i>Stegodon</i> sp. B), bovid? ( <i>Bubalus depressicornis</i> ), suid? ( <i>Celebochoerus</i> sp.)	–
<i>M. creticus</i>	Crete	EP	–	–
<i>M. exilis</i>	Santa Rosae	LP	–	–
<i>M. lamarmorai</i>	Sardinia	LP	Deer ( <i>Praemegaceros cazioti</i> )	–
<i>P. creutzburgi</i>	Crete	LP	Deer ( <i>Candiacervus</i> spp.)	–
<i>P. aff. creutzburgi</i>	Kassos	LP	Deer (' <i>Candiacervus</i> ' sp.)	–
<i>Palaeoloxodon cypriotes</i>	Cyprus	LP	–	–
<i>P. falconeri</i>	Sicily	early MP	–	–
<i>P. 'mnaidriensis'</i>	Sicily	late MP–LP	Deer ( <i>Cervus elaphus</i> , <i>Dama carburangelensis</i> ?), bovids ( <i>Bos primigenius</i> , <i>Bison priscus</i> )	Hyena ( <i>Crocuta crocuta</i> ), wolf ( <i>Canis lupus</i> )?
<i>P. sp.</i>	Rhodos	LP?	–	–
<i>P. lomoloini</i>	Naxos	LP	–	–
<i>P. tiliensis</i>	Tilos	LP–H	–	–
<i>S. aurorae</i>	Japan	EP	Deer ( <i>Elaphurus</i> spp., <i>Cervus</i> sp.), rhino ( <i>Rhinoceros</i> sp.)	Wolf ( <i>Xenocyon falconeri</i> )
<i>S. florensis florensis</i>	Flores	MP	–	Hominin ( <i>Homo floresiensis</i> )?
<i>S. florensis insularis</i>	Flores	LP	–	Hominin ( <i>Homo floresiensis</i> )?
<i>S. luzonensis</i>	Luzon	P	Elephants (Proboscidea indet., <i>Elephas beyeri</i> ), deer ( <i>Cervus</i> sp.), rhino ( <i>Rhinoceros luzonensis</i> )	–
<i>S. sompoensis</i>	Sulawesi	EP	Elephant ( <i>Stegoloxodon celebensis</i> ), suid ( <i>Celebochoerus heekereni</i> )	–
<i>S. sondaari</i>	Flores	EP	–	–
<i>S. timorensis</i>	Timor	MP	–	–
<i>S. sp. B</i>	Sulawesi	MP	Elephant ( <i>Elephas</i> sp.), bovid? ( <i>Bubalus depressicornis</i> ), suid? ( <i>Celebochoerus</i> sp.)	–
<i>S. trigonocephalus</i>	Trinil (Java)	late EP	Deer ( <i>Axis lydekkeri</i> , <i>Rusa timorensis</i> ), large bovids ( <i>Bubalus palaeokerabau</i> , <i>Epileptobos groeneveldti</i> , <i>Bibos palaeondaicus</i> ), elephant ( <i>E. hysudricus</i> ), rhino ( <i>Rhinoceros sondaicus</i> )	Tiger ( <i>Panthera tigris</i> ), hyena ( <i>Pachycrocuta brevirostris</i> ), hominin ( <i>Homo erectus</i> )?
<i>S. sumbaensis</i>	Sumba	MP–LP	–	–

(Sprintall *et al.*, 2014) that pass through these gaps may have significantly undermined overseas dispersal to Flores.

Body size appears positively but weakly correlated with island area for proboscideans as a whole. The relationship is much more robust for *Stegodon*, if we exclude the most northern species (*S. aurorae*), which appears much smaller than would be predicted, and the earlier chronosubspecies of Flores (*S. florensis florensis*).

Our results provide only limited support for a direct influence of predation on body size of insular proboscideans (but see Meiri *et al.*, 2008b; account of dwarfism in extant, large mammals, which may have been associated with reduced predator pressures of Borneo. This may stem at least in part from the limited variability in predator assemblages on the islands we studied. Nevertheless, temporal variation in ecological assemblages on Flores reveals the lack of a significant effect of predators on body size evolution of native stegodonts in the long run. Here, a geologically older pygmy stegodont (*Stegodon sondaari*) was replaced by a larger stegodont (*S. florensis*), coinciding with the arrival of a hominin. The latter apparently had access to carcasses as attested by cut marks on stegodont bones in the site Liang Bua (van

den Bergh *et al.*, 2009). Yet, even with this putative predator around, the larger *Stegodon* decreased in size between the middle Pleistocene (Soa Basin; subspecies *florensis*) and the latest Pleistocene (Liang Bua; subspecies *insularis*) (van den Bergh *et al.*, 2008), and eventually its degree of dwarfism practically rivalled that of *S. sondaari* ( $S_i = 0.17$  and  $0.15$  respectively). It seems, thus, that the effect of predation on proboscideans was either modest or, perhaps more likely, a function of the diversity and intensity of predators. After all, a single, hominin predator is likely not as significant as an assemblage of large felids and other mega-carnivores as present in mainland settings. In addition, *Stegodon* predation by hominins on Flores may have been a relatively late development, as there are no indications for active hunting or butchering of stegodont remains in the various middle Pleistocene assemblages from the So'a Basin (van den Bergh *et al.*, 2014).

The presence of other large herbivores on the other hand has a significant influence on the degree of dwarfism of insular elephants. Species with other proboscideans as a competitor all show a less dramatic size reduction than species with other competitors or no competitors at all. The strong influ-

ence of competition on body size evolution is in line with the model of Lomolino *et al.* (2012), and was also noticed for mediterranean elephants by Palombo (2009).

## CONCLUSIONS

This study provides further support for the applicability of the island rule to the largest terrestrial mammalian herbivorous taxa (see Benton *et al.*, 2010 and Stein *et al.*, 2010 for intriguing indications that very large herbivorous dinosaurs such as titanosaurian sauropods may have also followed the predicted pattern).

The patterns discussed here for body size variation in proboscideans over space and time do not support a significant relationship between isolation and body size decrease for proboscideans. Islands relatively close to the mainland may harbour extremely size-reduced proboscideans as well as normal-sized species. Islands further away than 48 km (the maximum reported distance an Asian elephant can swim) do not harbour the smallest species.

The correlation between area and body size is positive but weak with much scatter around the trend for all proboscideans, but more robust for *Stegodon* from low latitude islands. Our observations support an ecological hypothesis of body size evolution, inferred from the significant influence of competition on body size evolution (see also Palombo, 2009). On the mainland and on relatively balanced and ecologically rich palaeo-islands such as Sulawesi, interaction with ecologically relevant species resulted in a (relatively) large body size. On islands with more depauperate assemblages, however, release from these ecological interactions appears to have resulted in a smaller body size. In the absence of competitors, body size tends to trend towards a size positively correlated with island area, provided that deep geological time was available to allow the dwarfing process to proceed beyond the initial phases.

## ACKNOWLEDGEMENTS

We are grateful to George Iliopoulos, Victoria Herridge, Per Christianssen, Adrian Lister, Fachroel Aziz, Anwar Akib and Thomas Sutikna who provided useful information or support in one way or the other. We thank Elisabetta Cioppi, Reinier van Zelst, John de Vos, Loic Costeur, Martin Schneider, Andrew (Andy) Curren, Spyridoula Pappa, Jessica McDonald, Khairudin, Bambang Sulisyanto and Iwan Kurniawan for permission to access the collections under their care. Victoria Herridge kindly provided measurements of Siwalik stegodonts. We further thank Tama McGlenn and Susan Hayes for correcting the English of the final manuscript and Effie Mitsopoulou for her help with the statistical analyses. Lastly, we want to thank the anonymous referees for their suggestions and comments on earlier versions of this manuscript. This research has been co-financed by the European Union (European Social Fund – ESF) and Greek national funds through the Operational Program ‘Education and Lifelong

Learning’ of the National Strategic Reference Framework (NSRF) – Research Funding Program: THALIS – UOA-Island biodiversity and cultural evolution: Examples from the Eastern Mediterranean, Madagascar, Mauritius and Philippines during the past 800,000 years (MIS375910, KA:70/3/11669). Fieldwork and lab research in Indonesia of G.D. v.d.B., U.P.W. and E.S. was funded by the Australian Research Council (grants FT100100384 and DP1093342), and funds from the Geological Survey Institute of Indonesia.

## REFERENCES

- Ali, J.R. & Huber, M. (2010) Mammalian biodiversity on Madagascar controlled by ocean currents. *Nature*, **463**, 653–656.
- Ambrosetti, P. (1968) The Pleistocene dwarf elephant of Spinagallo. *Geologica Romana*, **7**, 277–398.
- Athanassiou, A., Reese, D., Iliopoulos, G., Herridge, V., Roussiakis, S., Mitsopoulou, V., Tsiolakis, E. & Theodorou, G. (2014) The endemic elephants of Cyprus: a reconsideration of their variation and taxonomy based on fossil finds. *Abstract Book of the VIth International Conference on Mammoths and their Relatives*. Scientific Annals, School of Geology, Aristotle University of Thessaloniki, Greece, Special Volume, **102**, 24.
- Athanassiou, A., Herridge, V., Reese, D.S., Iliopoulos, G., Roussiakis, S., Mitsopoulou, V., Tsiolakis, E. & Theodorou, G. (2015) Cranial evidence for the presence of a second endemic elephant species on Cyprus. *Quaternary International*, **379**, 47–57.
- Bard, E., Jouannic, C., Hamelin, B., Pirazzoli, P., Arnold, M., Faure, G., Sumosusastro, P. & Syaefudin (1996) Pleistocene sea levels and tectonic uplift based on dating of corals from Sumba Island, Indonesia. *Geophysical Research Letters*, **23**, 1473–1476.
- Benton, M.J., Csiki, Z., Grigorescu, D., Redelstorff, R., Sander, P.M., Stein, K. & Weishampel, D.B. (2010) Dinosaurs and the island rule: the dwarfed dinosaurs from Hațeg Island. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **293**, 438–454.
- van den Bergh, G.D. (1999) The Late Neogene elephantoid-bearing faunas of Indonesia and their palaeozoogeographic implications; a study of the terrestrial faunal succession of Sulawesi, Flores and Java, including evidence for early hominid dispersal east of Wallace’s line. *Scripta Geologica*, **117**, 1–419.
- vanden Bergh, G.D., Brumm, A., Puspaningrum, M., Setiabudi, E., Kurniawan, I. & Prasetyo, U.W. (2014). Taphonomy of *Stegodon florensis* remains from the early Middle Pleistocene archaeological site Mata Menge, Flores, Indonesia. *Abstract Book of the VIth International Conference on Mammoths and their Relatives*. S.A.S.G., Special Volume, **102**, 207–208.
- van den Bergh, G.D., Due, Rokus.Awe., Morwood, M.J., Sutikna, T., Jatmiko, P. & Wahyu Saptomo, E. (2008) The youngest *Stegodon* remains in south-east Asia from the

- Late Pleistocene archaeological site Liang Bua, Flores, Indonesia. *Quaternary International*, **181**, 16–48.
- van den Bergh, G.D., Meijer, H.J.M., Due, Rokus.Awe., Morwood, M.J., Szabó, K., van den Hoek Ostende, L.W., Sutikna, T., Saptomo, E.W., Piper, P.J. & Dobney, K.M. (2009) The Liang Bua faunal remains: a 95 k.yr. sequence from Flores, East Indonesia. *Journal of Human Evolution*, **57**, 527–537.
- van den Bergh, G.D., Li, B., Brumm, A., Grün, R., Yurnaldi, D., Moore, M.W., Kurniawan, I., Setiawan, R., Aziz, F., Roberts, R.G., Suyono, Storey, M., Setiabudi, E. & Morwood, M.J. (2016) Earliest hominin occupation of Sulawesi, Indonesia. *Nature*, **529**, 208–211.
- Bintjana, R., van de Wal, R.S.W. & Oerlemans, J. (2005) Modelled atmospheric temperatures and global sea levels over the past million years. *Nature*, **437**, 125–128.
- Bonfiglio, L., Marra, A.C. & Masini, F. (2000) The contribution of the Quaternary vertebrates to the palaeoenvironmental and palaeoclimatic reconstructions in Sicily. *Climates: past and present* (ed. by M.B. Hart), pp. 169–182. Special Publications, 181. Geological Society, London.
- Bonfiglio, L., Mangano, G., Marra, A.C., Masini, F., Pavia, M. & Petruso, D. (2002) Pleistocene Calabrian and Sicilian bioprovinces. *Geobios*, **24**, 29–39.
- Brumm, A., Jensen, G.M., van den Bergh, G.D., Morwood, M.J., Kurniawan, I., Aziz, F. & Storey, M. (2010) Hominins on Flores, Indonesia, by one million years ago. *Nature*, **464**, 748–752.
- de Bruyn, M., Stelbrink, B., Morley, R.J., Hall, R., Carvalho, G.R., Cannon, C.H., van den Bergh, G.D., Meijaard, E., Metcalfe, I., Boitani, L., Maiorano, L., Shoup, R. & von Rintelen, T. (2014) Borneo and Indochina are major evolutionary hotspots for Southeast Asian biodiversity. *Systematic Biology*, **63**, 879–901.
- Christiansen, P. (2004) Body size in proboscideans, with notes on elephant metabolism. *Zoological Journal of the Linnean Society*, **140**, 523–549.
- Defant, M.J. & Ragland, P.C. (1988) Recognition of contrasting magmatic processes using SB-systematics: an example from the western Central Luzon arc, the Philippines. *Chemical Geology*, **67**, 197–208.
- Dennell, R.W., Louys, J., O'Regan, H.J. & Wilkinson, D.M. (2014) The origins and persistence of *Homo floresiensis* on Flores: biogeographical and ecological perspectives. *Quaternary Science Reviews*, **96**, 98–107.
- Ferretti, M.P. (2008) The dwarf elephant *Palaeoloxodon mnaidriensis* from Puntali Cave. Carini (Sicily; late Middle Pleistocene): anatomy, systematics and phylogenetic relationships. *Quaternary International*, **182**, 90–108.
- van der Geer, A., Lyras, G., de Vos, J. & Dermitzakis, M. (2010) *Evolution of island mammals: adaptation and extinction of placental mammals on islands*. Wiley-Blackwell, Oxford.
- van der Geer, A., Lyras, G.A., Lomolino, M.V., Palombo, M.R. & Sax, D.F. (2013) Body size evolution of palaeo-insular mammals: temporal variations and interspecific interactions. *Journal of Biogeography*, **40**, 1440–1450.
- van der Geer, A.E., Lyras, G.A., van den Hoek Ostende, L.W., de Vos, J. & Drinia, H. (2014) A dwarf elephant and a rock mouse on Naxos (Cyclades, Greece) with a revision of the palaeozoogeography of the Cycladic Islands (Greece) during the Pleistocene. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **404**, 133–144.
- Hanebuth, T.J.J., Voris, H.K., Yokoyama, Y., Saito, Y. & Okuno, J. (2011) Formation and fate of sedimentary depocentres on Southeast Asia's Sunda Shelf over the past sea-level cycle and biogeographic implications. *Earth-Science Reviews*, **104**, 92–110.
- Heaney, L.R., Walsh, J.S., Jr & Peterson, A.T. (2005) The roles of geological history and colonization abilities in genetic differentiation between mammalian populations in the Philippine archipelago. *Journal of Biogeography*, **32**, 229–247.
- Herridge, V.L. (2010) *Dwarf elephants on Mediterranean islands: a natural experiment in parallel evolution*. PhD Thesis, University College, London.
- Herridge, V.L. & Lister, A.M. (2012) Extreme insular dwarfism evolved in a mammoth. *Proceedings of the Royal Society B: Biological Sciences*, **279**, 3139–3200.
- van Hinsbergen, D.J.J. & Meulenkaamp, J.E. (2006) Neogene supradetachment basin development on Crete (Greece) during exhumation of the South Aegean core complex. *Basin Research*, **18**, 103–124.
- Hooijer, D.A. (1955) Fossil Proboscidea from the Malay Archipelago and India. *Zoologische Verhandelingen*, **28**, 1–146.
- Itescu, Y., Karraker, N.E., Raia, P., Pritchard, P.C.H. & Meiri, S. (2014) Is the island rule general? Turtles disagree. *Global Ecology and Biogeography*, **23**, 689–700.
- Johnson, D.L. (1980) Problems in the land vertebrate zoogeography of certain islands and the swimming powers of elephants. *Journal of Biogeography*, **7**, 383–398.
- Joordens, J.C.A., d'Errico, F., Wesselingh, F.P., Munro, S. & deVos, J. (2014) *Homo erectus* at Trinil on Java used shells for tool production and engraving. *Nature*, **518**, 228–231.
- Lambeck, K. & Chappell, J. (2001) Sea level change through the last glacial cycle. *Science*, **292**, 679–686.
- Larick, R., Ciochon, R.L., Zaim, Y., Sudijono, Suminto., Rizal, Y. & Aziz, F. (2000) Lithographic context for Kln-1993.05-SNJ, a fossil colobine maxilla from Jokotingkir, Sangiran Dome. *International Journal of Primatology*, **21**, 731–759.
- Lister, A.M. (1989) Rapid dwarfing of red deer on Jersey in the Last Interglacial. *Nature*, **342**, 539–542.
- Lister, A.M. (1996) Dwarfing in island elephants and deer: processes in relation to time of isolation. *Symposia of the Zoological Society of London*, **69**, 277–292.
- Lomolino, M.V. (1985) Body size of mammals on islands: the island rule reexamined. *The American Naturalist*, **125**, 310–316.

- Lomolino, M.V. (2005) Body size evolution in insular vertebrates: generality of the island rule. *Journal of Biogeography*, **32**, 1683–1699.
- Lomolino, M.V., Sax, D.F., Palombo, M.R. & van der Geer, A.A. (2012) Of mice and mammoths: evaluations of causal explanations for body size evolution in insular mammals. *Journal of Biogeography*, **39**, 842–854.
- Lomolino, M.V., van der Geer, A.A., Lyras, G.A., Palombo, M.R., Sax, D.F. & Rozzi, R. (2013) Of mice and mammoths: generality and antiquity of the island rule. *Journal of Biogeography*, **40**, 1427–1439.
- Louys, J. (2014) The large terrestrial carnivore guild in Quaternary Southeast Asia. *Predators, Prey and Hominins, celebrating the scientific career of Alan Turner (1947-2012)* (ed. by H.J. O'Regan, S. Elton and D. Schreve). *Quaternary Science Reviews*, **96**, 86–97.
- Lyras, G.A., van der Geer, A. & Rook, L. (2010) Body size of insular carnivores: evidence from the fossil record. *Journal of Biogeography*, **37**, 1007–1102.
- Meijaard, E. (2004) *Solving mammalian riddles, a reconstruction of the Tertiary and Quaternary distribution of mammals and their palaeoenvironments in island South-East Asia*. PhD Thesis, Australian National University, Canberra, Australia.
- Meiri, S. (2007) Size evolution in island lizards. *Global Ecology and Biogeography*, **16**, 702–708.
- Meiri, S., Dayan, T. & Simberloff, D. (2004) Body size of insular carnivores: little support for the island rule. *The American Naturalist*, **163**, 469–479.
- Meiri, S., Dayan, T. & Simberloff, D. (2006) The generality of the island rule reexamined. *Journal of Biogeography*, **33**, 1571–1577.
- Meiri, S., Cooper, N. & Purvis, A. (2008a) The island rule: made to be broken? *Proceedings of the Royal Society B: Biological Sciences*, **275**, 141–148.
- Meiri, S., Meijaard, E., Wich, S.A., Groves, C.P. & Helgen, K.M. (2008b) Mammals of Borneo - small size on a large island. *Journal of Biogeography*, **35**, 1087–1094.
- Meiri, S., Raia, P. & Phillimore, A.B. (2011) Slaying dragons: limited evidence for unusual body size evolution on islands. *Journal of Biogeography*, **38**, 89–100.
- Morwood, M.J. & Jungers, W.L. (2009) Conclusions: implications of the Liang Bua excavations for hominin evolution and biogeography. *Journal of Human Evolution*, **57**, 640–648.
- Palombo, M.R. (2001) Endemic elephants of the Mediterranean Islands: knowledge, problems and perspectives. *The World of Elephants. Proceedings of the 1st International Congress; 16–20 Oct 2001, Rome* (ed. by G. Cavaretta, P. Gioia, M. Mussi and M.R. Palombo), pp. 486–491. CNR, Rome.
- Palombo, M.R. (2007) How can endemic proboscideans help us understand the “island rule”? A case study of Mediterranean islands. *Quaternary International*, **169–170**, 105–124.
- Palombo, M.R. (2009) Body size structure of the Pleistocene mammalian communities from Mediterranean Islands: what support for the “island rule”? *Integrative Zoology*, **4**, 341–356.
- Palombo, M.R. & Ferretti, M.P. (2005) The Italian elephant fossil record: knowledge, problems and perspectives. *Quaternary International*, **126–128**, 107–136.
- Raia, P. & Meiri, S. (2006) The island rule in large mammals: paleontology meets ecology. *Evolution*, **60**, 1731–1742.
- Roth, V.L. (1982) *Dwarf mammoths from the Santa Barbara, California Channel Islands: size, shape, development and evolution*. PhD Thesis, Yale University, New Haven, CT.
- Shapiro, S.S. & Wilk, M.B. (1965) An analysis of variance test for normality (complete samples). *Biometrika*, **52**, 591–611.
- Shoshani, J., Ferretti, M.P., Lister, A.M., Agenbroad, L.D., Saegusa, H., Mol, D. & Takahashi, K. (2007) Relationships within the Elephantinae using hyoid characters. *Quaternary International*, **169–170**, 174–185.
- de Smet, M.E.M., Fortuin, A.R., Troelstra, S.R., Van Marle, L.J., Karmini, M., Tjokrosapoetro, S. & Hadiwasastra, S. (1990) Detection of collision-related vertical movements in the Outer Banda Arc (Timor, Indonesia), using micropalaeontological data. *Journal of Southeast Asian Earth Sciences*, **4**, 337–356.
- Sondaar, P.Y. (1977) Insularity and its effect on mammal evolution. *Major patterns in vertebrate evolution* (ed. by M.N. Hecht and P.L. Goody), pp. 671–707. Plenum, New York.
- Sprintall, J., Gordon, A.L., Koch-Larrouy, A., Lee, T., Potemra, J.T., Pujiana, K. & Wijffels, S.E. (2014) The Indonesian seas and their role in the coupled ocean–climate system. *Nature Geoscience*, **7**, 487–492.
- Stein, K., Csiki, Z., Curry Rogers, K., Weishampel, D.B., Redelstorff, R., Carballido, J.L. & Sandera, P.M. (2010) Small body size and extreme cortical bone remodeling indicate phyletic dwarfism in *Magyarosaurus dacus* (Sauropoda: Titanosauria). *Proceedings of the National Academy of Sciences USA*, **107**, 9258–9263.
- Todd, N.E. (2010) New phylogenetic analysis of the family Elephantidae based on cranial and dental morphology. *Anatomical Record*, **293**, 74–90.
- Van Valen, L. (1973) Pattern and the balance of nature. *Evolutionary Theory*, **1**, 31–49.
- de Vos, J. & Bautista, A. (2003) Preliminary notes on the vertebrate fossils from the Philippines. *Semantics and systematics: Philippine archaeology* (ed. by A. de la Torre and V.J. Paz), pp. 42–62. National Museum, Manila.

## SUPPORTING INFORMATION

Additional Supporting Information may be found in the online version of this article:

**Appendix S1** Body mass estimations of palaeo-species, palaeo-island characteristics and statistical analyses.

## BIOSKETCH

**Alexandra A.E. van der Geer** is a palaeontologist and first author of *Evolution of island mammals: adaptation and extinction of placental mammals on islands*. Her current research focuses on evolutionary processes on islands, including body size evolution and morphological changes in the skeleton.

Author contributions: A.V.G., G.A.L., G.v.d.B conceived the ideas, planned the analysis and analysed the data; G.A.L., G.v.d..B., U.W., R.D.A., E.S. collected the data; G.A.L. prepared the figures; and A.V.G. led the writing assisted by G.A.L., G.v.d..B., U.W., R.D.A., E.S. and H.D.

---

Editor: Robert Whittaker