



Naturalis Repository

Ostreid herpesvirus OsHV-1 μ Var in Pacific oysters *Crassostrea gigas* (Thunberg 1793) of the Wadden Sea, a UNESCO world heritage site

A Gittenberger, M A Voorbergen-Laarman, M Y Engelsma

Downloaded from:

<https://doi.org/10.1111%2Fjfd.12332>

Article 25fa Dutch Copyright Act (DCA) - End User Rights

This publication is distributed under the terms of Article 25fa of the Dutch Copyright Act (Auteurswet) with consent from the author. Dutch law entitles the maker of a short scientific work funded either wholly or partially by Dutch public funds to make that work publicly available following a reasonable period after the work was first published, provided that reference is made to the source of the first publication of the work.

This publication is distributed under the Naturalis Biodiversity Center 'Taverne implementation' programme. In this programme, research output of Naturalis researchers and collection managers that complies with the legal requirements of Article 25fa of the Dutch Copyright Act is distributed online and free of barriers in the Naturalis institutional repository. Research output is distributed six months after its first online publication in the original published version and with proper attribution to the source of the original publication.

You are permitted to download and use the publication for personal purposes. All rights remain with the author(s) and copyrights owner(s) of this work. Any use of the publication other than authorized under this license or copyright law is prohibited.

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the department of Collection Information know, stating your reasons. In case of a legitimate complaint, Collection Information will make the material inaccessible. Please contact us through email: collectie.informatie@naturalis.nl. We will contact you as soon as possible.



Ostreid herpesvirus OsHV-1 μ Var in Pacific oysters *Crassostrea gigas* (Thunberg 1793) of the Wadden Sea, a UNESCO world heritage site

A Gittenberger^{1,2,3}, M A Voorbergen-Laarman⁴ and M Y Engelsma⁴

1 GiMaRIS, Marine Research Inventory & Strategy solutions, Leiden, The Netherlands

2 Institute of Biology Leiden (IBL), Leiden University, Leiden, The Netherlands

3 Department of Marine Zoology, Naturalis Biodiversity Center, Leiden, The Netherlands

4 Central Veterinary Institute, part of Wageningen UR, Lelystad, The Netherlands

Abstract

The Wadden Sea is an extensive wetland area, recognized as UNESCO world heritage site of international importance. Since the mid-1990s, the invasive Pacific oyster *Crassostrea gigas* (Thunberg 1793) population in the area has grown exponentially, having a distinct impact on the ecosystem. The recent spread of the emerging oyster pathogen Ostreid herpesvirus OsHV-1 μ Var worldwide and specifically in the oyster culture areas in the south of the Netherlands raised the question whether the virus may also be present in the Wadden Sea. In the summer of 2012 juvenile Pacific oysters were collected from five locations in the Dutch Wadden Sea. The virus was shown to be present in three of the five locations by real-time PCR and sequencing. It was concluded that OsHV-1 μ Var has settled itself in Pacific oyster reefs in the Wadden Sea. These results and the recent discoveries of OsHV-1 microvariants in Australia and Korea indicate that OsHV-1 μ Var and related variants might be more widespread than can be deduced from current literature. In particular in regions with no commercial oyster culture, similar to the Wadden Sea, the virus may go undetected as wild beds with mixed age classes hamper the detection of mortality among juvenile oysters.

Keywords: *Crassostrea gigas*, molecular detection, mollusc health, oyster herpesvirus, viral disease, Wadden Sea.

Introduction

The Wadden Sea is an extensive wetland area characterized by large intertidal flats, stretching from the Netherlands to Denmark. It is recognized as UNESCO world heritage site of international importance. The ecosystem in this area has an important role in a wide range of functions, varying from being a nursery area for commercially important fish and shrimp to providing a staging area for waterbirds migrating along the East-Atlantic flyway (Compton *et al.* 2013). The Pacific oyster *Crassostrea gigas* (Thunberg 1793) was first recorded in the Dutch Wadden Sea in 1983. Since then, the *C. gigas* population has grown exponentially from the mid-1990s onwards (Fey *et al.* 2009). The oysters are predominantly settling on intertidal blue mussel (*Mytilus edulis*) beds, which are increasingly transformed into *Crassostrea* reefs (Markert *et al.* 2013). The introduction of *Crassostrea gigas* has a distinct impact on the Wadden Sea ecosystem, among which a change in the foraging habitat of different bird species (Markert *et al.* 2013).

Mass mortalities of oysters are well-known phenomena occurring in different geographical regions for which various factors may play a role, including environmental factors (Petton *et al.* 2013; Green *et al.* 2014) and pathogens. Mass

Correspondence Adriaan Gittenberger, GiMaRIS, J.H. Oortweg 21, 2333 CH Leiden, The Netherlands (e-mail: Gittenberger@gimaris.com)

mortalities of adult Pacific oysters have occurred in the past in the Wadden Sea (Watermann *et al.* 2008; Fey *et al.* 2009; Büttger, Nehls & Witte 2011), usually linked to environmental causes like relatively warm summers or cold winters. The Pacific oyster is known to be susceptible to pathogens such as *Vibrio* spp. (Saulnier *et al.* 2010; Vezzulli *et al.* 2010) and Ostreid herpesvirus type 1 (OsHV-1; Nicolas, Comps & Cochennec 1992; Hine, Wesley & Hay 1992). Recently, a new genotype of oyster herpesvirus, OsHV-1 μ Var, was detected and associated with high mortalities, locally over 95%, of juvenile *C. gigas* (Segarra *et al.* 2010). OsHV-1 μ Var was first detected in France in 2008 and quickly spread over the main *C. gigas* culture areas in Europe. The virus was recorded along the coast of France (Segarra *et al.* 2010), Ireland (Peeler *et al.* 2012), England (OIE 2010), Spain (Roque *et al.* 2012), Italy (Dundon *et al.* 2011) and the Netherlands (Engelsma 2012) while outside Europe related OsHV-1 microvariants were detected in Australia (Jenkins *et al.* 2013) and Korea (Hwang *et al.* 2013). Infection with OsHV-1 microvariants has been recently listed as a notifiable disease for the World Organization of Animal Health (OIE 2014). In the Netherlands OsHV-1 μ Var has been detected since 2010 in Lake Grevelingen and the Oosterschelde, two Pacific oyster culture areas in the south-west of the Netherlands (Engelsma 2012). In the Dutch Wadden Sea, mortalities of juvenile Pacific oysters that could be linked to the presence of OsHV-1 μ Var have not been reported. However, mortality of juvenile oysters is difficult to assess in wild beds of mixed age classes. In 2011,

a preliminary attempt has been carried out to investigate the presence of OsHV-1 μ Var in the Wadden Sea with a very limited number of samples from Texel and Sylt (Thieltges *et al.* 2013). OsHV-1 μ Var could not be demonstrated in the samples at that time.

Taking into account the recent spread of the virus and its presence in the oyster culture areas in the Netherlands, it was questioned whether OsHV-1 μ Var could possibly be present in the oyster reefs in the Wadden Sea and if so, what the effect would be on the Pacific oyster population. To address the first question five oyster reefs in the Dutch Wadden Sea were screened in 2012 for the presence of OsHV-1 by real-time PCR. The OsHV-1 genotype was subsequently determined by sequencing of two OsHV-1 genome regions from positive samples.

Materials and methods

Samples were taken from five locations (see Fig. 1), selected to represent the various habitats in which oyster reefs are found in the Wadden Sea. They differed in being either sheltered or exposed, and in being either a reef dominated by Pacific oysters, *Crassostrea gigas*, or being a more mixed shellfish bed, which also included blue mussels, *Mytilus edulis* (L.) and common cockles, *Cerastoderma edule* (L.) (Table 1).

As outbreaks of OsHV-1 μ Var are reported at temperatures above 16 °C (Dégremont *et al.* 2013), the optimal temperature for the virus appears to be above this temperature. Hence, the sampling was conducted at the end of the summer in the last week




Figure 1 Locations sampled: 1, Het Abt; 2, Schiermonnikoog; 3, Vierhuistergat; 4, Eemshaven; 5, Termunterzijl.

Table 1 Sampling locations, habitat descriptions of the locations, number of Pacific oysters tested from each location and results of real-time PCR analysis on OsHV-1

Location	Location description	# Pacific oysters tested	# Tested positive for OsHV-1
1 Het Abt	Mixed shellfish bed with mainly oysters, cockles and mussels	30	0
2 Schiermonnikoog	Large shellfish bed with small tidal pools that lie close to the island. locally dominated by either mussels or oysters	30	9
3 Vierhuistergat	Seaweeds, oysters and mussels lying scattered around a jetty. Oysters of all ages were present, but the larger ones were rare	30	3
4 Eemshaven	The sheltered side of a riprap dike in the harbour, mostly overgrown by oysters of all sizes. A riprap dike is a loose assemblage of broken stones erected in water as a foundation	30	8
5 Termunterzijl	A jetty, locally covered with seaweeds. Large numbers of oysters of which only a few were found to be alive	17	0

of August and the first weeks of September of 2012, when water temperatures in the Wadden Sea are at their annual highest (around 20 °C).

At each location, 30 juvenile oysters (<2.0 cm shell length) were collected, with the exception of the location Termunterzijl where only 17 individuals were collected, as most of the oysters at that location appeared to have died recently by unknown causes. At the other locations, no abnormal mortalities were observed. All 137 individuals were screened for the presence of OsHV-1 μ Var by real-time PCR. In brief, after collection, the samples were frozen and stored at -20 °C until further processing. Of each individual oyster, the soft tissue was homogenized with a TeSeE PRECESS 24 homogenizer (Bio-Rad) in tissue disruption tubes equipped with ceramic beads. Total DNA was extracted using the automated MagNa Pure LC system (Roche). The TaqMan real-time PCR was performed using the primers and probes as described by Martenot *et al.* (2010): OsHV1 BF (5'-GTCGCATCTTTGGATTTAACA-3'), reverse B4 (5'-ACTGGGATCCGACTGACAAC-3') and the probe B (5'-TGCCCCTGTCATCTTGAGGTATAGACAATC-3'). The real-time PCR consisted of 1 \times TaqMan Fast Universal PCR Master Mix (Applied Biosystems) including 0.4 μ M of each primer, 0.2 μ M probe, 1.25 units of Uracil-DNA Glycosylase (New England Biolabs) and 5 μ L of template DNA to a final volume of 20 μ L per reaction. The real-time PCR was carried out in an AB 7500 with fast block (Applied Biosystems) according to the program: 37 °C for 10 min, 95 °C for 10 min followed by 40 cycles of 95 °C for 3 s and 60 °C for 30 s. Prior to testing of the samples, the limit of detection (LOD) of the assay was assessed by a 1/4

serial dilution of an OsHV-1 μ Var-positive field sample tested in eightfold. Based on this, the LOD of the assay was estimated to be around cycle threshold (Ct) 36.5. This value was subsequently used as cut-off value for the assay.

The genotype of OsHV-1 was determined by amplification and direct sequencing of the C2/C6 region and the IA1/IA2 region from positive samples according to Segarra *et al.* (2010). In brief, the conventional PCR mixture consisted of 0.2 μ M of each primer C2 (5'-CTC TTT ACC ATG AAG ATA CCC ACC-3')–C6 (5'-GTG CAC GGC TTA CCA TTT TT-3') or IA1 (5'-CGC GGT TCA TAT CCA AAG TT-3')–IA2 (5'-AAT CCC CAT GTT TCT TG CTG-3') (Eurogentec), PCR buffer at 1 \times concentration, 2.5 mM MgCl₂, 200 μ M of each deoxynucleotide (dNTP mixture, Takara Bio Inc.), 2 units Taq DNA Polymerase (Invitrogen) and 5 μ L of template DNA. The thermal profile of the PCR was an initial denaturation at 94 °C for 2 min followed by 35 cycles at 94 °C for 1 min, 50 °C for 1 min, 72 °C for 1 min and a final extension at 72 °C for 5 min. After purification with QIAquick PCR Purification Kit (Qiagen), the PCR products were sequenced in both directions using an ABI Prism BigDye v 1.1 Terminator Cycle Sequencing kit (Applied Biosystems) and the corresponding amplification primers and analysed on an ABI Prism 3100 Avant Genetic Analyzer (Applied Biosystems).

Results

As result of the analyses, juvenile Pacific oysters were found positive for OsHV-1 by real-time PCR at three of the five locations (Table 1). At the OsHV-1 positive sites, the number of oysters

in which genomic material of OsHV-1 could be detected ranged from 3 of 30 to 9 of 30. For most positive samples, high Ct values were obtained, above Ct 30, indicative for a low amount of viral copies. This hampered subsequent genetic analysis. From only five samples (one from location 3 and four from location 2), sequences could be obtained by amplification and direct sequencing of the C2/C6 and IA1/IA2 region. The sequences from both genomic regions were 100% identical to sequences of OsHV-1 μ Var as described by Segarra *et al.* (2010) and hence contained the characteristic substitutions and micro-satellite zone of OsHV-1 μ Var in both regions.

Discussion

From these results, it can be concluded that OsHV-1 μ Var has reached the Wadden Sea and OsHV-1 has settled itself in at least three Pacific oyster reefs. As the prevalence in each of these populations was low, it remains uncertain whether the virus was truly absent in the remaining two populations or was missed there by undersampling. Many recently dead oysters were recorded during the fieldwork at Termunterzijl where only 17 alive juvenile oysters could be collected. None of these individuals was tested positive for OsHV-1 μ Var. However, due to the limited sample size, this does not necessarily exclude the presence of the virus as cause of the mortalities. The results indicate that the virus may be widespread in at least the Dutch part of the Wadden Sea.

It remains uncertain when the virus was introduced into the Wadden Sea. Considering the observation at distinct sites, the virus may have been present undetected for several years. In natural populations with Pacific oysters forming *Crassostrea* reefs, mortalities among spat and juveniles only will be difficult to detect. Empty shells of spat and juveniles are easily lost or overgrown by other organisms. For example, in 2011, OsHV-1 μ Var was detected in Pacific oyster spat collected on disc collectors from two locations in Lake Grevelingen, the Netherlands (Engelsma 2012). Although no mortality was observed in wild beds of Pacific oysters at the same locations, on the collectors, the cumulative mortality of juveniles ranged up to 46%.

The route of introduction of the virus into the Wadden Sea remains unknown. Transport of the infected host is one of the major risk factors in spread of disease (Murray *et al.* 2012). However,

the only recent oyster imports from abroad (Britain) into the Wadden Sea took place in the German part, north of Sylt (Moehler *et al.* 2011). This area is a few hundred kilometres up-current from the Pacific oyster populations that were found to be infected and there are no indications that oysters have recently been transported from north to south in the Wadden Sea. Other possible routes of introduction into the Wadden Sea are via infected oysters attached to ship hulls or transfer with planktonic larvae of the Pacific oyster along the general south–north current of Dutch coast. With regard to the latter, this might be possible either directly, as larvae can drift along in the sea-currents for 2–4 weeks (Rohfritsch *et al.* 2013), or indirectly via settled oysters on hard substrates like embankments, buoys and structures in harbours and off-shore windmill parks that function as stepping stones in between the Delta and the Wadden Sea.

The impact of OsHV-1 μ Var on the *C. gigas* population in the Wadden Sea could not be deduced from this study. Possibly, losses by OsHV-1 μ Var can be compensated by the massive spat fall of *C. gigas* in the area, as observed over the last decade, but this would be a subject for further studies. Thus far, OsHV-1 μ Var has been primarily detected in the culture areas in Europe. In wild populations, observations of mortalities among spat and juvenile molluscs present difficulties. The present results indicate that OsHV-1 μ Var might be more widespread than can be concluded from records in literature alone. Similarly to the Wadden Sea, the virus may go undetected especially in regions with no commercial oyster culture.

Acknowledgements

Ineke Roozenburg is greatly acknowledged for the assistance with analysing the samples. We like to thank the Producentenorganisatie van de Nederlandse Mosselcultuur for their support of this project. Jaap Vegter is thanked for the collection of the samples.

Publication History

Received: 6 October 2014

Revision received: 10 October 2014

Accepted: 11 November 2014

This paper was edited and accepted under the Editorship of Professor Ron Roberts.

References

- Büttger H., Nehls G. & Witte S. (2011) High mortality of Pacific oysters in a cold winter in the North-Frisian Wadden Sea. *Helgoland Marine Research* **65**, 525–532.
- Compton T.J., Holthuijsen S., Koolhaas A., Dekinga A., ten Horn J., Smith J., Galama Y., Brugge M., van der Wal D., van der Meer J., van der Veer H.W. & Piersma T. (2013) Distinctly variable mudscapes: distribution gradients of intertidal macrofauna across the Dutch Wadden Sea. *Journal of Sea Research* **82**, 103–116.
- Dégremont L., Guyader T., Tourbiez D. & Pépin J.-F. (2013) Is horizontal transmission of the Ostreid herpesvirus OsHV-1 in *Crassostrea gigas* affected by unselected or selected survival status in adults to juveniles? *Aquaculture* **408–409**, 51–57.
- Dundon W.G., Arzul I., Omnes E., Robert M., Magnabosco C., Zambon M., Gennari L., Toffan A., Terregino C., Capua I. & Arcangeli G. (2011) Detection of Type 1 Ostreid Herpes variant (OsHV-1 μ Var) with no associated mortality in French-origin Pacific cupped oyster *Crassostrea gigas* farmed in Italy. *Aquaculture* **314**, 49–52.
- Engelsma M.Y. (2012) *Verslag monitoring oester herpesvirus 2011; Resultaten van onderzoek naar de aanwezigheid van oester herpesvirus in Japanse oesters van het Grevelingenmeer en de Oosterschelde. CVI-rapport 12/CVI0098*. Centraal Veterinair Instituut, Lelystad [In Dutch].
- Fey F., Dankers N., Steenbergen J. & Goudswaard K. (2009) Development and distribution of the non-indigenous Pacific oyster (*Crassostrea gigas*) in the Dutch Wadden Sea. *Aquaculture International* **18**, 45–59.
- Green T.J., Montagnani C., Benkendorff K., Robinson N. & Speck P. (2014) Ontogeny and water temperature influences the antiviral response of the Pacific oyster, *Crassostrea gigas*. *Fish & Shellfish Immunology* **36**, 151–157.
- Hine P.M., Wesney B. & Hay B.E. (1992) Herpesviruses associated with mortalities among hatchery-reared larval Pacific oysters *Crassostrea gigas*. *Diseases of Aquatic Organisms* **12**, 135–142.
- Hwang J.Y., Park J.J., Yu H.H., Hur Y.B., Arzul I., Couraleau Y. & Park M.A. (2013) Ostreid herpesvirus 1 infection in farmed Pacific oyster larvae *Crassostrea gigas* (Thunberg) in Korea. *Journal of Fish Diseases* **36**, 969–972.
- Jenkins C., Hick P., Gabor M., Spiers Z., Fell S.A., Gu X., Read A., Go J., Dove M., O'Connor W., Kirkland P.D. & Frances J. (2013) Identification and characterisation of an ostreid herpesvirus-1 microvariant (OsHV-1 μ -var) in *Crassostrea gigas* (Pacific oysters) in Australia. *Diseases of Aquatic Organisms* **105**, 109–126.
- Markert A., Esser W., Frank D., Wehrmann A. & Exo K.-M. (2013) Habitat change by the formation of alien *Crassostrea*-reefs in the Wadden Sea and its role as feeding sites for waterbirds. *Estuarine, Coastal and Shelf Science* **131**, 41–51.
- Martenot C., Oden E., Travaillé E., Malas J.P. & Houssin M. (2010) Comparison of two real-time PCR methods for detection of ostreid herpesvirus 1 in the Pacific oyster *Crassostrea gigas*. *Journal of Virological Methods* **170**, 86–89.
- Moehler J., Wegner K.M., Reise K. & Jacobsen S. (2011) Invasion genetics of Pacific oyster *Crassostrea gigas* shaped by aquaculture stocking practices. *Journal of Sea Research* **66**, 256–262.
- Murray A.G., Marcos-Lopez M., Collet B. & Munro L.A. (2012) A review of the risk posed to Scottish mollusc aquaculture from *Bonamia*, *Marteilia* and oyster herpesvirus. *Aquaculture* **370–371**, 7–13.
- Nicolas J.L., Comps M. & Cochennec N. (1992) Herpes-like virus infecting Pacific-oyster larvae, *Crassostrea gigas*. *Bulletin of the European Association of Fish Pathologists* **12**, 11–13.
- OIE (2010) Immediate notification Oyster herpesvirus (OsHV-1, μ Var), United Kingdom 21/07/2010. WAHID Interface - OIE World Animal Health Information Database. http://www.oie.int/wahis_2/public/wahid.php/Reviewreport/Review?page_refer=MapEventSummary&creportid=9527
- OIE (2014) *Manual of Diagnostic Tests for Aquatic Animals 2014*. World Organization for Animal Health, Paris. <http://www.oie.int/en/international-standard-setting/aquatic-manual/access-online/>
- Peeler E.J., Reese R.A., Cheslett D.L., Geoghegan F., Power A. & Thrush M.A. (2012) Investigation of mortality in Pacific oysters associated with Ostreid herpesvirus-1 μ Var in the Republic of Ireland in 2009. *Preventive Veterinary Medicine* **105**, 136–143.
- Petton B., Pernet F., Robert R. & Boudry P. (2013) Temperature influence on pathogen transmission and subsequent mortalities in juvenile Pacific oysters *Crassostrea gigas*. *Aquaculture Environmental Interactions* **3**, 257–273.
- Rohfritsch A., Bierne N., Boudry P., Heurtebise S., Cornette F. & Lapègue S. (2013) Population genomics shed light on the demographic and adaptive histories of European invasion in the Pacific oyster, *Crassostrea gigas*. *Evolutionary Applications* **6**, 1064–1078.
- Roque A., Carrasco N., Andree K.B., Lacuesta B., Elandaloussi L., Gairin I., Rodgers C.J. & Furones M.D. (2012) First report of OsHV-1 microvar in Pacific oyster (*Crassostrea gigas*) cultured in Spain. *Aquaculture* **324–325**, 303–306.
- Saulnier D., De Decker S., Haffner P., Cobret L., Robert M. & Garcia C. (2010) A large-scale epidemiological study to identify bacteria pathogenic to Pacific oyster *Crassostrea gigas* and correlation between virulence and metalloprotease-like activity. *Microbial Ecology* **59**, 787–798.
- Segarra A., Pépin J.F., Arzul I., Morga B., Faury N. & Renault T. (2010) Detection and description of a particular Ostreid herpesvirus 1 genotype associated with massive mortality outbreaks of Pacific oysters, *Crassostrea gigas*, in France in 2008. *Virus Research* **153**, 92–99.
- Thieltges D.W., Engelsma M.Y., Wendling C.C. & Wegner K.M. (2013) Parasites in the Wadden Sea food web. *Journal of Sea Research* **82**, 122–133.
- Vezzulli L., Previati M., Pruzzo C., Marchese A., Bourne D.G., Cerrano C. & the VibrioSea Consortium (2010) *Vibrio* infections triggering mass mortality events in a warming Mediterranean Sea. *Environmental Microbiology* **12**, 2007–2019.
- Watermann B.T., Herlyn M., Daehne B., Bergmann S., Meemken M. & Kolodzey H. (2008) Pathology and mass mortality of Pacific oysters, *Crassostrea gigas* (Thunberg), in 2005 at the East Frisian coast, Germany. *Journal of Fish Diseases* **31**, 621–630.