



OPEN

DATA DESCRIPTOR

Synthesised database of wild bee and hoverfly records in Europe

Ahlam Sentil *et al.*[#]

Wild bees (Hymenoptera: Anthophila) and hoverflies (Diptera: Syrphidae), the two major groups of insect pollinators, are undergoing alarming declines worldwide, including Europe. The lack of accessible and verified spatial and temporal occurrence records currently challenges efforts to understand and mitigate this decline. Here, we compiled datasets from diverse sources, including taxonomists, national experts, public repositories, museum collections, published literature, verified open-access platforms, and aggregated datasets from previous European projects. The collected data were standardised, cleaned and validated by taxonomists and national experts. This collective effort resulted in two databases comprising more than 4.34 million and 1.04 million records for wild bees and hoverflies, respectively. The databases cover 97% of the European bee fauna (2,083 species out of 2,138 recorded in Europe) and 97% of the European hoverfly fauna (886 species out of 913 recorded in Europe). These standardised databases constitute essential resources for future assessments of status and trends, habitat associations, and other research and conservation initiatives to protect and understand wild pollinators on the European continent.

Background & Summary

The current rate of biodiversity loss is an unprecedented threat in the history of humanity. The associated loss of ecosystem services is impacting food provision, air and water quality, waste remediation, flood regulation, and the diversity of plants and associated organisms¹. Pollinators are a key element of this global biodiversity, providing vital pollination ecosystem services to domesticated and wild plants². Among the pollinators that are increasingly threatened by global changes are bees (Hymenoptera: Anthophila) and hoverflies (Diptera: Syrphidae). Two European Red Lists formally encapsulate the worries regarding their conservation status in Europe^{3,4}. Altogether, these Red Lists indicate that 17% of the 2,834 assessed species are threatened with extinction, while data is deficient for over 40% of all species. Available national studies suggest an even more alarming situation in some countries — for example, in Belgium, 33% of the assessed bee species were classified as threatened⁵, and in the Netherlands, 46% of hoverfly species were assessed as threatened, with the national extinction rate of Syrphidae accelerating dramatically since 2000⁶. At a local scale a decrease of 80% in insect biomass and 65% of species was recorded in Germany and the Netherlands^{7,8}. The main identified drivers of decline include habitat destruction and fragmentation, environmental pollution, climate change, species invasion, pesticides and pathogens spillover^{9–11}.

In response to growing evidence of pollinator declines, the European Union has developed a range of measures — from long-standing biodiversity frameworks to more recent, targeted actions. Foundational instruments such as the Convention on Biological Diversity, the Bern Convention, and EU nature legislation, including the Birds and Habitats Directives (which underpin the Natura 2000 network), remain central to biodiversity conservation, though not originally designed with pollinators in mind. Broader strategies like the EU Biodiversity Strategy, the European Green Deal, the Farm to Fork Strategy, and the newly adopted Nature Restoration Regulation also support pollinators by promoting sustainable land use, ecosystem recovery, and reduced pesticide use. In recent years, targeted initiatives have emerged, including the EU Pollinator Initiative, the revised EU Pollinator Initiative - New Deal for Pollinators¹², interregional projects such as SAPOLL (*SAuvons nos POLLinisateurs*), and EU-funded research and coordination efforts like Life4Pollinators and SPRING. Species-specific action plans have also been developed (e.g.^{13–15}). However, only since the adoption of the Nature Restoration Regulation

[#]A full list of authors and their affiliations appears at the end of the paper.

(2024/1991) in 2024, with its Article 10, did both EU pollinator conservation and an EU wide Pollinator monitoring become legally binding for EU Member States¹⁶.

Despite the significant progress made in Europe to tackle pollinator decline, conservation efforts are still hampered by considerable knowledge gaps¹⁷. These gaps include low knowledge of the conservation status, limited predictive abilities regarding the impact of anthropogenic and environmental stressors, lack of taxonomic capacity, and absence of standardised tools for assessing pollinator status and trends^{12,18}. The phylogenetic and morphological diversity of insect pollinators, together with their wide variation in body size, mobility, and typically short lifespans, pose substantial challenges for the development and standardisation of robust population monitoring schemes in order to fill those gaps¹⁹. Additionally, a major challenge for effective pollinator conservation is that data on pollinator distribution are notoriously scattered across numerous European institutions and online platforms (e.g., GBIF and BWARS) making comprehensive assessment and coordinated action difficult. In addition, their quality and quantity vary broadly between regions, with northern and western European countries having good datasets, while southern and eastern European countries present important knowledge gaps²⁰.

A harmonised occurrence database is a prerequisite for developing a baseline of robust indicators of pollinator status, quantifying population declines, identifying their drivers, and assessing the resulting impacts on society and the economy^{21,22}. Verified and harmonised data, alongside spatial mapping, are also essential for guiding the transition towards more targeted, sustainable, and actionable conservation policies, including prioritisation of conservation areas²³, and serve as an efficient instrument to evaluate the progress of European Union conservation and restoration strategies¹². In this context, the aim of this work is to compile and synthesise available occurrence data on wild bees and hoverflies, providing a consolidated reference database to support future research and inform the development of robust indicators for the status of these key pollinator groups in Europe.

The present work was developed within the framework of multiple EU-funded projects: (i) the “Safeguard” project, initiated by the European Commission under the Horizon-2020 program for research and innovation; (ii) the “ORBIT” and “Taxo-Fly” projects, funded by the Directorate-General for Environment (DG ENV), which aim to centralise and generate taxonomic information (including ecology and distribution) for all the species of bees and hoverflies; (iii) the “Pulse” project, also funded by DG ENV, aiming to update the Red List of European bees; (iv) The European Red List of Hoverflies project, funded by the European Commission, aimed to assess the conservation status of European hoverflies. All these projects ultimately aim to reverse the losses of wild pollinators by joining the efforts of world-leading researchers, NGOs and policy experts. Their common task is delivering the most comprehensive open EU wild pollinator distribution database, a first step in consolidating European bee and hoverfly occurrence data. The next steps should focus on integrating any overlooked datasets, considering additional data sources, including national and regional monitoring programs, unpublished datasets, and contributions from citizen science initiatives. This will help expand the records of European pollinators, improve the representativeness of different groups, and improve knowledge on their abundance and distribution across the continent. To ensure long-term relevance and usability, it is crucial to move beyond a static database and work toward developing a dynamic, continuously updated resource. Thus, following its publication, the Safeguard occurrence databases will be deposited on GBIF, where it will receive a DOI and be maintained through versioned updates. A first update is already planned within the ongoing European project WildPosh²⁴, and future updates will be released as new versions under the same DOI to ensure continuity and prevent dataset duplication. Although long-term updates are currently project-driven, due to the absence of a permanent European structure dedicated to coordinating regular dataset revisions, the steady funding of pollinator-related European initiatives, the implementation of the EU Nature Conservation Law (i.e., Chapter 10), and the monitoring obligations imposed on Member States are expected to support sustained updating of this resource.

Methods

In the following work, we consider as “data” a record that includes at least the name of the species and the coordinates where the species was observed or collected. A “dataset” refers to a set of data received from each provider, while the term “database” represents the compilation of all the datasets for each pollinator group (bees and hoverflies) into a single, consolidated resource.

All the steps of the data processing workflow are presented in Fig. 1.

Data Collection

Wild bees (Hymenoptera: Anthophila). We explored multiple sources to establish a comprehensive database on European bees. The state of the art was the existing data from BDFGM (*Banque de Données Faunique de Gembloux-Mons*) and previous European projects, which presents the baseline of our database. We then supplemented these data by acquiring extensive and comprehensive records from four main sources: taxonomists with expertise in specific taxa at global level, regional or national experts on bee fauna as well as accessible public platforms and repositories. To address gaps in existing knowledge, especially for data-deficient groups and under-sampled regions, we explored three additional sources: literature, museum and private collections, and newly collected material (Table 1).

The BDFGM is a faunistic database developed through a long-standing collaboration between the Zoology Laboratory at the University of Mons and the Entomology Unit of Functional and Evolutionary Biology at Gembloux Agro-Bio Tech, University of Liège (formerly the Faculty of Agronomy of Gembloux University), which focused mainly on spatial data on wild bees (see <http://www.atlashymenoptera.net>). The BDFGM has served as a foundational data source for several major initiatives and projects focusing on pollinator monitoring and conservation.




Fig. 1 Summary of the data processing workflow of bees and hoverflies: data collection, data unification and compilation, standardisation, cleaning (character cleaning, taxonomic cleaning, spatial cleaning, and duplicate removal), and technical validation. The numbers in brackets indicate the number of raw data at each processing stage. The numbers outside the funnels indicate the number of raw data removed (preceded by “-”) or added (preceded by “+”).

The previous projects include the Status and Trends of European Pollinators (STEP project), the Bees, Wasps and Ants Recording Society (BWARS) program, and the SAuvons les POLLinisateurs sauvages (SAPOLL) project. The STEP project was supported by the European Commission under the 7th Framework Programme for Research and Technological Development. STEP compiled a large spatio-temporal dataset on pollinators to assess the extent of their decline, identify the main drivers, and evaluate the consequences for ecosystems and agriculture². SAPOLL is a cross-border initiative funded by the European Regional Development Fund and Wallonia through the INTERREG V France-Wallonie-Vlaanderen program. It aimed to develop an action plan

datasetName	dataAvailability	categoryOfDatasetSource	datasetSource	numberOfRowData	DateOfLastUpdate
Additional_data_PULSE_SAFEGUARD	published and unpublished data	Different sources	UMONS	22754	—
Literature_Andrena	published data	Literature	see list of references in metadata_2 ⁴⁷	453	20231016
Literature_Belarus	published data	Literature	Reverté, Miličić <i>et al.</i> ²⁸ ; https://doi.org/10.1111/icad.12680	338	20230900
Literature_Ebmer_Halictus	published data	Literature	see list of references in metadata_2, Sentil <i>et al.</i> ⁴⁷	777	—
Literature_Moldova	published data	Literature	Reverté, Miličić <i>et al.</i> ²⁸ ; https://doi.org/10.1111/icad.12680	217	20230900
Literature_Russia	published data	Literature	see list of references in metadata_2 ⁴⁷	8921	20230226
Literature_Russia2	published data	Literature	see list of references in metadata_2 ⁴⁷	6320	20230901
Grandi_Collection	unpublished data	Museum/historical collection	Lucia Lenzi & Fabio Sgolastra	5392	20230331
Linz_Museum	unpublished data	Museum/historical collection	UMONS	7423	20221228
NHMW_Museum	unpublished data	Museum/historical collection	Dominique Zimmermann	3192	20221200
Priore_Collection	unpublished data	Museum/historical collection	UMONS	3158	20230828
Sapienza_Museum	unpublished data	Museum/historical collection	Maurizio Mei	6743	20230223
BoguschP_CzechRepublic	unpublished data	National expert	Petr Bogusch	6348	20220519
BurdysE_Lithuania	unpublished data	National expert	Eduardas Budrys	3215	20230131
CasalM_MelittidaeSpain	unpublished data	National expert	Mercè Galbany Casal	40	20230831
DemeterI_Romania	unpublished data	National expert	Demeter Imre	602	20190000
EIS_Netherlands	unpublished data	National expert	European Invertebrate Survey (EIS) the Netherlands, Leiden	344572	20221122
FiordalisoW_Belgium	unpublished data	National expert	William Fiordaliso	4752	20221021
FlaminioS_Italy	unpublished data	National expert	Simone Flaminio	541	20231212
GasparH_Portugal	published data	National expert	Gaspar <i>et al.</i> ²⁹ ; https://www.researchgate.net/publication/367328436_New_regional_contributions_to_the_knowledge_of_the_portuguese_bee_fauna_Hymenoptera_Anthophila	901	20221200
KieratJ_Poland	unpublished data	National expert	Justyna Kierat	40	20230828
PetanidouT_Greece	unpublished data	National expert	Theodora Petanidou	13669	20230510
RuizC_MelittidaeCanaryIslands	unpublished data	National expert	Carlos Ruiz Carreira	59	20230729
SelisM_Italy	unpublished data	National expert	Marco Selis	1942	20220930
StojnicS_Serbia	unpublished data	National expert	Sonja Mudri Stoni	4183	20220900
VarnavaA_Cyprus	published data	National expert	Varnava <i>et al.</i> ²⁹ ; https://zookeys.pensoft.net/article/38328/download/suppl/31/	458	20220410
VesnicA_Bosnia	unpublished data	National expert	Adi Vesnic	459	20230228
YvesP_Corsica	unpublished data	National expert	Gibernau Marc and Maestracci Pierre-Yves	33	20230629
BenrezkallahJ_Cyprus	unpublished data	Newly collected material	UMONS	211	20230125
deManincorN_Greece	unpublished data	Newly collected material	UMONS, de Manincor ³¹ ; https://doi.org/10.5281/zenodo.17135537	4213	20250303
DevorsineR_Italy	unpublished data	Newly collected material	UMONS	351	20230906
GambarottoM_Belgium	unpublished data	Newly collected material	UMONS	910	20230906
ThulierJ_Peyresq	unpublished data	Newly collected material	UMONS	687	20230906
Artportalen_Sweden	published data	Open data	Artportalen: https://www.artportalen.se/Occurrence/TaxonOccurrence/16/2002991	283302	20220412
ArtsDataBanken_Norway	published data	Open data	ArtsDataBanken: https://artskart.artsdatabanken.no/	117261	20220320
BiodiversityMaps_Ireland	published data	Open data	Biodiversity Maps ³² ; https://maps.biodiversityireland.ie/Dataset/5	50902	2022
BWARS	published data	Open data	BWARS (Bees, Wasps & Ants Recording Society 2022): https://bwars.com/sites/default/files/diary_downloads/20230404%20BWARS%20public%20data.zip	491891	2022
FinBIF_AmateurEntomologist_etc_Finland	published data	Open data	FinBIF: https://laji.fi/en/observation/map?target=MX.289375&collectionId=HR.22	55030	20220325
Continued					

datasetName	dataAvailability	categoryOfDatasetSource	datasetSource	numberOfRowData	DateOfLastUpdate
FinBIF_LUOMUS_Finland	published data	Open data	FinBIF: https://laji.fi/en/observation/map?target=MX.289375&countryId=ML.206&collectionId=HR.3778,HR.1989,HR.3552,HR.3551,HR.1247,HR.1627,HR.1127,HR.1207,HR.121,HR.1915,HR.3811,HR.4171	98476	20230325
GBIF_Estonia	published data	Open data	GBIF: see list of sources in metadata_3, Sentil <i>et al.</i> 2025	39480	20220706
GBIF_Switzerlands	published data	Open data	GBIF ³³ : https://doi.org/10.15468/ksfmzj	432492	2022
mdataMNHN_Luxembourg	published data	Open data	mdata.MNHN: https://mdata.mnhn.lu/	10006	20220321
Zenodo_Iberia	published data	Open data	Bartomeus <i>et al.</i> ³⁴ : https://doi.org/10.5281/zenodo.6354502	71374	20221018
Zobodat_Austria	published data	Open data	Zobodat: https://www.zobodat.at/belege.php	13251	20220802
SAPOLL	unpublished data	Previous project	UMONS	197589	20230400
BDFGM_Andrenidae	unpublished data	Previous project	UMONS	592912	20230400
BDFGM_Anthophorini_Melectini	unpublished data	Previous project	UMONS	100971	20230116
BDFGM_Bombus	unpublished data	Previous project	UMONS	754797	20231211
BDFGM_Halictidae	unpublished data	Previous project	UMONS	241252	20230400
BDFGM_Megachilidae	unpublished data	Previous project	UMONS	127526	20230400
BDFGM_Melittidae	unpublished data	Previous project	UMONS	24987	20230400
CalafatJD_BalearicIslands	unpublished data	Private data	Joan Díaz Calafat	2912	20230127
CappellariA_Italy	unpublished data	Private data	Andree Cappellari	12044	20231117
JansenK	unpublished data	Private data	Kobe Jansen	1742	2022
KleijnD_Boshommel	unpublished data	Private data	David Kleijn	8784	20221000
KleijnD_Netherlands	unpublished data	Private data	David Kleijn	1403	20221000
ScheperJ_France	unpublished data	Private data	Jeroen Scheper	4385	20221200
ULB_CanaryIslands	unpublished data	Private data	Leon Marshall	410	20220705
BoguschP_Coelioxys	unpublished data	Taxonomist	Petr Bogusch	567	20221213
BoguschP_CuckooBees	unpublished data	Taxonomist	Petr Bogusch	11164	20220307
BoguschP_Dioxyini	unpublished data	Taxonomist	Petr Bogusch	15	20221213
DatheH_Hylaeus	unpublished data	Taxonomist	Holger Dathe	29312	20211031
GadoumS_Hylaeus	unpublished data	Taxonomist	Serge Gadoum	6497	20230216
KasperekM_Anthidiini	unpublished data	Taxonomist	Max Kasperek	1036	20220611
KulmannM_Colletes	unpublished data	Taxonomist	Michael Kuhlmann	8162	20231208
MichezD_Melittidae	unpublished data	Taxonomist	UMONS	14	20230924
MullerA_Osmiini	unpublished data	Taxonomist	Andreas Muller	14150	20220412
RadchenkoV_Colletes	unpublished data	Taxonomist	Radchenko Vladimir	332	20210101
RadchenkoV_Dasydopoda	unpublished data	Taxonomist	Radchenko Vladimir	172	20210101
RischS_Eucera	unpublished data	Taxonomist	Stephan Risch	11302	20220629
SmitJ_Nomada	unpublished data	Taxonomist	Jan Smit	50903	20220400
StrakaJ_Nomada	unpublished data	Taxonomist	Jakub Straka	3211	20220418
WoodTJ_Andrena	unpublished data	Taxonomist	Thomas James Wood	15541	20220800
WoodTJ_Panurginae	unpublished data	Taxonomist	Thomas James Wood	253	20231121

Table 1. Overview of the bee datasets included in this study, with information on the dataset name, availability (published or unpublished), source category (e.g., literature, museum, national expert), specific source, number of raw data records, and date of last update.

for wild pollinators in Belgium and northern France. One of the project's objectives was to create a shared database of bees and hoverflies²⁵. BWARS is a UK-based organisation, operating under the aegis of the UK Biological Records Centre. It is dedicated to studying and conserving bees, wasps, and ants (Hymenoptera) to promote the recording and understanding of these insect groups²⁶. The datasets produced and compiled by these projects provided a range of information, including the spatio-temporal distribution of European bees.

We acquired 16 datasets from 12 taxonomists, covering most of the main bee clades recorded in Europe: Melittidae (Denis Michez and Vladimir Radchenko), Eucerini (Stephan Risch), *Nomada* (Jan Smit and Jakub Straka), *Sphecodes*, *Ammobates*, *Ammobatoides*, *Biastes*, *Chiasmognathus*, *Epeoloides*, *Parammobatodes*, *Pasites*, *Schmiedeknechtia*, *Epeolus*, *Coelioxys*, *Dioxyini* and *Paradioxys* (Petr Bogusch and Jakub Straka), Osmiini (Andreas Müller), *Andrena* and Panurginae (Thomas James Wood), *Hylaeus* (Holger H. Dathe and Serge Gadoum), Anthidiini (Max Kasperek) and *Colletes* (Michael Kuhlmann and Vladimir Radchenko). Additionally, 16 datasets from national and local experts across various European countries were obtained:

Czech Republic (Petr Bogusch), Lithuania (Eduardas Budrys), Spain (Mercè Galbany Casal), Romania (Demeter Imre), Belgium (William Fiordaliso), Italy (Marco Selis and Simone Flaminio), Portugal (Hugo Gaspar), Poland (Justyna Kierat), Greece (Theodora Petanidou), The Netherlands (Menno Reemer, European Invertebrate Survey EIS), Canary Islands (Carlos Ruiz Carreira), Serbia (Sonja Mudri Stojnić), Cyprus (Androulla Varnava), Bosnia and Herzegovina (Adi Vesnić) and Corsica (Gibernau Marc and Maestracci Pierre-Yves). In some countries, centralised national datasets have already been made publicly accessible through: (i) repositories: Zenodo (the Iberian Peninsula), (ii) National platforms: Zobodat (Austria), Artportalen (Sweden), FinBIF (Finland), ArtsDataBanken (Norway), mdata MNHN (Luxembourg), Biodiversity Maps (Ireland), and (iii) international platforms: Global Biodiversity Information Facilities GBIF (Estonia and Switzerland) (Table 1, metadata 1²⁶).

For some target genera (*Andrena* and *Halictus*) and countries (Russia, Belarus and Moldova) for which we did not have enough records from the previous sources, we digitised part of the available literature, following the recommendations of expert taxonomists (see the list of references in metadata 2²⁶).

Historical collections were also used as sources of data. We digitised information from voucher specimens housed in five European museums, including the *Oberösterreichisches Landesmuseum* (Linz, Austria), the Sapienza University of Rome Museum of Zoology (Rome, Italy), the Natural History Museum of Vienna NHMW (Vienna, Austria), the Priore historical collection hosted by the University of Naples Federico II (Naples, Italy), and the Guido Grandi Collection at the University of Bologna (Italy). Data from many museums had already been included in the STEP dataset as part of taxonomic revisions (e.g.²⁷). The selection of museums was guided by the taxonomic expertise of the co-authors on the diversity and originality of the collections (e.g., number of species and rarity), the accessibility of the material, as some European museums offer limited access, and the accuracy of specimen identifications (i.e., collections that had recently undergone taxonomic revision). Considering limited time and resources, material from data deficient species or less studied regions was prioritised for encoding.

Lastly, we targeted some Southern European countries, particularly Italy, Greece and Cyprus, where significant gaps in occurrence data were identified²⁸. Field trips were organized in collaboration with taxonomists to address these gaps and collect new material from these under-sampled areas. This category (i.e., newly collected material) also includes two datasets from Belgium and France that were collected by the UMONS team throughout the Safeguard project period.

Through the exchanges we had with experts, we gained access to seven additional private datasets (Table 1).

We considered that enough data had been collected for a species when taxonomists, national and regional experts, confirmed during maps' validation that the available records represented the distribution of the target species.

Hoverflies. Regarding hoverflies, two projects have significantly contributed to compiling spatial data. The first is the European Red List of Hoverflies (EU and pan-Europe), the first assessment of the conservation status of European hoverflies. This project gathered and validated a large amount of spatial data and also highlighted gaps and geographical biases in the data. The second is the EU funded Service Contract 'Taxonomic Resources for EU Pollinator Monitoring Schemes (2 Lots), Lot 2 - Development of Taxonomic Tools for Hoverflies (Insecta, Syrphidae) (Taxo-Fly)', which focused on compiling accurate taxonomic, occurrence and ecological data on European hoverflies. Furthermore, the STEP project provided a certain quantity of data related to hoverflies.

Similar to the approach used for bees, the core hoverfly database was expanded by integrating extensive and detailed records obtained from taxonomists specialising in particular hoverfly taxa, experts with regional or national knowledge of hoverfly fauna, publicly accessible platforms and biodiversity repositories, and institutional databases. We incorporated additional data from published literature, entomological collections housed in museums and private holdings, and recently gathered field material.

IUCN Red List Assessors compiled part of the data included in this work: Andrea Aracil, Ana Grković, Antonio Ricarte, Axel Ssymank, Anja Šebić, Ante Vujić, Celeste Pérez Bañón, Dubravka Milić, Gerard Pennards, Gunilla Ståhls, Jeroen van Steenis, Laura Likov, Leendert-Jan van der Ent, Libor Mazánek, Marija Miličić, Marina Janković, Rita Földesi, Santos Rojo, Sanja Veselić, Snežana Radenković, Tamara Tot, Zorica Nedeljković and Wouter van Steenis. Some of these data records were already included in the previous projects (e.g., STEP), but many records were newly collected in the framework of this project.

Historical collections were also valuable sources of data for Syrphidae. Data from several museums visited over the years (visits undertaken in the context of other research objectives) have been digitised and deposited in the FSUNS database (see the list of museums and historical collections in metadata 2²⁶).

Regarding field trips, research efforts focused on areas known for their high species richness (e.g., Greece and Spain), as these regions are believed to host species that are either new to science or to Europe.

The nature of the contributed data varied considerably across sources. The IUCN assessment relied on datasets compiled by assessors from published literature and their own private databases, while Taxo-Fly brought together additional occurrence information from a wide range of sources and systematised these records for taxonomic use. Earlier material generated during the STEP project also contributed to the dataset. These project-based datasets were complemented by expert-contributed files, publicly accessible biodiversity platforms such as GBIF and FinBIF, digitised material from historical museum collections, and newly gathered field records from regions of high species richness. Because these sources differ in scope and content, their proportional contributions are not summarised here; the origin of each individual record is preserved in the database and can be directly identified by users. The present project integrates all of these inputs into a single harmonised database^{29–34}.

Data unification and compilation

A thorough exploration of the datasets was conducted to gain a general insight into their contents and formats before the standardisation and cleaning process. Based on this, we developed a data template, which defines the structure of our databases by specifying field names (i.e., column names) and their expected values or formats. To ensure compatibility with existing standards referenced in Simon Delso, *et al.*³⁵ and to adhere to FAIR

(Findable, Accessible, Interoperable, and Reusable) principles³⁶, we developed the “Pollinator Core” template. This template aligns with Darwin Core where applicable. However, Darwin Core is a general-purpose biodiversity standard and does not cover all the data elements required for pollinator monitoring, such as *data-setProvider*, *datasetSource*, *dataSource*, *individualCountInterval* and *startIndividualCount*, *endIndividualCount*. The standards compiled in the Pollinator Metadata Standard³⁵, extend Darwin Core with additional terms. Our “Pollinator Core” follows this approach, incorporating Darwin Core and Pollinator Metadata Standard terms and additional fields needed to capture the full scope of our database. Table 2 presents the fields of the template. The file metadata 3²⁶ describes the template fields and provides a mapping between their names and the existing standards.

The contents of the bee datasets were placed separately in the standardised template, and then merged into one unique database.

For hoverflies, the initial step involved merging all datasets to retain relevant information into FileMaker Pro³⁷, a cross-platform relational database application. It integrates a database engine with a graphical user interface (GUI) and security features, allowing users to visually modify a database. It supports the use of sub-databases, which are essentially related tables within a main database, enabling users to structure and organise their data efficiently. These sub-databases are linked through relationships, allowing for easy access to related information. Additionally, FileMaker Pro incorporates calculations (for calculation details see metadata 4²⁶), which can be used to automate tasks, perform mathematical operations, or manipulate data dynamically within fields, layouts, and scripts. A dedicated database was developed to integrate all separate datasets (including complete information provided from each source). Most data wrangling was carried out within this platform. Four sub-databases were created - covering dates, specimen sex, species names, and presence in Europe based on Reverté, Miličić, *et al.*²⁸ - as detailed later in the text. In certain instances, the original information was retained as is and incorporated into the final database (e.g., nearest place and locality, included in *verbatimLocality*). However, the majority of variables underwent transformation, involving multiple calculations to standardise and clean the data. Only data that met all predefined criteria—such as valid species names, valid dates, valid geographic coordinates, and confirmed presence in Europe according to Reverté, Miličić, *et al.*²⁸—were included in the final database presented in this study. Clean hoverfly datasets contained following fields: Collection, Code, Genus, Species, Genus_Species, Country, Nearest place, Locality, Latitude, Longitude, Collector, Date_Start_original, Date_End_original, Starting_day, Starting_month, Starting_year, Ending_day, Ending_month, Ending_year, Source, Data_provider, without_year (Boolean, true/false), which were then mapped to Pollinator Core template, same as for wild bees.

Because the two databases were prepared independently, each team relied on the tools and workflows they had already established, which explains the differences in software used. Nevertheless, the objectives and the final outputs were fully harmonized.

Data Standardisation

Coordinates. *Georeferencing.* We collected data both with and without geographical coordinates. When records lacked original coordinates, but included locality descriptions that clearly referred to a single identifiable place (e.g. city name), we inferred the coordinates by locating and verifying the site using standard digital mapping resources. Information on whether coordinates were provided originally or added manually is available in the “manualGeoreferencing” field in both databases.

For bees, localities were classified into two categories: those with clear spatial extents and those with undefined boundaries. We considered a locality to have undefined boundaries when its limits are not mapped (e.g. regions). We determined the boundaries for localities with well-defined spatial extents by drawing an imaginary polygon encompassing the locality. For localities with undefined boundaries, we estimated the boundaries by identifying and including features of similar type and size (i.e., localities that share the same characteristics, such as forests or protected areas). Once the boundaries were established, we obtained the coordinates of the centroid of the polygon.

For hoverflies, two different approaches were used when coordinates were unavailable. For well-documented localities that have been consistently studied over the years, we assigned coordinates from the FSUNS database, an institutional repository compiling locality information from previous research projects, expert surveys, and published sources, selecting those known to be accurate and reliably associated with the respective locality. These coordinates correspond to previously verified records bearing the same locality name. For localities that had not been previously investigated, we used the locality name as a reference and identified the nearest ecologically meaningful site, such as a park, forest edge, or other natural or semi-natural area. The coordinates of this nearby area were then assigned to the record.

Google Maps³⁸ was used as a source of coordinates for both bees and hoverflies. All coordinates were recorded using the WGS 1984 geographic coordinate system.

Considering the large number of non-georeferenced records, we prioritized georeferencing for data-deficient groups and poorly studied regions. The remaining non-georeferenced records were excluded but will be explored in future projects.

Coordinates’ conversion. The original geographic coordinates were presented in various formats: decimal degrees (DD), degrees minutes and seconds (DMS), and degrees decimal minutes (DDM). First, we identified the format of each set of coordinates. We considered the DD format in our data due to its widespread usage, its scientific format and global applicability. After identifying the formats, we automatically converted the other coordinate formats to DD in Excel or R using custom functions and the conversion formula ($DD = DD + MM/60 + SS/3600$). For hoverflies, the conversion was done with the help of a calculation field for the coordinate conversion within FileMaker Pro ($DD = DD + MM/60 + SS/3600$).

Field	Description
databaseName	Name of the database that includes data on bees or hoverflies.
institutionName	Name of the institution in charge of data management.
projectSource	Name of the projects that have funded the curation process.
projectUpdate	Year of the last database update.
datasetName	Name of the dataset.
datasetProvider	The entity or person who shared or made available the dataset for this project.
datasetSource	Entity or person who originally produced or generated the dataset.
dataSource	Entity or person who produced the data.
license	Creative Common License provides information concerning official permission to use the database.
occurrenceID	A new unique identifier number that was assigned to each data.
verbatimOccurrenceID	The original unique identifier of the data.
scientificNameAuthorship	Author credited for the description of the species (scientificName) and the year of publication.
order	Name of the order to which the species belongs.
family	Name of the family to which the species belongs.
subfamily	Name of the subfamily to which the species belongs.
tribe	Name of the tribe to which the bee species belongs. For hoverflies, we did not use tribes since it is currently undergoing substantial changes.
genus	Name of the genus to which the species belongs.
subgenus	Name of the subgenus to which the species belongs.
infraspecificEpithet	Scientific name of the infraspecific epithet, which corresponds to the third part of the scientific name, is used to classify a rank below speci
scientificName	Full binomial name of a species, consisting of a combination of the genus name and the specific epithet.
individualCountInterval	Indicates if an individual count interval occurred (between startIndividualCount and endIndividualCount). If the interval occurred is TRUE other
startIndividualCount	Minimum number of individuals recorded, used when a range is provided to indicate uncertainty in the count.
endIndividualCount	Maximum number of individuals recorded. If no range is specified, endIndividualCount is used as the default.
sex	Sex of the individual(s) represented in the data.
eventDateInterval	Indicates if a sampling interval occurred (between startEventDate and endEventDate). If the interval occurred is TRUE otherwise FALSE.
startEventDate	Starting date of the interval during which the record was done, used when a range is specified to indicate the earliest possible date of the eve
endEventDate	Ending date of the interval during which the record was done. If no range is specified, endEventDate is used by default.
startDay	Earliest day of the interval during which the record was done, used when a range is specified.
startMonth	Earliest month of the interval during which the record was done, used when a range is specified.
startYear	Earliest year of the interval during which the record was done, used when a range is specified.
endDay	Latest day of the interval during which the record was done. If no range is specified, endDay is used by default.
endMonth	Latest month of the interval during which the record was done. If no range is specified, endMonth is used by default.
endYear	Latest year of the interval during which the record was done. If no range is specified, endYear is used by default.
country	Name of the country where the record was observed or collected.
stateProvince	Name of the next smaller administrative region than the country (state, province, canton, department, region, etc.) in which the event occurs.
county	Name of the next smaller administrative region than stateProvince (county, shire, department, etc.) in which the event occurs.
verbatimLocality	Original textual description of the place.
decimalLatitude	Geographic latitude (in decimal degrees) of the data, based on the WGS 1984 geographic coordinate reference system. Positive values are north of
decimalLongitude	Geographic longitude (in decimal degrees) of the data, based on the WGS 1984 geographic coordinate reference system. Positive values are east of
coordinateUncertaintyInMeters	The maximum horizontal distance (in meters) representing the uncertainty around the provided coordinates.
manualGeoreferencing	Indicates whether the coordinates in the bee and hoverfly databases were originally provided in the datasets or manually georeferenced by our te
isPseudodata	Indicates records for which coordinates are not directly tied to individual specimens (see "Pseudodata" section for more details). If a record i
verbatimElevation	The original elevation (altitude, usually above sea level) of the verbatimLocality.

Table 2. Description of the fields included in the template of the bee and hoverfly databases.

Sex. Since the sex of the specimens was recorded in various formats (e.g., male, M, ♂, or in languages other than English), we standardised the information by reducing it to three main characters: M for males, F for females and M&F for both (records that includes both males and females). In the bee database, additional characters were: G (gynandromorph: specimens that have externally features of both sexes); Q (queen) and W (worker). For the hoverfly database a sub-database was created in FileMaker Pro to standardise the information. If the sex was unknown, the data record was marked as “NA”.

Number of specimens. For bees, we extracted all non-Arabic numerical values, including ranges, textual descriptions, and Roman numerals. Using this extraction of unique values, we created a dictionary that mapped the original values to their corresponding Arabic numerical values. For data presented as intervals, we split the information into two separate columns: one for the minimum value and one for the maximum value. If only one value was provided, it was recorded in the maximum value column. We then integrated the new numerical values using the created dictionary and the `left_join` function from the ‘dplyr’ package³⁹. The invalid values were replaced with 1. A similar procedure was applied to the hoverflies within FileMaker Pro, with the added step of extracting information from the sex column. Data providers sometimes reported sex and specimen counts together (e.g., “32 males, 12 females”), which required additional processing. These numbers were extracted and added to the “number of specimens” column. If only one sex and its count were reported, that number was used directly. When counts for both sexes were provided, and the exact numbers of males and females were known, the total number of specimens was calculated as their sum.

Geographical assignment. We assigned the current country, stateProvince and county name to all data using the function `st_intersection` from the package ‘sf’^{40,41}. This function assigns administrative divisions by overlaying the geographic coordinates of each occurrence with shapefiles that define the corresponding administrative boundaries.

Coordinate uncertainty. Similar to the number of specimens, the non-Arabic numerical values of precision were extracted. We created a dictionary to map the original precision values to their corresponding new Arabic numerical values (in meters). For precision data presented as intervals, we split the range into two separate columns: one for the minimum value and one for the maximum value. If only one value was provided, it was recorded in the maximum value column. We then integrated the new precision values using the dictionary and the `left_join` function from the ‘dplyr’ package³⁹. The original precision field was subsequently excluded from the database, and unreasonable values (e.g., negative numbers, extremely large distances) were replaced with “NA”. For hoverflies, no information on precision was available.

Date. For bees, dates were recovered when accurate date events were provided. Illogical or invalid dates that fell outside acceptable ranges, such as a year after the present year, a day greater than 31, or a month greater than 12, were omitted from the database. Various date formats (e.g., YYYY/MM/DD, YYYY-MM-DD, YY-MM-DD, DDMMYYYY, DD/MM/YYYY, DD-MM-YYYY, DD-MM-YY, DD.MM.YYYY, YYYY-MM, YY-MM, YYYY/MM) were standardised to the YYYY-MM-DD, in accordance with the ISO 8601–1:2019 standard. Due to the substantial differences in date formats, we conducted date inspection and cleaning automatically whenever possible, and manually in cases where automation was not applicable. When dates were displayed as intervals (e.g., “2024/07/25–29”), we split these intervals into two fields: the starting date and the ending date. If only one date was provided, the date was duplicated and the presence/absence of an interval is indicated in the `eventDateInterval` field.

January 1st and December 31st are sometimes used as placeholders for `startEventDate` and `endEventDate` within the same record when only the year of collection is known, which may affect phenology studies. To address this issue, we replaced the day and month with “NA” for records with YYYY-01-01 and YYYY-12-31 as a start date and end date (7463 records), respectively.

Cleaning

Characters cleaning. Initially, we standardised all columns to the Latin-1 format when importing the datasets. We then replaced special characters, including “?” and various unicode whitespace characters, that could have caused issues during database importation in R. Subsequently, we cleaned the data using the `mutate()` and `gsub()` functions from the ‘dplyr’ package in R³⁹.

Taxonomic cleaning. Species name validation was done in two steps: first, we created the taxonomic backbone with the most up-to-date bee and hoverfly taxonomy, and then we proceeded with the taxonomic cleaning.

For bees, we used the most recent European checklist, which counts 2,138 species⁴². We used the EU checklist to replace the junior synonyms with the accepted senior synonyms. We then prepared a second list (hereafter called “wild bee taxonomic dictionary”) of species names where we compiled all other types of mistaken names, such as names with typos, wrong spaces, orthographic variants, other misspellings, and old synonyms. This wild bee taxonomic dictionary has been reviewed and corrected by expert taxonomists in the coauthorship list to make sure it has the most updated species names. Finally, the two lists, the European checklist and the wild bee taxonomic dictionary, have been merged into one final list.

The bee taxonomic cleaning was done in R using a two-match process: (i) first, we matched and replaced all invalid species names (e.g. misspellings, synonyms) that were present in the bee database, with their valid scientific names present in the final list (i.e., the EU checklist of wild bees and the wild bee taxonomic dictionary); (ii) secondly, species names that were not verified and replaced during the first match, have been checked using a fuzzy search approach. This approach used the European checklist as a backbone and matched the uncorrected




Fig. 2 Map of Europe showing the geographical framework of the compiled database of bees and hoverflies (based on Ghisbain, Rosa, *et al.*⁴² and Reverté, Miličić, *et al.*²⁸).

species names to it using the `stringdist_join` function in the R package ‘fuzzyjoin’^{43,44}, that assigned a likelihood score using the Jaro-Winkler distance method. This helped us to manually correct the most likely names, i.e. species names that had a likelihood score below 0.2. Records with wrong or incomplete names (e.g., resolved at the Genus or Family level) have been deleted from the aggregated database (71,517 data).

For hoverflies, we used the species list published in Reverté, Miličić, *et al.*²⁸, which counts 913 species. Similar to the bee database, a taxonomic dictionary was created for hoverfly names, addressing misspellings, synonyms, blank spaces, and other issues (referred to as the ‘hoverfly taxonomic dictionary’). This hoverfly taxonomic dictionary was used to clean and standardise species names and replace them with valid hoverfly nomenclature. Entries with incorrect or incomplete species names were removed (30,337 data). The taxonomic cleaning of the hoverfly database was conducted in FileMakerPro.

These dictionaries are “static” in time and do not take into account the variable use of names over time, and also the variable use of names by region. Because names are sometimes used in a *sensu auctorum* (i.e. in a way that is not consistent with the type material), a straightforward taxonomic dictionary will not capture uses that deviates from a chresonomic framework. In these cases, consultation with taxon authorities is required to ensure data quality. For example, the use of “*Andrena trimmerana*” from 1802–1917 is different from its use after 1917, and between 1917–2022 the use of this name in the published literature and in specimen datasets has also been variable and inconsistent. However, when possible and necessary, these changes in names have been taken into account during the data validation process (see paragraph “Technical validation” below) and ambiguous records have been either deleted or have been assigned to another taxon during the distribution map revision with experts.

Nevertheless, for the synonyms cleaning, we strongly encourage referring to the most up-to-date taxonomic literature. These dictionaries are still useful in the taxonomic cleaning process when manipulating large amounts of records, particularly in relation to the correction of species names typos, misspelling and incomplete names that could not have been cleaned otherwise.




Fig. 3 Map of Europe showing the spatial distribution of data for (a) bees and (b) hoverflies at a 10 km grid cell.

Spatial cleaning. The geographical range considered for the spatial cleaning is based on the IUCN Red List for geographical Europe and for the 27 Member States of the European Union (EU27) (detailed discussion on the European borders is given in Rasmont, *et al.*⁴⁵ (Fig. 2). This geographic range included continental Europe, extending from Iceland and the Azores Islands in the west to the Urals in the east, and from Franz Josef Land in the north to the Canary Islands in the south. The United Kingdom is also included at the geographical Europe level. The Caucasus region is not included, along with the Russian Northern Caucasus. The European part of Türkiye is included, but not the Asian part.

Records with spatial issues were deleted from the final database using the following conditions: (i) data without coordinates, (ii) data with incomplete coordinates, (iii) data with coordinates that fall outside the IUCN

shape file described above for Europe, which included non-European data and data that fall into the water (e.g., sea or lakes), and (iv) only for hoverflies, data for which presence was not reported in the country records list²⁸.

After the spatial cleaning, 296,310 records for bees and 484 records for hoverflies have been removed.

Duplicate removal. One of the first steps in data cleaning for hoverflies was removing duplicates. Duplicates came from overlapping sources, for example, parts of the FSUNS hoverfly database had been used in previous works, including IUCN assessments, and additional records were retrieved from external repositories such as GBIF, where some contributors had already deposited their data.

When two data in the initial datasets had exactly the same locality name, coordinates, date, combination of sex and number of specimens, they were regarded as duplicates and one of them was removed from the final database. The cleaning process was done using Statistica for Windows v. 13⁴⁶. After the duplicate removal, 429,606 data were removed.

Duplicate removal was not performed for the bee database because most of the provided data lacks a unique identifier for tracking. The encoding of different specimens within the same dataset varies: in some cases, multiple rows may contain identical information for each field, while in others, all data is merged into a single row, with the number of specimens recorded. Unlike the hoverfly database, it was not possible to establish consistent association across all fields (e.g., location, coordinate, date) without the risk of removing records that were not true duplicates. Due to data wrangling and simplification made by data providers in their original datasets, an exact match between records was not achievable, preventing reliable duplicate identification. We acknowledge that this may pose a limitation; however, we have accounted for this issue throughout the data validation and cleaning process (see below). Nevertheless, this issue is limited to a few datasets and only affects certain species groups. Therefore, we consider the bee database to remain reliable.

Data Records

The final cleaned databases and the additional material used during the data collection and cleaning processes are available for download from Zenodo, a free and open-access research repository. For simplicity, two repositories have been created, and the material is organised as follows:

1. Spatio-temporal distribution of bees and hoverflies (Zenodo⁴⁷). It includes:
 - (i) Two occurrence databases of bees and hoverflies in “Pollinator Core” template (as presented in section 2. Data unification and compilation), saved as RDS and CSV files.
 - (ii) Metadata files:
 - Metadata 1: information about the datasets for bees.
 - Metadata 2: a list of explored museums and private collections, and the references reviewed to acquire occurrence data.
 - Metadata 3: a detailed description of the fields for both databases.
 - Metadata 4: a detailed description of the calculations used in FileMakerPro to prepare and clean the hoverfly database.
 - Metadata 5: the data used to produce Figs. 4 and 5.
 - Metadata 6: the list of the dataset sources of the Estonian data (GBIF_Estonia dataset) retrieved from GBIF.
2. Taxonomic dictionaries and cleaning scripts for wild bees and hoverflies (Zenodo⁴⁷). This Zenodo repository provides a zip folder for the GitHub repository implemented in Zenodo, including:
 - (i) a README file with information concerning all the files and scripts provided in the repository and how to use them;
 - (ii) a “scripts” folder with the scripts used to prepare, check and clean the databases;
 - (iii) a “data” folder containing the files used for taxonomic cleaning for both bees and hoverflies, including the *WildBeeChecklist*, the *WildBeeDictionary*, the *HoverflyDictionary* and the *pollinatorClassification*. The *WildBeeChecklist* is the list of species names reported in Ghisbain, Rosa *et al.* 2023. The two taxonomic dictionaries contain the list of species names used for taxonomic cleaning, and include three columns: the first column contains the original names with errors (e.g., misspelling, typos, etc.), the second column lists the correct names, and the third column indicates the presence of each species in Europe based on the European checklists (for bees Ghisbain, Rosa, *et al.*⁴², for hoverflies Reverté, Miličić, *et al.*²⁸). The *pollinatorClassification* file is the taxonomic backbones used to associate all relative taxonomic information, such as Order, Family, Authorship, etc.

In addition to these files, there are three additional folders:

- (a) in the first folder “shapefile”, we provided the shape files to associate the administrative level (country, stateProvince and county);
- (b) in the second folder “working_directory”, we provided an example of a dataset that can be used to run the scripts;
- (c) in the third folder “ToFill”, we provided a detailed description on how to collect and organise the data using the same standard template provided in this manuscript (“ToFill_FieldDescription.xlsx”), and an empty template ready to use (“ToFill_fill_Me.xlsx” or “ToFill.csv”). A README file (“ToFill_ReadMe.csv”) is also provided.




Fig. 4 Map of Europe showing the proportion of (a) bee and (b) hoverfly species covered by our database compared to the national checklists of European bees and hoverflies²⁸. Colour intensity indicates the proportion of species covered by our database relative to the total number of species reported per country. The data used to produce this figure are available in metadata 5 in Sentil, *et al.*⁴⁷.

Data Overview

This work resulted in two comprehensive databases counting 4,341,684 and 1,036,181 data for bees and hoverflies, respectively. Overall, the databases covered 97% of the European bee species (2,083 species out of 2,138) and 97% of the European hoverfly species (886 species out of 913) (see Fig. 3). The countries with the highest number of bee species in our database are Spain (1,050), Greece (1,024), Italy (907) and France (902) (Fig. 4a) and countries with the highest number of hoverfly species are France (546), Serbia (443), Germany




Fig. 5 Visual representation of the database taxonomic composition for **(a)** wild bees and **(b)** hoverflies. Each panel combines a donut chart and a pie chart. The donut chart illustrates the proportion of species present in Europe, while the pie chart shows the proportion of records in the database, grouped by taxonomic rank (tribes for bees, subfamilies for hoverflies). Taxonomic groups with fewer than 60,000 records per family were grouped under “Others.” Numbers in the pie charts refer to the following taxonomic groups: 1 = Melittidae, 2 = Other Megachilidae, 3 = Sphecodini, 4 = Other Halictidae, 5 = Other Andrenidae. The data underlying this figure are available in Reverté, Miličić, *et al.*²⁸ and metadata 5 in Sentil, *et al.*⁴⁷. Bee photographic material provided by the Beelibre team (beelibre.lu), and selected by Antonio Cruz and Fernanda Herrera-Mesías. Hoverfly photographic material provided by Sander Bot.

(426) and Italy (414) (Fig. 4b). The bee families with the highest number of records in our database (Fig. 5a) are Apidae (1,837,090), Andrenidae (1,078,322), and Halictidae (732,843), followed by Megachilidae (423,336), Colletidae (197,751), and Melittidae (72,342). The hoverfly subfamily with the highest number of records (Fig. 5b) is Eristalinae (645,502), followed by Syrphinae (375,541), Pipizinae (13,647) and Microdontinae (1,491).

Technical Validation

The validation of the spatial distribution was conducted by taxonomists and national experts. During this process, doubtful records not confirmed by experts were removed from the database.

For bees, the technical validation process followed these steps: (i) all data with spatial coordinates were split into different groups based on the Family (e.g., Halictidae) or Tribe (e.g., Anthidiini), (ii) for each species within each group a static map was made, with an associated Excel file where the information of the records was provided (e.g., taxonomic information, spatial and temporal information, etc.) and where a validation column was added for the following steps, (iii) each group has been assigned to one or more experts to review the maps and to validate or flag the incorrect points, (iv) if the distribution of the species was incomplete, new overlooked records were provided by experts and included to fill the gaps on the maps, (v) based on the validation column (filled with “OK” for valid records or “DELETE” for flagged records or “ADD” for additional records) a new version of the maps and the Excel files were made for each species, in which the incorrectly flagged points were removed. Steps from (iii) to (v) were conceived as an iterative process until all experts approved the final version of the maps. Only validated and additional records were retained and merged into a final database. At the end of this process, 775,590 records were removed and 886,684 were added (Fig. 1).

For hoverflies, technical validation was carried out by Ante Vujic for all species through visual inspection of maps created in QGIS version 3.22.4⁴⁹. These maps included associated files with geographic coordinates, and any doubtful data points were removed. For certain species, additional experts familiar with the regions from which the questionable data originated were consulted to verify the issues. In total, 53 records were removed during this process.

Databases limitations. Despite the great effort that has been done to collect and validate the occurrence records in the bee database, it presents some limitations due to the presence of what can be termed “pseudodata” concerning the genus *Andrena*. Specifically, “pseudodata” records originate from an unpublished *Andrena* monograph of Klaus Warncke which included maps with distribution points added by hand; these maps were later published by Gusenleitner & Schwarz⁵⁰. These distribution maps were scanned and digitally transformed in order to generate “occurrence points” on a map (forming a grid pattern easily recognisable when plotting the points), and to create machine-readable records in a “specimen” database. Therefore, these records are not directly tied to individual specimens or collection dates (all records refer to the year 1989, which is the year that this monograph was last worked on), and hence could misrepresent the true size of the dataset in terms of occurrence records. However, these records are useful for range delineation (e.g., creation of minimum convex polygons) and have been then included in the database to calculate the maximum area extent in the updated Red List of European Bees.

Pseudodata records were identified by filtering the *Andrena* records from the BDFGM dataset, which contain no collection dates other than the general publication year of 1989. Subsequently, using QGIS mapping software, records that did not align with the expected grid pattern were manually removed to avoid excluding valid data. Records that are aligned with the grid patterns have been removed. For this purpose and to help data users to easily locate these records, a new column named “isPseudodata” has been added to the bee database, with the value TRUE indicating the presence of pseudodata records.

The Syrphidae database is a compilation of data from multiple sources. While it includes most known species, there are significant disparities in coverage between different species and countries. For the rarest species, the majority of existing records have been incorporated, largely thanks to efforts related to the European Red List. In certain genera—particularly *Brachyopa* and, to a lesser extent, *Merodon*—the database captures a substantial proportion of all known records. However, for most other genera, the records are more fragmented and inconsistently represented. Similarly, geographic coverage varies. Some countries, such as Serbia, Greece, and other parts of the Balkans, are well represented in the database. Others, like the Netherlands and the United Kingdom, contain large numbers of records; however, these are predominantly recent observations of photographed flies submitted through platforms like Observation.org and iNaturalist. This results in a high number of records for large, easily identifiable species, while older records and sightings of smaller or more cryptic species are underrepresented. Meanwhile, many countries still have only a limited number of records. In France, for instance, the INPN hosts around 160,000 hoverfly records, but the absence of a centralized national database means that many datasets from institutions and private collections remain unincorporated. This lack of coordination continues to limit access to existing information. Efforts to address these gaps are ongoing. In Italy, for example, a project launched in 2020 is monitoring pollinators—bees, hoverflies, and butterflies—across all 24 National Parks. Initial results are now being published, contributing with important new data and demonstrating the value of coordinated national initiatives in improving our understanding of hoverfly distributions.

Data availability

The bee and hoverfly databases, with their associated metadata, are available at <https://doi.org/10.5281/zenodo.17107215> (Zenodo repository). The taxonomic dictionaries and cleaning scripts for wild bees and hoverflies are available at <https://doi.org/10.5281/zenodo.16412138> (Zenodo repository).

Code availability

The code used for the analyses in this study is available on GitHub at [<https://github.com/Safeguarding-European-wild-pollinators/SafeguardEuropeanWildBee>], implemented in the Zenodo repository⁴⁸. This repository includes all scripts and documentation necessary to reproduce the results presented in this paper. The analyses conducted in this study used several R packages for data wrangling, visualisation, and spatial analysis. Key packages included: 'dplyr'³⁹ (v1.1.4): for data wrangling and transformation; 'sf'^{40,41} (v1.0-16): for handling spatial data; 'ggplot2'⁵¹ (v3.5.0): for data visualisation; 'lubridate'⁵² (v1.9.3): for working with date-time data; and 'tidyr'⁵³ (v1.3.1): for tidying and reshaping data. Refer to the Github repository for a complete list of packages and their respective versions used in the analysis.

Received: 22 September 2025; Accepted: 14 January 2026;

Published online: 30 January 2026

References

- Oliver, T. H. *et al.* Declining resilience of ecosystem functions under biodiversity loss. *Nat. Commun.* **6**, 10122, <https://doi.org/10.1038/ncomms10122> (2015).
- Potts, S. G. *et al.* Safeguarding pollinators and their values to human well-being. *Nature* **540**, 220–229, <https://doi.org/10.1038/nature20588> (2016).
- Nieto, A. *et al.* *European Red List of Bees* (Luxembourg: Publication Office of the European Union, Belgium, 2014).
- Vujić, A. *et al.* *Pollinators On The Edge: Our European Hoverflies. The European Red List of Hoverflies* (Brussels, Belgium: European Commission <https://doi.org/10.2779/359875>, 2022).
- Drossart, M. *et al.* *Belgian Red List of bees*. (Mons: Presse universitaire de l'Université de Mons, 2019).
- Zeegers, T., van Steenis, W., Reemer, M. & Smit, J. Drastic acceleration of the extinction rate of hoverflies (Diptera: Syrphidae) in the Netherlands in recent decades, contrary to wild bees (Hymenoptera: Anthophila). *J. van Syrphidae* **3**, 1–11, <https://doi.org/10.55710/1/YDSJ1547> (2024).
- Barendregt, A., Zeegers, T., van Steenis, W. & Jongejans, E. Forest hoverfly community collapse: Abundance and species richness drop over four decades. *Insect Conserv. Divers.* **15**, 510–521, <https://doi.org/10.1111/icad.12577> (2022).
- Hallmann, C. A., Ssymank, A., Sorg, M., de Kroon, H. & Jongejans, E. Insect biomass decline scaled to species diversity: General patterns derived from a hoverfly community. *Proc. Natl. Acad. Sci. USA* **118**, e2002554117, <https://doi.org/10.1073/PNAS.2002554117> (2021).
- Dicks, L. V. *et al.* A global-scale expert assessment of drivers and risks associated with pollinator decline. *Nat. Ecol. Evol.* **5**, 1453–1461, <https://doi.org/10.1038/s41559-021-01534-9> (2021).
- Gekière, A., Gérard, M., Potts, S. G., Michez, D. & Ghisbain, G. Underlying mechanisms shaping wild bee decline. *Biol. J. Linn. Soc.* **145**, blaf043, <https://doi.org/10.1093/biolinnean/blaf043> (2025).
- Wan, N. F. *et al.* Pesticides have negative effects on non-target organisms. *Nat. Commun.* **16**, 1360, <https://doi.org/10.1038/s41467-025-56732-x> (2025).
- European Commission. *Revision of the EU Pollinators Initiative: A New Deal for Pollinators*. Brussels, Belgium. Official Journal of the European Union, (2023).
- Monasterio, Y. L. *et al.* *Canarian Islands endemic pollinators of the Laurel Forest zone—Conservation plan 2023–2028*. (2023).
- Michez, D. *et al.* *Teasel-plant specialised bees in Europe – Conservation action plan 2023–2030* (2023).
- Vujić, A. *et al.* *Hoverflies specialised to veteran trees in Europe—Conservation action plan 2023–2030* (2023).
- European Commission. *Regulation (EU) 2024/1991 on nature restoration and amending Regulation (EU) 2022/869* (Official Journal of the European Union, 2024).
- European Commission. *Opportunities and knowledge gaps for EU research and innovation on pollinators* (Publications Office of the European Union, 2023).
- European Court of Auditors. *Special Report 15/2020: Protection of wild pollinators in the EU – Commission initiatives have not borne fruit* (Publications Office of the European Union, 2020).
- Aubouin, L. *et al.* BeeFunc, a comprehensive trait database for French bees. *Sci. Data* **12**, 1302, <https://doi.org/10.1038/s41597-025-05626-0> (2025).
- Marshall, L. *et al.* Understanding and addressing shortfalls in European wild bee data. *Biol. Conserv.* **290**, 110455, <https://doi.org/10.1016/j.biocon.2024.110455> (2024).
- Garibaldi, L. A., Bartomeus, I., Sáez, A., Aizen, M. A. & Fijen, T. Crop pollination management needs flower-visitor monitoring and target values. *J. Appl. Ecol.* **57**, 664–670, <https://doi.org/10.1111/1365-2664.13574> (2020).
- IPBES (Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services). *Summary for policymakers of the assessment report of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services on pollinators, pollination and food production*. Potts, S. G. *et al.* (eds). (IPBES Secretariat, Bonn, Germany, 2016).
- Nania, D. *et al.* Insects as indicators of key biodiversity areas. *Insect Conserv. Divers.* **17**, 464–473, <https://doi.org/10.1111/icad.12712> (2024).
- Michez, D. *et al.* WildPosh: Pan-European assessment, monitoring, and mitigation of chemical stressors on the health of wild pollinators. *Research Ideas and Outcomes* **11**, e156185 (2025).
- Folschweiller, M. *et al.* *Plan d'action transfrontalier en faveur des pollinisateurs sauvages*. In *Projet Interreg V SAPOLL – Sauvons nos Pollinisateurs – Samenwerken voor Pollinators* (Université de Mons, Mons, Belgium, 2019).
- BWARS (Bees, Wasps & Ants Recording Society) <https://www.bwars.com/index.php?q=bee/apidae/anthophora-plumipes> (2022).
- Michez, D. & Eardley, C. Monographic revision of the bee genus *Melitta* Kirby 1802 (Hymenoptera: Apoidea: Melittidae). *Ann. soc. entomol. Fr.* **43**, 379–440, <https://doi.org/10.1080/00379271.2007.10697535> (2007).
- Reverté, S. *et al.* National records of 3000 European bee and hoverfly species: A contribution to pollinator conservation. *Insect Conserv. Divers.* **16**, 758–775, <https://doi.org/10.1111/icad.12680> (2023).
- Gaspar, H. *et al.* New regional contributions to the knowledge of the Portuguese bee fauna (Hymenoptera: Anthophila). *Bol. Soc. Entomol. Aragonesa* **71**, 54–62 (2022).
- Varnava, A. I. *et al.* The wild bees (Hymenoptera, Apoidea) of the island of Cyprus. *ZooKeys* **924**, 1–114, <https://doi.org/10.3897/zookeys.924.38328> (2020).
- de Manincor, N. In praise of non-standardized collecting: on the wild bee fauna (Hymenoptera: Anthophila) of Greece (Version v1) [Data set]. *Zenodo* <https://doi.org/10.5281/zenodo.17135537> (2025).
- National Biodiversity Data Centre. *Bees of Ireland*. National Biodiversity Data Centre, Ireland (<https://maps.biodiversityireland.ie/Dataset/5>, 2022).
- Praz, C. *et al.* *Swiss National Apoidea Databank*, Version 1.7. (Swiss National Biodiversity Data and Information Centres – infosppecies.ch, <https://doi.org/10.15468/ksfmzj>, 2022).

34. Bartomeus, I. *et al.* Iberian bees database - Base de datos de abejas ibéricas. *Ecosistemas* **31**, 2380–2380, <https://doi.org/10.7818/ECOS.2380> (2022).
35. Simon Dello, N. S., Otalora, S., Sušan, G., Rubinigg, M. & San Martin, G. *The EU Pollinator Hub: Operationalisation of the EU Bee Partnership Platform for Harmonised Data Collection and Sharing Among Stakeholders on Bees and Pollinators* (EFSA Support. Publ. 22, EN-9219, <https://doi.org/10.2903/sp.efsa.2025.EN-9219> (2025).
36. Wilkinson, M. *et al.* The FAIR Guiding Principles for scientific data management and stewardship. *Sci. Data* **3**, 160018, <https://doi.org/10.1038/sdata.2016.18> (2016).
37. Claris International Inc. FileMaker Pro (Version 19) [Computer software]. <https://www.claris.com/filemaker/> (2019).
38. Google. *Google Maps (Online map)* <https://maps.google.com> (2022).
39. Wickham, H., François, R., Henry, L., Müller, K. & Vaughan, D. Package ‘dplyr’: a grammar of data manipulation, <https://doi.org/10.32614/CRAN.package.dplyr> (2023).
40. Pebesma, E., Bivand, R. *Spatial data science: With applications in R*. Chapman and Hall/CRC (2023).
41. Pebesma, E. Simple features for R: standardized support for spatial sector data. *The R J* **10**, 439–446, <https://doi.org/10.32614/RJ-2018-009> (2018).
42. Ghisbain, G. *et al.* The new annotated checklist of the wild bees of Europe (Hymenoptera: Anthophila). *Zootaxa* **5327**, 1–147, <https://doi.org/10.11646/zootaxa.5327.1.1> (2023).
43. Robinson, D., Bryan, J. & Elias, J. Fuzzyjoin: join tables together on inexact matching <https://doi.org/10.32614/CRAN.package.fuzzyjoin> (2022).
44. Van der Loo, M. *et al.* Package ‘stringdist’: approximate string matching, fuzzy text search, and string distance functions, <https://doi.org/10.32614/CRAN.package.stringdist> (2023).
45. Rasmont, P., Devalez, J., Pauly, A., Michez, D. & Radchenko, V. G. Addition to the checklist of IUCN European wild bees (Hymenoptera: Apoidea). *Ann. la Soc. Entomol. Fr.* **53**, 17–32, <https://doi.org/10.1080/00379271.2017.1307696> (2017).
46. TIBCO Software Inc. Data Science Workbench, version 14, <http://tibco.com> (2020).
47. Sentil *et al.* European Bee and Hovefly Database (EBHD). *Zenodo* <https://doi.org/10.5281/zenodo.17107215> (2025).
48. Benrezkallah, J., Sentil, A., Michez, D. & de Manincor, N. Safeguarding-European-wild-pollinators/SafeguardEuropeanWildBee: Spatial data script analysis (V1.0). *Zenodo* <https://doi.org/10.5281/zenodo.14884373> (2025).
49. QGIS Development Team. QGIS Geographic Information System. *Open Source Geospatial Foundation Project* <http://qgis.osgeo.org> (2024).
50. Gusenleitner, F. & Schwarz, M. Weltweite Checkliste der Bienengattung *Andrena*: mit Bemerkungen und Ergänzungen zu paläarktischen Arten (Hymenoptera, Apidae, Andreninae, *Andrena*). *Entomofauna*, Suppl. **12**, 1–1280 (2002).
51. Wickham, H. *ggplot2: Elegant graphics for data analysis* (Springer-Verlag, New York, 2016).
52. Wickham, G. & Wickham, H. Dates and times made easy with lubridate. *J. Stat. Softw.* **40**, 1–25, <https://doi.org/10.18637/jss.v040.i03> (2011).
53. Wickham, H., Vaughan D. & Girlich, M. tidy: Tidy Messy Data. R package version 1.3.1, <https://doi.org/10.32614/CRAN.package.tidy> (2024).

Acknowledgements

A. Sentil, M. Miličić, D. Michez, I. Steffan-Dewenter, I. Bartomeus, L. Marini, A. Andrić, J. Ačanski, S. Veselić, A. Vujić, S. G. Potts and T. Skendžić are funded by the H2020 project Safeguard (Safeguarding European wild pollinators, grant agreement No. 101003476). A. Andrić, M. Miličić, J. Ačanski, S. Veselić, A. Vujić, S. Popov, S. Mudri Stojnić, T. Tot, A. Grković, M. Janković Milosavljević, S. Radenković and L. Likov acknowledge the financial support of the Ministry of Science, Technological Development and Innovation of the Republic of Serbia (Grants No. 451-03-137/2025-03/200125, 451-03-136/2025-03/ 200125 and 451-03-136/2025-03/200358). M. Miličić, A. Vujić, T. Tot, A. Grković, M. Janković Milosavljević, S. Radenković, L. Likov and A. Đorđević are supported by Taxo-Fly (Taxonomic resources for European hoverflies, contract No. 09.029901/2021/850402/SER/ENV.D.2). A. Andrić, M. Miličić, J. Ačanski, A. Vujić, D. Michez, J. Benrezkallah, S. Reverté and S. GPotts are supported by the Horizon Europe Project WildPosh (Pan European assessment, monitoring, and mitigation of chemical stressors on the health of wild pollinators, Grant No agreement No. 101135238). S. G. Potts, P. Rosa, D. Michez and V. Radchenko are funded by the DG ENV project ORBIT (Taxonomic resources for European bees, contract No. 09.029901/2021/848268/SER/ENV.D.2). P.Y. Maestracci and M. Gibernau are funded by the CPER project No. 40137 “BiodivCorse—Explorer la biodiversité de la Corse” (Collectivité de Corse—Ministère de la Cohésion du territoire et des Relations avec les Collectivités territoriales/les Collectivités territoriales). I. Bartomeus, L. Marini, D. Michez and M. Boustani are funded by the DG ENV project PULSE (Providing technical and scientific support in measuring the pulse of European biodiversity using the Red List Index contract No. 07.027755/2020/840209/SER/ENV.D.2.). M. Miličić and J. Ačanski are supported by the H2020 project ANTARES, Grant/Award Number: 739570. S.G. Potts, D. Michez, T. Petanidou and S. Reverté are funded by the DG ENV project SPRING (Strengthening Pollinator Recovery through Indicators and monitoring, contract No. 09.02001/2021/847887/SER/ENV.D.2). C. Tourbez and B. DeTandt are funded by the F.R.S.-FNRS PhD grant ‘Aspirant’. F. Sgolastra and L. Lenzi are funded by the the Agritech National Research Center and received funding from the European Union Next-GenerationEU (PIANO NAZIONALE DI RIPRESA E RESILIENZA (PNRR) – MISSIONE 4 COMPONENTE 2, INVESTIMENTO 1.4 – D.D. 1032 17/06/2022, CN00000022). P. Y. Maestracci is funded by the CIFRE doctoral program of the ANRT (No. 2022/0393) between the ENGIE—Lab CRIGEN, the University Pantheon-Assas Paris and the University of Corsica. L. Marini is supported by the European Union Next-GenerationEU (PIANO NAZIONALE DI RIPRESA E RESILIENZA (PNRR) – MISSIONE 4 COMPONENTE 2, INVESTIMENTO 1.4 – D.D. 1032 17/06/2022, CN00000022) and the Agritech National Research Center. V. Radchenko is funded by the Program “Support for the development of priority areas of scientific research in Ukraine”, KPKVK 6541230. C. Siopa is funded by a doctoral grant under the fellowships SFRH/BD/145962/2019 and COVID/BD/153552/2024, financed by the Portuguese Foundation for Science and Technology. C. Ruiz is funded by the AEI-Spanish State Research Agency (PID2020-117758GA-I00). J. Loureiro is funded by the Aligning Reference Collections with tAxonomic Development Efforts (ARCADE) for pollinator conservation in Portugal (3PP Project under TETTRIs project, reference nr. 2/T1). R. Santerre received a doctoral scholarship from Wallonie-Bruxelles International (Excellence WBI.World). N. de Manincor is funded by H2020 Marie Skłodowska-Curie Actions COFUND | C2W (Marie Skłodowska Curie grant agreement No. 101034383). S. Castro is funded by PolinizAÇÃO - Action Plant for the Conservation and Sustainability of Pollinators in

Portugal (Fundo Ambiental, Ministry of Environment and Energy). H. Gaspar is funded by the Portuguese Foundation for Science and Technology – FCT – through the fellowship 2023.01736.BD. M. Gérard is funded by a postdoctoral grant “Chargé de Recherches” from the F.R.S.-FNRS (n° 1.B.168.23 F). D. Sommaggio is funded by the National Recovery and Resilience Plan (NRRP), Mission 4 Component 2 Investment 1.4 - Call for tender No. 3138 of 16 December 2021, rectified by Decree n.3175 of 18 December 2021 of Italian Ministry of University and Research funded by the European Union – NextGenerationEU. Project code CN_00000033, Concession Decree No. 1034 of 17 June 2022 adopted by the Italian Ministry of University and Research, CUP E93C22001090001. Project title “National Biodiversity Future Center – NBFC”. P. Cerretti is funded by The European Union–NextGenerationEU as part of the National Biodiversity Future Center, Italian National Recovery and Resilience Plan (NRRP) Mission 4 Component 2 Investment 1.4 (CUP: B83C22002950007). M. DeGroot is funded by the program core group P4-0107 financed by the Slovenian Research and Innovation Agency. The Finnish data collection was partially funded through the Synthesys + Virtual Access program (Call 1 2020). A significant part of Luxembourg’s recent hoverfly data were obtained via the Wild pollinator monitoring program Luxembourg (MONIPOL), funded by the Ministry of the Environment, Climate and Biodiversity (MECB) and the Luxembourg Institute of Science and Technology (LIST). The Polish data collection was partly funded by the 2018-2019 BiodivERsA3 ERA-Net COFUND program and by the funding organization NCN UMO-2019/32/Z/NZ8/00006. The Greek data are mostly funded by POL-AEGIS – Pollinators of the Aegean: Biodiversity and Threats (Greek-ESF funded THALES program, Grant # MIS 376737), SERAPIS – Self-restoration of pollination services in Mediterranean post-fire communities co-considering fire traits and grazing stress (Hellenic Foundation for Research and Innovation, Project # HFRI-FM17-1139) and the project ALARM - Assessing LArge scale environmental Risks for biodiversity with tested Methods (Grant # GOCE-CT-2003-506675). We thank GRoupe d’Étude des Invertébrés Armoricaïns (GRETIA) for sharing data from north-western France from programs funded by various public agencies as well as numerous volunteers. We thank Nicolas Vereecken, Léon Marshal, Nicolas Leclercq, Francisco La Roche and Szentgyörgyi Hajnalka for data provision and Romain Le Divelec for data validation. We thank Andreas Müller (andreas.mueller@naturumweltwissen.ch) for providing his private dataset on European Osmiini and the European Invertebrate Survey (EIS), Leiden, The Netherlands for the provision of the national dataset of Netherlands (menno.reemer@naturalis.nl). We thank Yvan Barbier for his guidance in data processing and Dimitri Evrard for his technical support. We thank the Beelibre team (beelibre.lu) and Fernanda Herrera-Mesías for the photographic material and photos selection. We thank Georgios Nakas and Konstantinos Minachilis for the insect collection. We thank Yulia V. Astafurova (Zoological Institute, St. Petersburg, Russia) and Timofey V. Levchenko (State Darwin Museum, Moscow, Russia) for providing Russian literature. We thank Véronique Sarthou for the identification of the Corsican hoverfly material and Alexandre Cornuel-Willermoz for the identification of the Corsican bee material.

Author contributions

A. Sentil, J. Acanski, J. Benrezkallah, M. Milicic and N. de Manincor provided, processed, and validated data, and wrote the manuscript. D. Michez and G. Ghisbain provided, and validated data, and wrote the manuscript. B. DeTandt, F. Carion, I. Bartomeus, J. Diaz-Calafat, J. Thulier, M. Boustani, S. Lemaire, V. Forgeois and W. Fiordaliso provided and processed data. A. Perrard, A. Ssymank, A. Vujic, A. Zmuda, B. Geslin, C. Ruiz, D. Zimmermann, J. Devalez, J. Kierat, J. Litman, J. Paukkunen, J. Straka, M. Aubert, M. Kasperek, M. Kuhlmann, P. Bogusch, P. Rasmont, R. Santerre, S. Flaminio, S. Gadoum, T. Brau, T. J. Wood, T. Petanidou, E. Budrys, E. Dufrêne, S. M. Stojnić, T. Tot and V. Radchenko provided and validated data. A. Ricarte, M. JankoviMilosavljević and W. vanSteenis provided data and wrote the manuscript. A. Andric, A. Cappellari, A. Đorđević, A. Dormal, A. Grkovic, A. Herinckx, A. Prokhorov, A. Varnava, A. Vesnic, C. Dussaix, C. Geppert, C. Siopa, C. Tourbez, D. Imre, D. Kleijn, D. Sommaggio, E. Gazzea, E. Ockermueller, F. Sgolastra, M. Garrin, G. Pennards, G. Popov, G. Stahls, H. H. Dathe, H. Gaspar, H. Heimburg, I. Steffan-Dewenter, J. Loureiro, F. J. Ortiz-Sanchez, J. Scheper, J. Smit, J. vanSteenis, K. Janssen, L. A. FedorovaZaytseva, L. B. Carstensen, L. Cantú-Salazar, L. Lenzi, L. Likov, L. Marini, M. DeGroot, M. Gibernau, M. Jacobs, M. Mei, M. Mignot, M. Selis, M. Stavrinides, P. Biella, P. Cerretti, P. Rosa, P. Sagot, P. Y. Maestracci, R. Devorsine, R. Foldesi, S. Bot, S. Castro, S. B. Lopes, S. Patiny, S. Popov, S. Potts, S. Pruner, S. Radenkovic, S. Reverte, S. Rojo, S. Veselic, S. Vray, T. Romig, U. Maritano, V. Shparyk, V. Soon, S. Cavallès, G. Ståhls, X. Lair, T. Skendžić and Z. Nedeljkovic provided data. M. Gerard and P. Williams validated data. All authors have read, revised, and approved the final version of the manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Correspondence and requests for materials should be addressed to A.S.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher’s note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License, which permits any non-commercial use, sharing, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if you modified the licensed material. You do not have permission under this licence to share adapted material derived from this article or parts of it. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

© The Author(s) 2026

Ahlam Sentil^{1,2,60} , Marija Miličić^{3,60} , Jordan Benrezkallah^{1,4,60} , Jelena Ačanski³, Andrijana Andrić³, Matthieu Aubert⁵, Ignasi Bartomeus⁶, Paolo Biella⁷, Mira Boustani¹, Leif Bloss Carstensen⁵⁹, Petr Bogusch⁸, Sander Bot⁵⁹, Thomas Brau¹, Eduardas Budrys⁹, Lisette Cantú Salazar¹⁰ , Andree Cappellari¹¹, Frédéric Carion¹, Silvia Castro¹², Simon Cavaillès⁵⁹, Pierfilippo Cerretti¹³, Holger H. Dathe¹⁴, Maarten De Groot¹⁵ , Bastien De Tandt^{1,16} , Jelle Devalez^{17,18}, Rémi Devorsine¹ , Joan Díaz-Calafat¹⁹ , Aleksandra Đorđević²⁰, Ambre Dormal¹, Eric Dufrêne⁵, Cyrille Dussaix⁵⁹, Liuba A. Fedorova Zaytseva¹, William Fiordaliso⁴, Simone Flaminio¹, Rita Földesi⁵⁹, Vianney Forgeois¹, Serge Gadoum²¹, Mael Garrin²², Hugo Gaspar¹² , Elena Gazzea¹¹, Costanza Geppert¹¹, Maxence Gérard¹, Benoît Geslin²³ , Guillaume Ghisbain¹, Marc Gibernau²⁴, Ana Grković²⁰, Helge Heimburg²⁵, Arthur Herinckx¹, Imre Demeter²⁶ , Marina Janković Milosavljević²⁰, Kobe Janssen⁵⁹, Max Kasperek⁵⁹, Justyna Kierat²⁷, David Kleijn²⁸, Michael Kuhlmann²⁹, Xavier Lair⁵⁹, Shawn Lemaire¹, Lucia Lenzi³⁰, Laura Likov²⁰, Jessica Litman³¹, Sara Brito Lopes¹², João Loureiro¹², Jacobs Maarten³², Pierre-Yves Maestracci²⁴, Lorenzo Marini¹¹ , Umberto Maritano⁵⁹, Maurizio Mei¹³, Maud Mignot¹, Sonja Mudri Stojnić²⁰, Zorica Nedeljković³³, Esther Ockermueller³⁴, F. Javier Ortiz-Sanchez³⁵, Sébastien Patiny¹, Juho Paukkunen³², Gerard Pennards⁵⁹, Adrien Perrard³⁶, Theodora Petanidou¹⁷, Grigory Popov^{37,38}, Snežana Popov²⁰, Simon G. Potts³⁹ , Alex Prokhorov³⁷, Stefan Pruner⁵⁹, Vladimir Radchenko⁴⁰ , Snežana Radenković²⁰, Pierre Rasmont¹, Sara Reverté¹, Antonio Ricarte³³, Santos Rojo³⁸, Thomas Romig⁴¹, Paolo Rosa¹, Carlos Ruiz⁴², Philippe Sagot⁵⁹, Rémi Santerre¹, Jeroen Scheper²⁸, Marco Selis⁵⁹, Fabio Sgolastra³⁰ , Viktor Shparyk⁴³, Catarina Siopa¹² , Tea Skendžić²⁰, Jan Smit⁵⁹, Daniele Sommaggio^{44,45}, Villu Soon⁴⁶, Axel Ssymanck⁴⁷, Gunilla Ståhls³², Menelaos Stavrinides^{48,49}, Ingolf Steffan-Dewenter⁵⁰, Jakub Straka⁵¹, Joffrey Thulier¹, Tamara Tot²⁰, Clément Tourbez¹, Jeroen van Steenis⁵², Wouter van Steenis⁵³, Androulla Varnava⁴⁸, Sanja Veselić²⁰, Adi Vesnic⁵⁴, Sarah Vray¹⁰, Paul Williams⁵⁵, Thomas James Wood^{1,56}, Dominique Zimmermann⁵⁷, Aleksandra Žmuda^{27,58}, Ante Vujić^{20,61}, Denis Michez^{1,61} & Natasha de Manincor^{1,61}

¹Laboratory of Zoology, University of Mons, Research Institute for Biosciences, Place du parc 20, 7000, Mons, Belgium. ²Department of Plant and Environment Protection, Agroecology Unit, National School of Agriculture, Meknes, Morocco. ³BioSense Institute – Research Institute for Information Technologies in Biosystems, University of Novi Sad, Novi Sad, Serbia. ⁴Interaction Ecology and Global Change, University of Mons, Research Institute for Biosciences, Place du parc 20, 7000, Mons, Belgium. ⁵Observatoire des Abeilles, Bretagne, France. ⁶Doñana Biological Station (EBD-CSIC), Seville, Spain. ⁷Department of Biotechnology and Biosciences, University of Milano-Bicocca, Milano, Italy. ⁸Department of Biology, Faculty of Science, University of Hradec Kralove, Rokitanského 62, CZ-500 03, Hradec Kralove, Czechia. ⁹State Scientific Research Institute Nature Research Centre (NRC), Akademijos g. 2, LT-08412, Vilnius, Lithuania. ¹⁰Environmental Sensing and Modelling unit, Biodiversity Monitoring and Assessment group Luxembourg Institute of Science and Technology, 41 rue du Brill, L-4422, Belvaux, Luxembourg. ¹¹Department of Agronomy, Food, Natural Resources, Animals and Environment (DAFNAE), University of Padova, Viale dell'Università 16, 35020, Legnaro, Padua, Italy. ¹²Department of Life Sciences, Centre for Functional Ecology, Associate Laboratory TERRA, University of Coimbra, Calçada Martim de Freitas, 3000-456, Coimbra, Portugal. ¹³Department of Biology and Biotechnologies "Charles Darwin", Sapienza University of Rome, Rome, Italy. ¹⁴Senckenberg Deutsches Entomologisches Institut, Eberswalder Straße 90, 15374, Müncheberg, Germany. ¹⁵Slovenian Forestry Institute, Ljubljana, Slovenia. ¹⁶Spatial Epidemiology Lab (SpELL), Université Libre de Bruxelles,

1000, Brussels, Belgium. ¹⁷Laboratory of Biogeography & Ecology, University of the Aegean, 81100, Mytilene, Greece. ¹⁸University of Tartu Natural History Museum, Tartu, Estonia. ¹⁹Department of Biology, University of the Balearic Islands, Balearic Islands, 07001, Palma, Spain. ²⁰Department of Biology and Ecology, University of Novi Sad, Faculty of Sciences, Novi Sad, Serbia. ²¹Office pour les insectes et leur environnement (Opie), BP 30, 78041, Guyancourt, France. ²²Groupe d'Étude des Invertébrés Armoricains (GRETIA), Rennes, France. ²³Université de Rennes (UNIR), UMR 6553 ECOBIO, CNRS, Rennes, France. ²⁴CNRS – University of Corsica - Laboratory Sciences for the Environment (UMR 6134 SPE), Natural Resources Project, Ajaccio, France. ²⁵Department of Biology, University of Graz, Graz, Austria. ²⁶Lendület Ecosystem Services Research Group, Institute of Ecology and Botany, HUN-REN Centre for Ecological Research, Budapest, Hungary. ²⁷Institute of Botany, Faculty of Biology, Jagiellonian University, Gronostajowa 9, 30-387, Kraków, Poland. ²⁸Plant Ecology and Nature Conservation Group, Wageningen University, 6708PB, Wageningen, The Netherlands. ²⁹Zoological Museum, University of Kiel, Hegewischstr. 3, D-24105, Kiel, Germany. ³⁰Department of Agricultural and Food Sciences, Alma Mater Studiorum – University of Bologna, Bologna, Italy. ³¹Natural History Museum of Neuchâtel, Terreaux 14, 2000, Neuchâtel, Switzerland. ³²Zoology Unit, Finnish Museum of Natural History, P.O. Box 17, FI-00014 University of Helsinki, Helsinki, Finland. ³³Research Institute CIBIO (Centro Iberoamericano de la Biodiversidad), Science Park, University of Alicante, Alicante, Spain. ³⁴Biodiversity Centre Linz of the Upper Austrian State Museum, Johann-Wilhelm-Klein-Straße 73, 4040, Linz, Austria. ³⁵Research Group “Technology Transfer in the Area of Natural Resources”, University of Almería, Ctra. de Sacramento s/n, E-04120 La Cañada, Almería, Spain. ³⁶Institute of Ecology and Environmental Sciences, Sorbonne Université, INRAE, IRD, UPEC, Université Paris Cité, 4 Place Jussieu FR-75005, Paris, France. ³⁷Schmalhausen Institute of Zoology, NAS of Ukraine, Kyiv, Ukraine. ³⁸Department of Environmental Sciences & Natural Resources, Faculty of Sciences, University of Alicante, Alicante, Spain. ³⁹Centre for Agri-Environmental Research (CAER), School of Agriculture, Policy and Development, University of Reading, RG6 6AR, Reading, United Kingdom. ⁴⁰Institute for Evolutionary Ecology of the National Academy of Sciences of Ukraine, acad. Lebedev, 37, 01001, Kyiv, Ukraine. ⁴¹University of Hohenheim, Center for Biodiversity and Integrative Taxonomy, Stuttgart, Germany. ⁴²Department of Animal Biology, Edaphology, and Geology, University of La Laguna, San Cristóbal de La Laguna, Spain. ⁴³Vasyl Stefanyk Precarpathian National University, Ivano-Frankivsk, Ukraine. ⁴⁴Department of Life Sciences, Laboratory of Applied Entomology, University of Modena and Reggio Emilia, via Giovanni Amendola 2, 4112, Modena, Italy. ⁴⁵National Biodiversity Future Center (NBFC), Piazza Marina, 61, Palermo, Italy. ⁴⁶Natural History Museum, University of Tartu, Vanemuise 46, 50090, Tartu, Estonia. ⁴⁷Bundesamt für Naturschutz, Bonn, Germany. ⁴⁸Cyprus University of Technology, Archbishop Kyprianos 30, 3036, Limassol, Cyprus. ⁴⁹Eratos.thenes Centre of Excellence, Franklin Roosevelt 82, 3012, Limassol, Cyprus. ⁵⁰Department of Animal Ecology and Tropical Biology, Biocenter, University of Würzburg, Am Hubland, Germany. ⁵¹Department of Zoology, Faculty of Science, Charles University, Vinicna 7, Prague 2, CZ-12844, Staré Město, Czechia. ⁵²Syrphidae Foundation, Baarn, The Netherlands. ⁵³Natuurmonumenten, Amersfoort, The Netherlands. ⁵⁴Department of Biology, Faculty of Science, University of Sarajevo, Sarajevo, Bosnia and Herzegovina. ⁵⁵Natural History Museum, Cromwell Road, SW7 5BD, London, United Kingdom. ⁵⁶Naturalis Biodiversity Center, Darwinweg 2, 2333 CR, Leiden, The Netherlands. ⁵⁷Second Zoological Department, Natural History Museum Vienna, Burgring 7, 1010, Vienna, Austria. ⁵⁸Doctoral School of Exact and Natural Sciences, Jagiellonian University, prof. S. Łojasiewicza 11, 30-348, Kraków, Poland. ⁵⁹Independent researcher: Leif Bloss Carstensen, Sander Bot, Simon Cavaillès, Cyrille Dussaix, Rita Földesi, Kobe Janssen, Max Kasperek, Xavier Lair, Umberto Maritano, Gerard Pennards, Stefan Pruner, Philippe Sagot, Marco Selis, Jan Smit. ⁶⁰These authors contributed equally: Ahlam Sentil, Marija Miličić, Jordan Benrezkallah. ⁶¹These authors jointly supervised this work: Ante Vujić, Denis Michez, Natasha de Manincor. ⁶²e-mail: ahlam.sentil@umons.ac.be