

## Drivers of amphibian population dynamics and asynchrony at local and regional scales

Hugo Cayuela, Richard A. Griffiths, Nurul Zakaria, Jan W. Arntzen, Pauline Priol, Jean-Paul Léna, Aurélien Besnard, Pierre Joly

Downloaded from:

<http://dx.doi.org/10.1111/1365-2656.13208>

### Article 25fa Dutch Copyright Act (DCA) - End User Rights

This publication is distributed under the terms of Article 25fa of the Dutch Copyright Act (Auteurswet) with consent from the author. Dutch law entitles the maker of a short scientific work funded either wholly or partially by Dutch public funds to make that work publicly available following a reasonable period after the work was first published, provided that reference is made to the source of the first publication of the work.

This publication is distributed under the Naturalis Biodiversity Center 'Taverne implementation' programme. In this programme, research output of Naturalis researchers and collection managers that complies with the legal requirements of Article 25fa of the Dutch Copyright Act is distributed online and free of barriers in the Naturalis institutional repository. Research output is distributed six months after its first online publication in the original published version and with proper attribution to the source of the original publication.

You are permitted to download and use the publication for personal purposes. All rights remain with the author(s) and copyrights owner(s) of this work. Any use of the publication other than authorized under this license or copyright law is prohibited.

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the department of Collection Information know, stating your reasons. In case of a legitimate complaint, Collection Information will make the material inaccessible. Please contact us through email: [collectie.informatie@naturalis.nl](mailto:collectie.informatie@naturalis.nl). We will contact you as soon as possible.

# Drivers of amphibian population dynamics and asynchrony at local and regional scales

Hugo Cayuela<sup>1</sup>  | Richard A. Griffiths<sup>2</sup> | Nurul Zakaria<sup>2</sup> | Jan W. Arntzen<sup>3</sup> |  
Pauline Priol<sup>4</sup> | Jean-Paul Léna<sup>5</sup> | Aurélien Besnard<sup>6</sup> | Pierre Joly<sup>5</sup> 

<sup>1</sup>Institut de Biologie Integrative et des Systèmes (IBIS), Université Laval, Québec, QC, Canada; <sup>2</sup>Durrell Institute of Conservation and Ecology, School of Anthropology and Conservation, University of Kent, Canterbury, Kent, UK; <sup>3</sup>Naturalis Biodiversity Center, Leiden, The Netherlands; <sup>4</sup>StatiPOP, Ganges, France; <sup>5</sup>UMR 5023 LEHNA, Université de Lyon, Lyon1, CNRS, ENTPE, Villeurbanne, France and <sup>6</sup>CNRS, PSL Research University, EPHE, UM, SupAgro, IRD, INRA, UMR 5175 CEFE, Montpellier, France

## Correspondence

Hugo Cayuela

Email: hugo.cayuela51@gmail.com

Handling Editor: Mark Hewison

## Abstract

1. Identifying the drivers of population fluctuations in spatially distinct populations remains a significant challenge for ecologists. Whereas regional climatic factors may generate population synchrony (i.e. the Moran effect), local factors including the level of density dependence may reduce the level of synchrony. Although divergences in the scaling of population synchrony and spatial environmental variation have been observed, the regulatory factors that underlie such mismatches are poorly understood.
2. Few previous studies have investigated how density-dependent processes and population-specific responses to weather variation influence spatial synchrony at both local and regional scales. We addressed this issue in a pond-breeding amphibian, the great crested newt *Triturus cristatus*. We used capture–recapture data collected through long-term surveys in five *T. cristatus* populations in Western Europe.
3. In all populations—and subpopulations within metapopulations—population size, annual survival and recruitment fluctuated over time. Likewise, there was considerable variation in these demographic rates between populations and within metapopulations. These fluctuations and variations appear to be context-dependent and more related to site-specific characteristics than local or regional climatic drivers. We found a low level of demographic synchrony at both local and regional levels. Weather has weak and spatially variable effects on survival, recruitment and population growth rate. In contrast, density dependence was a common phenomenon (at least for population growth) in almost all populations and subpopulations.
4. Our findings support the idea that the Moran effect is low in species where the population dynamics more closely depends on local factors (e.g. population density and habitat characteristics) than on large-scale environmental fluctuation (e.g. regional climatic variation). Such responses may have far-reaching consequences for the long-term viability of spatially structured populations and their ability to respond to large-scale climatic anomalies.

## KEYWORDS

amphibian, crested newt, demography, density dependence, population synchrony, weather

## 1 | INTRODUCTION

Understanding the mechanisms driving population dynamics is a critical challenge in ecology and conservation biology. In recent decades, researchers have paid particular attention to the mechanisms involved in the synchronization of population dynamics at different scales (Bjørnstad, Ims, & Lambin, 1999; Koenig & Liebhold, 2016; Liebhold, Koenig, & Bjørnstad, 2004; Ranta, Kaitala, Lindstrom, & Linden, 1995). Three mechanisms have been widely recognized to cause population synchrony: dispersal between populations (Kendall, Bjørnstad, Bascompte, Keitt, & Fagan, 2000; Ylikarjula, Alaja, Laakso, & Tesar, 2000); trophic interactions with other species that are themselves either synchronized or mobile (Ims & Andreassen, 2000; Korpimäki, Norrdahl, Huitu, & Klemola, 2005); and synchronous stochastic environmental effects, called the Moran effect (Koenig, 2002; Moran, 1953). While the first two factors usually drive population synchrony at relatively local spatial scales, the Moran effect is considered a major mechanism for generating population synchrony at regional level (Koenig, 2002).

Moran's (1953) concept states that synchrony in the dynamics of populations regulated by the same density-dependent processes will reflect correlated environmental perturbations. Consequently, population synchrony is expected to decay with increasing distance between populations in a manner similar to that of the synchrony of a potential underlying environmental driver. However, population synchrony usually occurs at a far smaller spatial scale than that at which environmental drivers operate (Bjørnstad, Liebhold, & Johnson, 2008; Koenig, 2001; Peltonen, Liebhold, Bjørnstad, & Williams, 2002; Trenham, Koenig, Mossman, Stark, & Jagger, 2003). For example, simulation studies have revealed that the degree of population synchrony is influenced by the form of density regulation. Specifically, population-specific density dependence and nonlinear relationships between demographic rates and density typically reduce the synchronizing effect of spatially correlated environmental noise (Bjørnstad et al., 1999; Ranta et al., 1995; Royama, 2005). Secondly, theoretical models predict that spatial variation of the demographic parameters contributing to population growth rate negates the Moran effect and then reduces population synchrony (Engen & Sæther, 2005; Hugueny, 2006). In particular, variation in survival and recruitment caused by local environmental factors such as predation, interspecific competition and habitat quality could decrease population synchrony. Simulations have also demonstrated that the contribution of a covariate to spatial synchrony strongly depends on spatial heterogeneity in the covariate or on its effect on local dynamics (Engen & Sæther, 2005; Hugueny, 2006). Importantly, local expression of large-scale climatic phenomena could lead to context-dependent demographic responses to weather variation

(Selwood, McGeoch, & Mac Nally, 2015), which could then result in low spatial synchrony.

Several empirical studies have reported divergences in the scaling of population synchrony and spatial environmental variation (Koenig, 2001; Peltonen et al., 2002; Trenham et al., 2003). However, few have empirically examined the factors that could underlie such mismatches (Haynes, Bjørnstad, Allstadt, & Liebhold, 2013). No previous work has investigated how population-specific density-dependent processes and population-specific responses to environmental variation (e.g. weather) could affect spatial synchrony at both local and regional scales. Pond-breeding amphibians are excellent biological models to address this issue. First, population synchrony has already been highlighted in these organisms (Trenham et al., 2003) and spatially correlated environmental weather variables (i.e. temperature and precipitation) could have strong synchronizing effects. Yet, several factors with potentially desynchronizing effects also play an important role in amphibian demography. Variation in pond characteristics may dramatically affect demographic rates (Unglaub, Steinfartz, Drechsler, & Schmidt, 2015; Unglaub, Steinfartz, Kühne, Haas, & Schmidt, 2018). Furthermore, although amphibians may be highly sensitive to climate, their demographic responses to variation in weather seems highly context-dependent (Cayuela et al., 2016a; Muths et al., 2017). Moreover, density dependence has been often reported as a key driver of amphibian population dynamics (Altwegg & Reyer 2003, Harper & Semlitsch, 2007).

In this study, we examined population synchrony, demographic responses to weather and density dependence in the great crested newt *Triturus cristatus*. We used capture-recapture data collected through long-term surveys on five *T. cristatus* populations in Western Europe over a 22-year period (1995–2016). Two of these five populations (i.e. POP1 and POP5) were spatially structured (i.e. composed of subpopulations occupying different ponds or group of ponds). This nested design allowed us to examine demographic processes at both local (i.e. within population) and regional (between population, over western Europe) levels. We predicted a relatively low level of spatial synchrony at both local and regional levels due to the characteristics of amphibian demography stated above. We then examined how local demographic responses to climate and density dependence may trigger population asynchrony. In particular, we investigated how weather fluctuation and population density may affect survival, recruitment and population growth rate in the five populations and in the four subpopulations of POP1 and POP5. Based on previous studies (Cayuela et al., 2016a; Muths et al., 2017), we expected heterogeneous demographic responses (survival, recruitment and population growth) to temperature and rainfall within and between populations. In addition, we predicted negative density dependence to be a common phenomenon. At the regional level,

we assumed that such density-dependent processes would have a desynchronizing effect since populations have different histories, are located in different geographical areas and are not connected by dispersing individuals. At the local level, we also predicted a desynchronizing effect of density as dispersal rates are low between subpopulations (see Material and method section) and because ponds may differ in terms of local characteristics. These multiple-level desynchronizing processes should lead to a negligible Moran effect in our study system.

## 2 | MATERIALS AND METHODS

### 2.1 | Biological model

*Triturus cristatus* populations vary in age structure, with median age varying from 3 to 8 years (Jehle, Thiesmeier, & Foster, 2011). Females usually produce between 150 and 300 eggs per year (Arntzen & Hedlund, 1990; Hagström, 1980). Larval stage lasts 2.5–3 months or more and sexual maturity is reached at 2–3 years (Griffiths, 1996; Miaud, Joly, & Castanet, 1993). Reproduction occurs in still waters such as ponds, flooded quarries, lakes, irrigation canals and ditches where the adults spend usually 2–5 months during spring (from February to June). Outside this period, individuals occupy terrestrial habitats such as hedgerows, shrub, rough grassland or forest edges, and overwinter in underground refugia such as mammal burrows (Jehle & Arntzen, 2000). The crested newt is a species in decline that is on the IUCN Red List in many countries (IUCN, 2020), and it is protected by the European Habitats Directive (92/43/CEE).

### 2.2 | Studied populations, capture–recapture data and weather variables

The study was conducted on five populations in France and the United Kingdom using the capture–recapture method (for a map, see Appendix S1). Two populations (POP1 and POP2) are located in southeastern England and are 2.5 km apart and separated from each other by dispersal barriers, such as roads and unsuitable habitat (Zakaria, 2018). Another population (POP3) is located in western France. The last two populations (POP4 and POP5) are located in southeastern France. The populations POP2, POP3 and POP4 occupy single breeding sites. POP1 and POP5 are spatially structured populations, both composed of four distinct subpopulations. In POP1, the four subpopulations occupy four distinct ponds, separated from each other by distances ranging from 200 to 800 m. According to Griffiths, Sewell, and McCrea (2010), there is a low level of adult movement among subpopulations with dispersal mainly occurring during the subadult phase. In POP5, the four subpopulations occupy four distinct pond groups. Each group was composed of three very close (15–30 m) ponds between which annual dispersal rates are high ( $<0.20$ ; Cayuela, Grolet, & Joly, 2018). The pond groups were separated from each other by a distance ranging from 60 to

430 m; the number of breeding dispersal events among pond groups was low (i.e. 12 out of 2,282 individuals captured during the period 1996–2015), resulting in a mean dispersal rate of  $<0.01$ . These 12 individuals were discarded from our analyses.

Newts were surveyed over periods ranging from 8 to 20 years between 1995 and 2016. The newts were captured using bottle traps (POP1 and POP2, see Griffiths et al., 2010), funnel traps (POP3), dipnets (POP4, see Cayuela et al., 2017) and seine net (POP5, see Cayuela et al., 2017). The number of capture sessions performed each year varied from 1 (POP5) to 33 (POP1) during the newt activity period in ponds. Note that in capture–recapture analyses, we merged the observations of intra-annual sessions, considering one single session per year. Newts were individually identified using pit-tags in POP3 and POP5, and by photographs of belly pattern markings in POP1, POP2 and POP4. Providing the image quality is high and the identification is done by trained personnel, belly patterns are a highly reliable method for identifying individual newts (Zakaria, 2018). Each individual was classified as adult or juvenile, and adults were sexed on the basis of the presence of a swollen cloaca and a large crest on the back in males (Hedlund, 1990). As the number of juveniles was low in our datasets (juveniles only occasionally found in the ponds), the following analyses were restricted to adults. Detailed information about the capture–recapture surveys (samples size, the number of captured newts, etc.) is provided in Appendix S1.

In Western Europe, both precipitation and temperatures exhibit high synchrony declining with distance; the spatial autocorrelation remains statistically significant over a very large distance (up to 2,500 km; Koenig, 2002). We therefore assumed that these two meteorological parameters could act as potential population synchronizers and could underlie a Moran effect. We considered in our analyses the effect of four weather variables on survival, recruitment and population growth: cumulative precipitation during breeding (March–May; 'rainMM') and non-activity periods (June–February; 'rainJF'); the minimum ('tempDFI') and the maximum temperature ('tempDFS') during winter (December–February). The weather variables were collected at meteorological stations (in France, Météo France stations; in Great Britain, UK Meteorological Office) located in the vicinity (within 3–25 km) of the surveyed populations. These variables were selected based on the biology of the studied species. During the breeding period, low cumulative rainfall can increase mortality and decrease recruitment due to pond desiccation in amphibians in general (Muths et al., 2017) and in newts and salamanders in particular (Church, Bailey, Wilbur, Kendall, & Hines, 2007). On the other hand, high rainfall usually stimulates breeding activities (Wells, 2010) that are energetically demanding, which could increase adult mortality during the reproduction period. During the non-activity period when newts have left the pond and adopted a terrestrial lifestyle, high rainfall and high temperature could increase mortality due to respiratory and energetic constraints (Griffiths et al., 2010; Reading, 2007). These effects were investigated in POP1, POP2, POP3 and POP5 (at the whole population level), as well as in two subpopulations of POP1 (POP1.1 and POP1.2) and

POP5 (POP5.3 and POP5.4); too few individuals were captured in the other subpopulations to perform analyses. As POP4 has a different phenology due to Mediterranean climatic conditions, we considered the four following variables: the cumulative rainfall during the breeding period (October–April, 'rainOA') and the non-activity period (May–September, 'rainMS'); the minimum ('tempMSI') and maximum ('tempMSS') monthly temperature during the non-activity period (May–September). Additional information about weather variables (e.g. descriptive statistics, covariation patterns) is provided in Appendix S1.

### 2.3 | Weather and density-dependent effects on survival and recruitment

We examined transience and trap-dependence by performing goodness-of-fit (GOF) tests in the program U-CARE (Choquet, Lebreton, Gimenez, Reboulet, & Pradel, 2009; Choquet Rouan & Pradel, 2009) and investigated the potential occurrence of recapture heterogeneity using the GOF test proposed by Jeyam, McCrea, Bregnballe, Frederiksen, and Pradel (2018). Using U-CARE, we performed an overall GOF test that was statistically significant in only two populations, namely POP1 ( $df = 118$ ,  $\chi^2 = 320.08$ ,  $p < 0.0001$ ) and POP3 ( $df = 19$ ,  $\chi^2 = 35.81$ ,  $p = 0.01$ ; Appendix S1). In these two populations, test 2CT (POP1,  $\chi^2 = 33.65$ ,  $p = 0.02$ ; POP3,  $\chi^2 = 28.34$ ,  $p = 0.03$ ) detected trap-dependence; that was not the case for 2CI (POP1,  $\chi^2 = 4.84$ ,  $p = 0.93$ ; POP3, not estimated). In parallel, the tests 3SR (POP1,  $\chi^2 = 40.90$ ,  $p = 0.0001$ ; POP3,  $\chi^2 = 25.80$ ,  $p = 0.0002$ ) and 3SM (POP1,  $\chi^2 = 31.69$ ,  $p = 0.01$ ; POP3,  $\chi^2 = 5.13$ ,  $p = 0.16$ ) detected an excess of transients. In parallel, we detected recapture heterogeneity in these two populations by using Jeyam's GOF test (2018). The global test for heterogeneity was highly significant in POP1 (test statistic = 3.46,  $p < 0.001$ ) and POP3 (test statistic = 3.98,  $p < 0.001$ ). In summary, the fact that GOF tests simultaneously detected trap-dependence, transience and heterogeneity indicated overall recapture heterogeneity rather than transience or trap-dependence alone (Jeyam et al., 2018). Following the recommendations of Jeyam et al. (2018) and the approach used in previous studies with a similar situation (Cayuela, Cruickshank, Brandt, Ozgul, & Schmidt, 2019; Chambert et al., 2012; Crespin, Choquet, Lima, Merritt, & Pradel, 2008; Cubaynes et al., 2010), we included Pledger's heterogeneity mixtures (Pledger, 2000) in the multi-event models to take account of heterogeneity in recapture probability.

To investigate whether survival was influenced by conspecific density and weather variables in each of the five populations, we considered a model with three states  $A_1$ ,  $A_2$  and  $D$ , where  $A$  means alive and  $D$  dead and where 1 and 2 correspond to the first and second heterogeneity class respectively. We considered two possible field observations, captured or not captured. The model contained three pieces of information: (a) the vector of initial states probabilities; (b) the survival matrix; and (c) the event matrix linking observations to individual latent states. At their first capture, individuals

could be alternatively in two states:  $A_1$  and  $A_2$ , resulting in the following vector of initial state probabilities:

$$\begin{pmatrix} i_1 & 1 - i_1 \end{pmatrix},$$

where  $i$  is the initial state probability. Then, information about survival was updated: individuals in the state  $A_1$  at  $t - 1$  could reach the state  $A_1$  at  $t$  (i.e. survive) with a probability  $\phi_1$  or die with a probability  $1 - \phi_1$ ; individuals in the state  $A_2$  at  $t - 1$  could reach the state  $A_2$  at  $t$  with a probability  $\phi_2$  or die with a probability  $1 - \phi_2$ . This results in the following matrix:

$$\begin{pmatrix} \phi_1 & 0 & 1 - \phi_1 \\ 0 & \phi_2 & 1 - \phi_2 \\ 0 & 0 & 1 \end{pmatrix}.$$

We did not consider heterogeneity mixtures on survival (usually used to investigate actuarial senescence and variation in life history strategies; Denoël, Dalleur, Langrand, Besnard, & Cayuela, 2018; Péron et al., 2010) and we hold  $\phi_1$  and  $\phi_2$  to be equal in all the models. Lastly, observations were modelled: individuals in the state  $A_1$  could be captured with a probability  $p_1$  or not captured with a probability  $1 - p_1$ ; individuals in the state  $A_2$  could be captured with a probability  $p_2$  or not captured with a probability  $1 - p_2$ . This leads to the following event matrix:

$$\begin{pmatrix} 1 - p_1 & p_1 \\ 1 - p_2 & p_2 \\ 1 & 0 \end{pmatrix}.$$

To examine whether annual adult recruitment (i.e. the proportion of sexually mature individuals recruited each year) was affected by conspecific density and weather variables in each of the five populations, we built a model following the structure of Pradel's (1996) model, in which recruitment was modelled by reversing capture histories and analysing them backwards. The model is conditional upon the first capture. The recruitment probability  $\psi$  was estimated as the probability that an individual present at time  $t$  was not present at  $t - 1$ , that is, the proportion of 'new' individuals in the population at  $t$ , while accounting for capture heterogeneity. Note that in our multi-event model, survival cannot be estimated along with recruitment. This model has the same structure as the previous model and only the recruitment matrix was modified as follows:

$$\begin{pmatrix} 1 - \psi_1 & 0 & \psi_1 \\ 0 & 1 - \psi_2 & \psi_2 \\ 0 & 0 & 1 \end{pmatrix}.$$

We tested our hypotheses about survival and recruitment separately and by following a similar procedure. The models were implemented in the E-SURGE program (Choquet, Rouan, et al., 2009). The datasets from the five populations of *T. cristatus* differed in terms of the number of study years and study periods (Appendix S1). Hence, we separately analysed the five datasets and then compared the

population's responses to weather variations and density using the outputs of the best-fitting models for each population. Models were ranked through a model selection procedure using Akaike information criterion adjusted for a small sample size (AICc). The analyses were carried out in two steps. First, from the most general model [ $\phi(t + \text{sex})$ ,  $p(\text{het} + \text{sex})$ ], we evaluated the three following effects: sex ('sex'), heterogeneity mixture (low vs. high capture rates of individuals; 'het') and year-specific variation ('t'). Only additive models were considered in our analyses; interactions were avoided to limit the problem of parameter identifiability and to increase model stability. We also avoided including year-specific variation on recapture rates due to recurrent convergence issues when the model [ $\phi(t + \text{sex})$ ,  $p(t + \text{het} + \text{sex})$ ] was considered; in many populations, the number of individuals captured each year was relatively low. Time-specific recapture probability was supported in two populations (POP1 and POP4; Appendix S3), however the models [ $\phi(t)$ ,  $p(t)$ ] and [ $\phi(t)$ ,  $p(.)$ ] provided very similar estimates of survival in POP1 and POP4 (Appendix S3). Therefore, [ $\phi(t + \text{sex})$ ,  $p(\text{het} + \text{sex})$ ] was kept as a general model as time-specific survival is required in ANODEV analyses (see below). We examined all the possible combinations of effect, resulting in the consideration of 16 models per population and subpopulation.

In a second step, after determining the best-fitting model, we examined the effect of weather and density variables on survival probability using ANODEV as recommended in Grosbois et al. (2008). This approach allowed us to evaluate the fit of a model including a single meteorological covariate ( $M_{\text{cov}}$ ) relative to both the constant ( $M_{\text{cst}}$ ) and the time-dependent ( $M_t$ ) models. The statistic  $F_{\text{test}_{\text{cst}/\text{cov}/t}}$  has been derived as follows:

$$F_{\text{test}_{\text{cst}/\text{cov}/t}} = \left( \frac{\text{Dev}(M_{\text{cst}}) - \text{Dev}(M_{\text{cov}})}{\zeta - 1} \right).$$

This statistic tests the null hypothesis  $H_0$  that the meteorological covariate in  $M_{\text{cov}}$  has no significant effect on survival or recruitment. It follows under  $H_0$  a Fisher-Snedecor distribution with  $\zeta - 1$  degrees of freedom, where  $\zeta$  is the number of parameters required to describe the relationship between survival and the focal climatic covariate. To examine the effect of population density at  $t - 1$  on survival and recruitment, we estimated population size at  $t - 1$  by using the Horvitz-Thompson estimator. For POP1 and POP3 where recapture heterogeneity was detected using GOF tests, we used the modified version of the Horvitz-Thompson estimator proposed by Cubaynes et al. (2010). All variables (weather and density) were scaled (i.e. z-scored) when they were entered in the models.

## 2.4 | Weather and density-dependent effects on population growth rate

To assess the effect of density dependence and weather on population growth rate, we used Gompertz state-space (GSS) models (de Valpine & Hastings, 2002). We followed the approach described

in Schaub and Kéry (2012) and Băncilă, Ozgul, Hartel, Sos, and Schmidt (2015). The abundance time series of the five populations and the four subpopulations of POP1 and POP5 were analysed separately. The abundances were corrected by the time-constant recapture probability (i.e. Horvitz-Thompson estimator) obtained from the capture-recapture models. As recapture probability fluctuates over time in several populations (Appendix S3), we verified that using population sizes corrected for time-specific recapture probability did not affect our results (Appendix S5). The log-transformed ( $\log(X + 1)$ ) population sizes  $X_t$  in a pond at time  $t$  are described by:

$$X \sim \text{Normal}(N_t, \sigma_{\text{obs}}^2),$$

where  $N_t$  is the unobserved true population size at time  $t$  and  $\sigma_{\text{obs}}^2$  is the observation variance. This piece of the GSS model described the observation process and allows for observation error while assuming that new abundance may overestimate or underestimate true population size.  $N_t$  was defined from a normal distribution:

$$N_t \sim \text{Normal}(N_{\text{exp},t}, \sigma_{\text{proc}}^2),$$

where  $\sigma_{\text{proc}}^2$  designates the process variance (i.e. stochastic variability in new abundance) and  $N_{\text{exp},t}$  the expected population size at time  $t$ . The effect of population density at  $t - 1$  was modelled as follows:

$$N_{\text{exp},t} \sim \text{Normal}(N_{t-1} + a + b_1 N_{t-1}),$$

where  $a$  is the intercept and  $b$  the coefficient slope for the effect density at  $t - 1$ . To evaluate the effect of weather variables on population growth rate, we considered models including one weather variable at a time. The model had the following form:

$$N_{\text{exp},t} \sim \text{Normal}(N_{t-1} + a + b_2 w_t),$$

where  $w_t$  is the z-scored weather covariate at time  $t$  and  $b_2$  is the slope coefficient.

The models were fitted in JAGS (Plummer, 2003), using the R package JAGSUI (Kellner, 2018), with vague normal priors with a mean 0 and a precision of 0.0001 for  $a$  and  $b$  parameters. Vague uniform priors ranged within an interval 0–10 were applied for the standard deviation of  $\sigma_{\text{obs}}^2$  and  $\sigma_{\text{proc}}^2$ . We used a normal prior with a mean equal to the log-transformed population size and a variance of 100 for the first value of the time series. Three Markov Chain Monte Carlo were run with 2,000,000 iterations and a burn-in of 1,000,000. Chains were thinned by a factor of 20. We assessed model convergence with the Gelman-Rubin statistic  $R$ -hat; we assumed that the model convergence was satisfactory when  $R$ -hat values were less than 1.1 (Brooks & Gelman, 1998; Gelman & Hill, 2006). We considered that the effect of a variable on population growth rate was significant if the credible intervals (CRI) of the slope coefficient did not include 0 (Băncilă et al., 2015; Kéry & Schaub, 2011).

## 2.5 | Population synchrony within and between populations

The temporal synchrony among populations was assessed using the variance partitioning methodology (Grosbois et al., 2009; Schaub, Hirschheydt, & Gruebler, 2015). More specifically, we built a GLMM treating the estimate of population size as the dependent variable using a Poisson distribution to specify the error term and the  $\ln$  transformation as the link function. The study population was introduced as a fixed explanatory term in the model. We estimated yearly abundance by dividing the number of captured newts by the time-constant recapture probability obtained from the multi-event capture–recapture models described below (i.e. Horvitz–Thompson estimator). For subpopulations for which capture–recapture models were not built, we used the mean recapture probability estimated for the whole population. As recapture probability may vary over time in several populations (Appendix S3), we verified that using population sizes corrected with time-specific recapture probability did not affect our results. Given the uncertainty of population size estimates, we also controlled for potential underestimation of our estimates of population synchrony. Neither of these adjustments substantially changed our results (Appendix S2). Both time  $t$  and its interactive effect with the population  $t \times p$  were introduced as random effects in the model. The model was thus specified as follows:

$$N_{pt} \sim \text{Poisson}(\lambda_{pt})$$

where  $N_{pt}$  is the estimated demographic size of population  $p$  at time  $t$  following a Poisson distribution with  $\lambda_{pt}$ , the expected demographic size is

$$\ln(\lambda_{pt}) = \sum_{p=1}^k \beta_p x_p + \alpha_t + e_{pt},$$

with

$$\alpha_t \sim \text{Normal}(0, \sigma_\alpha^2),$$

where  $\sigma_\alpha^2$  is the shared temporal variance between the  $k$  populations, and

$$e_{pt} \sim \text{Normal}(0, \sigma_e^2),$$

where  $\sigma_e^2$  is the residual temporal variance specific to each  $p$  population.  $\beta_p$  is the main population effect (associated to the dummy variable  $x_p$  specifying the partial intercept for the population  $p$ ) accounting for the overall difference of demographic size between the  $k$  populations.

The temporal synchrony among populations was then measured using intraclass correlation coefficient (ICC) that expresses the ratio of the temporal variation that is common across all studied populations and the entire amount (i.e. spatiotemporal) of variation. ICC values close to 1 indicate a quasi-perfect temporal

synchrony among populations; values close to 0 indicate complete asynchrony, and values below 0.5 indicate relatively low synchrony. We computed the adjusted intraclass correlation coefficient ( $\text{ICC}_{\text{adj}}$ ) using the delta method to accommodate the latent variance inherent to the Poisson distribution associated to the GLM model (see Nakagawa, Johnson, & Schielzeth, 2017, for justification and implementation):

$$\text{ICC}_{\text{adj}} = \frac{\sigma_\alpha^2}{\sigma_\alpha^2 + \sigma_e^2 + \sigma_d^2},$$

where  $\sigma_d^2$  is the latent variance inherent to the Poisson distribution estimated as

$$\ln\left(1 + \frac{1}{\lambda_{pt}}\right),$$

with


$$E(\lambda_{pt}) \approx \frac{1}{k} \sum_{p=1}^k \beta_p x_p + 0.5(\sigma_\alpha^2 + \sigma_e^2).$$

To estimate population synchrony at the regional scale we restricted the time series of each population to the 2000–2015 period to ensure a balanced dataset across populations. In the case of POP1 and POP5, we used the estimated total population size to compute this estimation. Estimation of population synchrony at the local scale was performed separately for POP1 and POP5 using the population size estimated for each of their subpopulations (i.e. replacing population by subpopulation in the formula above) during the whole study period.

## 3 | RESULTS

### 3.1 | Population synchrony at local and regional scales

Population size drastically varied over time and across populations and subpopulations (Figure 1). The level of synchrony at the local scale varied greatly between populations (Table 1): it was relatively high in POP5 but rather low in POP1 as indicated by their respective ICC. At the regional scale, we found a relatively low level of population synchrony as indicated by the estimated ICC (0.212, Table 1). We further used another GLMM to decompose the total variance as the sum of the shared temporal variance between POP1 and POP5, the shared temporal variance between subpopulations of each population and the residual temporal variance specific to each subpopulation. The level of synchrony between populations was estimated at the lower bound (i.e. 0, Appendix S2), and the average level of synchrony between subpopulations (within each population) was moderate (i.e. 0.488, Appendix S2). In sum, the level of synchrony was variable between subpopulations at a local scale, and low between populations on a regional scale.



**FIGURE 1** Population sizes and their standard errors (derived from survival models) in the five studied populations (POP1, POP2, POP3 and POP4) of great crested newt and their respective subpopulations (POP1: POP1.1, POP1.2, POP1.3 and POP1.4; POP5: POP5.1, POP5.2, POP5.3 and POP5.4)

**TABLE 1** Synchrony among the five populations (i.e. regional level) and within two populations (i.e. local level, POP1 and POP5) of great crested newt. ICC and shared/specific temporal variance of population size.  $N$  is the total number of observations used,  $\sigma_{\alpha}^2$  is the shared temporal variance among populations,  $\sigma_{\epsilon}^2$  is the temporal variance specific to each population,  $E(\lambda_{pt})$  is the expected mean demographic size over all populations


| Scale      | $N$ | $\sigma_{\alpha}^2$ | $\sigma_{\epsilon}^2$ | $E(\lambda_{pt})$ | ICC  |
|------------|-----|---------------------|-----------------------|-------------------|------|
| Regional   | 69  | 0.15                | 0.56                  | 100               | 0.21 |
| Local POP1 | 76  | 0.43                | 1.04                  | 39                | 0.28 |
| Local POP5 | 80  | 0.87                | 0.53                  | 47                | 0.61 |

### 3.2 | Survival and recruitment variation within and between populations


Within populations, our analyses revealed that mean survival (provided by the model where survival was constrained to be constant over time) displayed slight subpopulation-specific variation in

POP1 and POP5. In POP1, the survival was  $0.61 \pm 0.02$  ( $M \pm SE$ ) in POP1.1 and  $0.66 \pm 0.02$  in POP1.2. In POP5, survival ranged from  $0.23 \pm 0.01$  in POP5.3 to  $0.36 \pm 0.01$  in POP5.4. In the different subpopulations of POP1 and POP5, survival varied widely between years (Figure 2). At the population level, survival displayed broad population-specific variation. Mean survival was relatively high in POP2 and POP3,  $0.83 \pm 0.02$  and  $0.87 \pm 0.02$  respectively. Survival was lower ( $0.62 \pm 0.01$ ) in both POP1 and POP4 and drastically lower in POP5 ( $0.29 \pm 0.01$ ). Survival varied between years to a lesser extent in POP2 and POP3 (Figure 2) where mean survival was higher. By contrast, it broadly varied between years in POP1, POP4 and POP5 (Figure 2). The effect of sex on survival was absent or marginal in all the populations and subpopulations (Figure 2, and model selection procedures in Appendix S3).

At the within-population level, recruitment displayed slight subpopulation-specific variation in POP1 and POP5. In POP1, recruitment was  $0.35 \pm 0.02$  in POP1.1 and  $0.32 \pm 0.02$  in POP1.2. In POP5, recruitment was  $0.51 \pm 0.03$  in POP5.3 and  $0.65 \pm 0.02$  in POP5.4. In the different subpopulations of POP1 and POP5, recruitment fluctuated among years (Figure 3). At the population level, mean




**FIGURE 2** Survival probability and their 95% CI in five populations (POP1, POP2, POP3, POP4 and POP5) of great crested newt in Western Europe. Survival has also been estimated in subpopulations (i.e. ponds or groups of ponds) of POP1 (POP1.1 and POP1.2) and of POP5 (POP5.3 and 5.4). In subpopulation POP5.3, we excluded the first five years of survey (1996–2002) because few newts were captured. Males are shown in full circles; females are shown in empty circles



**FIGURE 3** Recruitment probability and their 95% CI in five populations (POP1, POP2, POP3, POP4 and POP5) of great crested newt in Western Europe. Survival has been also estimated in subpopulations (i.e. ponds or groups of ponds) of POP1 (POP1.1 and POP1.2) and of POP5 (POP5.3 and 5.4). In subpopulation POP5.3, we excluded the first five years of survey (1996–2002) because few newts were captured. Males are shown in full circles; females are shown in empty circles

recruitment (estimated in the model where recruitment was constrained to be constant over time in each population) was relatively similar in POP1 ( $0.33 \pm 0.01$ ), in POP2 ( $0.25 \pm 0.03$ ) and in POP3 ( $0.29 \pm 0.03$ ). Yet, it was higher in POP4 ( $0.43 \pm 0.01$ ) and drastically

increased in POP5 ( $0.71 \pm 0.01$ ). In all five populations, recruitment was highly variable between years (Figure 3). Sex had little effect on recruitment in all the populations and subpopulations (Figure 3, and model selection procedures in Appendix S4).




**FIGURE 4** Effects of density (i.e. population size at  $t - 1$ ) and weather variables on survival (a–e) and recruitment (f–j) in five populations of great crested newt in Europe. In POP1, POP2, POP3 and POP5, four meteorological factors were considered in the analyses: cumulative rainfall during the breeding period (March–May) and the non-activity period (June–February) as well as minimum and maximum monthly temperature during the winter (December–February). As POP4 displays a phenological shift, we considered the four meteorological factors: cumulative rainfall during the breeding period (October–April) and the non-activity period (May–September) as well as minimum and maximum monthly temperature during the non-activity period. The figure shows all the significant relationships (assessed using ANODEVs) between demographic parameters and weather variables. For each relationship, the time-specific estimates of the demographic parameters are fitted against the considered covariate (circles) and lines show the predictions provided by the model including the weather covariate. In the cases where a sex-specific effect was retained, parameter estimates and predictions for males are shown in full circles and full lines; those of females are shown in empty circles and broken lines

### 3.3 | Influence of density dependence and weather on survival and recruitment

Our analyses revealed a weak effect of density dependence and a heterogeneous effect of weather on survival at both subpopulation and population levels (Figure 4)—ANODEV results are summarized in Figure 5, and complete ANODEV outputs can be found in Appendix S3. The only statistically significant effects were found in POP1 (Figure 4). At the subpopulation level, survival was negatively influenced by the cumulative rainfall during the March–May period (RainMM,  $F = 11.67$ ,  $p = 0.003$ ) and during the June–February period in subpopulation POP1.1 (RainJF,  $F = 5.13$ ,  $p = 0.04$ ) but not in POP1.2.

In this subpopulation, we detected a negative effect of population density on survival (density,  $F = 4.73$ ,  $p = 0.04$ ). At the population level, the two weather effects detected in POP1.1 were also detected at the whole population level (RainMM,  $F = 12.97$ ,  $p = 0.002$ ; RainJF,  $F = 5.64$ ,  $p = 0.03$ ) but no effect of population density was found. In POP4, we also detected a negative effect of the cumulative rainfall during the non-activity period (RainMS,  $F = 5.91$ ,  $p = 0.03$ ) on survival.

Our results revealed heterogeneous effects of weather and density dependence on adult recruitment (Figure 4)—ANODEV results are summarized in Figure 5, and complete ANODEV outputs can be found in Appendix S4. At the subpopulation level, we detected a negative density-dependent effect on recruitment in POP1.1



**FIGURE 5** Effects of density (i.e. population size at  $t - 1$ ) and weather variables on population growth survival and recruitment in five populations of great crested newt (POP1 to POP5) in Europe. POP1 and POP5 includes four subpopulations (POP1.1 to POP1.4, and POP5.1 to POP5.4). In POP1, POP2, POP3 and POP5, four meteorological factors were considered in the analyses: cumulative rainfall during the breeding period (RainMM) and the non-activity period (RainJF) as well as minimum (TempDFI) and maximum (TempDFS) monthly temperature during the winter. As POP4 displays a phenological shift, we considered the four meteorological factors: cumulative rainfall during the breeding period (RainOA) and the non-activity period (RainMS) as well as minimum (TempMSI) and maximum (TempMSS) monthly temperature during the non-activity period. The coefficient slope of the relationship between demographic rates and variables is given and appears in bold when the effect is significant (capture-models:  $p$ -value of the ANODEV  $< 0.05$ ; GSS models: the CRI does not include 0). For recruitment and survival, a complete description of the coefficient slopes (and their 95% CI) and the outputs of ANODEVs is provided in Appendices S3 and S4. The complete outputs of GSS models (posterior distribution, coefficient slopes) are provided in Appendix S5

(density,  $F = 6.06$ ,  $p = 0.02$ ) while no effect was detected in POP1.2. In POP5.3, we found a negative relationship between recruitment and the maximum mean temperature during winter (TempDFS,  $F = 8.14$ ,  $p = 0.02$ ) whereas no effect was detected in POP5.4. At the population level, we detected a negative density-dependent effect on recruitment in POP1 (density,  $F = 8.47$ ,  $p = 0.01$ ) and POP4 (density,  $F = 5.93$ ,  $p = 0.03$ ). By contrast, we found a positive effect of density on recruitment in POP3 (density,  $F = 8.30$ ,  $p = 0.03$ ). We detected an effect of weather on recruitment in two populations. In POP5, recruitment was negatively affected by cumulative rainfall during the aquatic period (RainMM,  $F = 7.35$ ,  $p = 0.01$ ); in POP4, the minimum temperature during the non-activity period had a negative impact on recruitment.

### 3.4 | Influence of density dependence and weather on population growth rate

Negative density dependence was a common phenomenon within populations (for the GSS outputs, see Appendix S5). In POP5, we found negative density-dependent effects on population growth rate in the four subpopulations POP5.1 ( $b_1 = -0.49 \pm 0.29$ ), POP5.2 ( $b_1 = -0.36 \pm 0.18$ ), POP5.3 ( $b_1 = -0.56 \pm 0.22$ ) and POP5.4 ( $b_1 = -0.96 \pm 0.39$ ). In POP1, we detected a negative density-dependent effect on population growth in POP1.2 ( $b_1 = -1.06 \pm 0.43$ ) and a trend (i.e. large size effect, but CRI including 0) in subpopulation POP1.1 ( $b_1 = -0.41 \pm 0.29$ ), POP1.3 ( $b_1 = -0.12 \pm 0.08$ ) and POP1.4 ( $b_1 = -0.39 \pm 0.28$ ). By contrast, weather effects were slight and less important (Figure 5). In POP1, cumulative rainfall during the aquatic period negatively affected population growth rate in subpopulation POP1.4 ( $b_2 = -0.55 \pm 0.23$ ). In addition, we detected a trend of a detrimental effect of cumulative rainfall during the non-aquatic period in POP1.2 ( $b_2 = -0.21 \pm 0.18$ ) and POP1.3 ( $b_2 = -0.30 \pm 0.17$ ); the sign was also negative in POP1.1 ( $b_2 = -0.24 \pm 0.23$ ) and POP1.4 ( $b_2 = -0.30 \pm 0.26$ ). In POP5, we only detected trends of weather effects in the subpopulation POP5.1 where population growth was positively affected by the minimum temperature during winter ( $b_2 = 0.16 \pm 0.12$ ) and by rainfall during the activity period ( $b_2 = 0.17 \pm 0.14$ ).

At the population level, density-dependent regulation was also widespread (for the GSS outputs, see Appendix S5). We detected negative effects of population density on population growth in POP4 ( $b_1 = -0.91 \pm 0.46$ ) and POP5 ( $b_1 = -0.41 \pm 0.23$ ). In addition, a similar trend was found in POP1 ( $b_1 = -0.28 \pm 0.24$ ) and POP3 ( $b_1 = -0.39 \pm 0.30$ ). By contrast, we failed to detect any effect of density dependence in POP2 ( $b_1 = -0.06 \pm 0.18$ ). Overall, weather had a negligible influence on population growth (Figure 5).

## 4 | DISCUSSION

We found a low level of demographic synchrony at both local and continental levels in *T. cristatus* populations across western Europe.

Our study also revealed that weather had weak and spatially variable effects on survival, recruitment and population growth rate. In contrast, density dependence was a common phenomenon (at least for population growth rate) in all the populations and subpopulations. Our results suggest that population dynamics more closely depend on habitat-specific factors rather than regional weather fluctuations, resulting in a low synchrony within and between populations.

### 4.1 | Weak and spatially heterogeneous effects of weather on population dynamics

Overall, we found that weather variation has relatively weak effects on demographic rates at the subpopulation and population levels. At the population scale, we detected only three statistically significant relationships between survival and weather factors (cumulative rainfall during the breeding and non-activity periods) in two populations (POP1 and POP4). In POP1, the two relationships (rainfall during the breeding and non-aquatic periods) were also detected in one of the two subpopulations (POP1.1). The negative relationship between survival and rainfall during non-breeding activity (POP1 and POP4) confirms the outcomes of a previous study over a shorter timeframe on this population (Griffiths et al., 2010). High rainfall often results in waterlogged soils, which could impair respiration during overwintering. It could also stimulate newt activity during periods with unsuitable weather conditions (i.e. summer in POP4), resulting in a negative impact on their survival. Moreover, our results indicate that high rainfall during the breeding period also negatively affects adult survival (POP1). As in many other amphibians, high rainfall levels during the reproduction period increase the level of energetically demanding activities related to breeding (breeding migration, mate searching activities, competition for mates; Wells, 2010), which may increase mortality rates. Our findings therefore suggest that these weather effects on adult survival were highly context-dependent and variable across the distribution range of *T. cristatus*. This confirms recent work that highlighted strong context-dependent effects of weather on survival in other amphibian species (Cayuela et al., 2016a, 2017; Muths et al., 2017).

Weather also had a low influence on adult recruitment. Indeed, we detected only two statistically significant relationships between adult recruitment and one weather factor in POP5 and POP4. In POP5, we found a negative relationship between recruitment and cumulative rainfall during the breeding period in year  $y - 1$ . A higher level of energetically demanding activities in immatures, such as natal dispersal during the previous year, might increase mortality rates. Such increases could then result in lower recruitment the following year. We also recorded a negative relationship between recruitment and the minimum monthly temperature during the non-activity period in POP4 and the subpopulation POP5.3. This pattern is in accordance with a previous study in amphibians showing that higher temperatures may increase mortality rates due the depletion of energy reserves during

overwintering (e.g. Reading, 2007). Lower survival of immatures in year  $y - 1$  could then affect recruitment. Like survival, the weather effects on adult recruitment varied across the distribution range of *T. cristatus*.

Finally, GSS models revealed highly marginal effects of weather on population growth rate at either subpopulation or population levels. This suggests that the three weather variables considered in our study that have been previously reported as important drivers for amphibians (e.g. Cayuela et al., 2016a; Muths et al., 2017) have a relatively weak effect on *T. cristatus* population dynamics. It is also congruent with our capture–recapture analyses that showed slight and spatially heterogeneous effects of weather on adult survival and recruitment. The spatial inconsistency of the weather effects within and between-population levels is thus a potential mechanism that might reduce the population synchrony at landscape and regional scales.

## 4.2 | Density dependence, a central driver of population dynamics

Our study revealed that density dependence during the adult stage was a common driver in the dynamics of *T. cristatus* populations. We detected negative effects of density on population growth rates in almost all the subpopulations and the populations, although the detection of density-dependent effects on survival and recruitment was more variable. We only found a negative effect of density on survival in subpopulation POP1.2. Yet, we detected a negative effect of density on adult recruitment in two of the five studied populations (POP1 and POP4). By contrast, we detected a positive effect of density dependence on recruitment in POP3. This population occupies a pond created at the beginning of the sampling, and positive density dependence could result from mate-finding Allee effect (Cayuela, Schmidt, Weinbach, Besnard, & Joly, 2019; Gascoigne, Berec, Gregory, & Courchamp, 2009).

As density effects on adult recruitment and survival varies across populations, it is very likely that density-dependent mechanisms at larval and juvenile stages cause the widespread negative effects of density on population growth. In amphibians, large population size usually results in high larval density in ponds (Beebee, Denton, & Buckley, 1996; Berven, 1990). High density of larvae negatively affects larval growth and survival as well as postmetamorphic performances (Anderson & Whiteman, 2015; Ousterhout & Semlitsch, 2016; Petranks, 1989; Van Buskirk & Smith, 1991). Such density-dependent processes during early life-history stages may lead to a decrease in juvenile recruitment and a reduction of population size at the next generation (Ousterhout & Semlitsch, 2016; Vonesh & De la Cruz, 2002), and can therefore produce the pattern of density-dependent population observed here. Further studies are therefore warranted to investigate how the effects of density-dependent mechanisms at different life stages (i.e. larvae, juveniles and adults) drive population growth in *T. cristatus*, and in amphibians in general. Although frequently difficult to achieve in

the field, increasing capture rates may increase the precision of demographic inferences and avoid the convergence issues encountered in our study when both time-specific survival and recapture rates are considered.

Although density dependence was a widespread phenomenon in *T. cristatus*, the strength of density dependence on population growth differed among populations. Indeed, we failed to detect density dependence in few populations and, when it was different from 0, the coefficient slope (i.e. size effect) describing the density effect varied among populations and subpopulations. This indicates that both geographically distant populations as well as connected subpopulations may have contrasting density dependence structures. In such cases, the differences are likely driven by local environmental characteristics (e.g. trophic resources, interspecific competition in ponds; Griffiths & De Wijer, 1994; Griffiths, Wijer, & May, 1994) and transient population dynamics (i.e. increase or decline over the study period), along with very low dispersal rates (Cayuela, Schmidt, et al., 2019).

## 4.3 | Weak population synchrony within and between populations

Our study highlighted a higher level of population synchrony within populations than between populations. However, population synchrony was overall weak, suggesting a negligible Moran effect, even at a relatively small spatial scale: the correlation coefficients were relatively low among the subpopulations of POP1 (0.28) and POP5 (0.61). A higher ICC in POP5 than in POP1 is likely due to lower Euclidean distances between subpopulations (from 200 to 800 m in POP1 and from 60 to 430 m in POP5). At this spatial scale, the meteorological conditions and the landscape characteristics are expected to be relatively homogeneous. Consequently, the low subpopulation synchrony may result from the effects of local demographic and environmental factors. The contrasting density dependence structures observed among subpopulations in our study system may broadly explain the low subpopulation synchrony in POP1 and POP5. Moreover, local variation in survival and recruitment driven by pond characteristics (Unglaub et al., 2015, 2018) also probably reduces the level of subpopulation synchrony.

At the regional scale, the level of population synchrony was particularly weak (i.e. 0.2 when performing the analysis only at the regional scale, close to 0 when performing the analysis at both the regional and local scale using POP1 and POP5, see Appendix S2). In addition to the mechanisms discussed above, regional climatic variation could further reduce population synchrony. Moreover, the very low level of synchrony could be also caused by inter-population differences in demographic strategies (in amphibians, see for instance Cayuela et al., 2016b). Indeed, our results show broad variation in survival and recruitment rates, several showing 'fast' life histories (low survival and high recruitment, POP1) while others displayed decelerated life histories (high survival and low recruitment, POP2 and POP3).

## 5 | CONCLUSIONS

The effects of weather on *T. cristatus* were weak and highly variable both within and between populations. In contrast, density dependence had a strong influence on the dynamics of *T. cristatus* populations. Our study demonstrated that context-dependent weather effects as well as local variation in density dependence structure are drivers of low population synchrony within and between populations. To date, our study is one of the few that have linked population synchrony, density dependence and heterogeneous demographic responses to weather in natural populations. Although our analyses were focused on one newt species, the low levels of population synchrony reported may be a widespread phenomenon in pond-breeding amphibians in which density dependence (Anderson & Whiteman, 2015; Petranka, 1989; Van Buskirk & Smith, 1991), as well as both aquatic (Morin, 1987; Pearman, 1995; Relyea, 2004) and terrestrial habitat characteristics (Patrick, Harper, Hunter, & Calhoun, 2008) have a strong impact on larval and juvenile survival. By contrast, the effect of climate on demographic rates seems highly context-dependent (Cayuela et al., 2016a; Muths et al., 2017), which should inevitably lead to a weak Moran effect among populations. Moreover, it is likely that low population synchrony is a common demographic feature in many freshwater organisms. Indeed, high population asynchrony has been reported in a broad range of freshwater fishes in which demographic parameters strongly depend on a combination of local aquatic factors and density dependence (Chevalier, Laffaille, Ferdy, & Grenouillet, 2015). In such taxa, a relatively low level of synchrony might have far-reaching consequences for the long-term viability of spatially structured populations. In particular, low population synchrony likely limits the risk of extinction of spatially structured populations, allowing rescue effects between subpopulations and the colonization of patches after local extinction. In addition, it is expected to have important consequences for population response to climate change, by mitigating extinction risk caused by large-scale climatic anomalies. To generalize our findings, we encourage further studies to examine the relative contribution of synchronizing (dispersal, weather variation) and desynchronizing (density, local environmental factors) drivers of population synchrony and dynamics in a broad range of taxa.

## ACKNOWLEDGEMENTS

We thank the many staff, students and volunteers who have contributed to data collection in populations from England over the past 25 years, particularly C. Williams, S. Young, D. Sewell, B. Lewis, A. Wright, P. Walsh and J. Phillips. All surveys in England were carried out under licence from Natural England. We also warmly thank the Conservatoire d'Espaces Naturels Languedoc-Roussillon who collected the data on POP5, especially Thomas Gendre and Pauline Bernard.

## AUTHORS' CONTRIBUTIONS

H.C. and P.J. conceived the ideas and designed methodology; J.W.A., R.A.G., N.Z. and P.P. collected the data; H.C. and J.-P.L. analysed the

data; H.C. led the writing of the manuscript. All authors contributed critically to the drafts and gave final approval for publication.

## DATA AVAILABILITY STATEMENT

Data available from the Dryad Digital Repository: <https://doi.org/10.5061/dryad.573n5tb46> (Cayuela et al., 2020).

## ORCID

Hugo Cayuela  <https://orcid.org/0000-0002-3529-0736>

Pierre Joly  <https://orcid.org/0000-0003-4918-3277>

## REFERENCES

- Anderson, T. L., & Whiteman, H. H. (2015). Non-additive effects of intra- and interspecific competition between two larval salamanders. *Journal of Animal Ecology*, 84, 765–772. <https://doi.org/10.1111/1365-2656.12335>
- Arntzen, J. W., & Hedlund, L. (1990). Fecundity of the newts *Triturus cristatus*, *T. marmoratus* and their natural hybrids in relation to species coexistence. *Ecography*, 13, 325–332. <https://doi.org/10.1111/j.1600-0587.1990.tb00625.x>
- Băncilă, R. I., Ozgul, A., Hartel, T., Sos, T., & Schmidt, B. R. (2015). Direct negative density-dependence in a pond-breeding frog population. *Ecography*, 39, 449–455. <https://doi.org/10.1111/ecog.01584>
- Beebee, T. J., Denton, J. S., & Buckley, J. (1996). Factors affecting population densities of adult natterjack toads *Bufo calamita* in Britain. *Journal of Applied Ecology*, 33, 263–268. <https://doi.org/10.2307/2404748>
- Berven, K. A. (1990). Factors affecting population fluctuations in larval and adult stages of the wood frog (*Rana sylvatica*). *Ecology*, 71, 1599–1608. <https://doi.org/10.2307/1938295>
- Bjørnstad, O. N., Ims, R. A., & Lambin, X. (1999). Spatial population dynamics: Analyzing patterns and processes of population synchrony. *Trends in Ecology & Evolution*, 14, 427–432. [https://doi.org/10.1016/S0169-5347\(99\)01677-8](https://doi.org/10.1016/S0169-5347(99)01677-8)
- Bjørnstad, O. N., Liebhold, A. M., & Johnson, D. M. (2008). Transient synchronization following invasion: Revisiting Moran's model and a case study. *Population Ecology*, 50, 379–389. <https://doi.org/10.1007/s10144-008-0105-5>
- Brooks, S. P., & Gelman, A. (1998). General methods for monitoring convergence of iterative simulations. *Journal of Computational and Graphical Statistics*, 7, 434–455.
- Cayuela, H., Griffiths, R. A., Zakaria, N., Arntzen, J. W., Priol, P., Léna, J.-P., ... Joly, P. (2020). Data from: Drivers of amphibian population dynamics and asynchrony at local and regional scales. *Dryad Digital Repository*, <https://doi.org/10.5061/dryad.573n5tb46>
- Cayuela, H., Arsovski, D., Thirion, J. M., Bonnaire, E., Pichenot, J., Boitaud, S., ... Besnard, A. (2016a). Demographic responses to weather fluctuations are context dependent in a long-lived amphibian. *Global Change Biology*, 22, 2676–2687. <https://doi.org/10.1111/gcb.13290>
- Cayuela, H., Arsovski, D., Thirion, J. M., Bonnaire, E., Pichenot, J., Boitaud, S., ... Besnard, A. (2016b). Contrasting patterns of environmental fluctuation contribute to divergent life histories among amphibian populations. *Ecology*, 97, 980–991. <https://doi.org/10.1890/15-0693.1>
- Cayuela, H., Cruickshank, S. S., Brandt, H., Ozgul, A., & Schmidt, B. R. (2019). Habitat-driven life history variation in an amphibian metapopulation. *Oikos*, 128, 1265–1276. <https://doi.org/10.1111/oik.06286>
- Cayuela, H., Grolet, O., & Joly, P. (2018). Context-dependent dispersal, public information and heterospecific attraction in newts. *Oecologia*, 188, 1069–1080. <https://doi.org/10.1007/s00442-018-4267-3>

- Cayuela, H., Joly, P., Schmidt, B. R., Pichenot, J., Bonnaire, E., Priol, P., ... Besnard, A. (2017). Life history tactics shape amphibians' demographic responses to the North Atlantic Oscillation. *Global Change Biology*, 23, 4620–4638. <https://doi.org/10.1111/gcb.13672>
- Cayuela, H., Schmidt, B. R., Weinbach, A., Besnard, A., & Joly, P. (2019). Multiple density-dependent processes shape the dynamics of a spatially structured amphibian population. *Journal of Animal Ecology*, 88, 164–177. <https://doi.org/10.1111/1365-2656.12906>
- Chambert, T., Pardo, D., Choquet, R., Staszewski, V., McCoy, K. D., Tveraa, T., & Boulinier, T. (2012). Heterogeneity in detection probability along the breeding season in black-legged kittiwakes: Implications for sampling design. *Journal of Ornithology*, 152, 371–380. <https://doi.org/10.1007/s10336-010-0542-8>
- Chevalier, M., Laffaille, P., Ferdy, J. B., & Grenouillet, G. (2015). Measurements of spatial population synchrony: Influence of time series transformations. *Oecologia*, 179, 15–28. <https://doi.org/10.1007/s00442-015-3331-5>
- Choquet, R., Lebreton, J. D., Gimenez, O., Reboulet, A. M., & Pradel, R. (2009). U-CARE: Utilities for performing goodness of fit tests and manipulating CAPTURE–RECAPTURE data. *Ecography*, 32, 1071–1074.
- Choquet, R., Rouan, L., & Pradel, R. (2009). Program E-SURGE: A software application for fitting multilevel models. In D. L. Thomson, E. G. Cooch, & M. J. Conroy (Eds.), *Modeling demographic processes in marked populations* (pp. 845–865). Boston, MA: Springer.
- Church, D. R., Bailey, L. L., Wilbur, H. M., Kendall, W. L., & Hines, J. E. (2007). Iteroparity in the variable environment of the salamander *Ambystoma tigrinum*. *Ecology*, 88, 891–903.
- Crespin, L., Choquet, R., Lima, M., Merritt, J., & Pradel, R. (2008). Is heterogeneity of catchability in capture–recapture studies a mere sampling artifact or a biologically relevant feature of the population? *Population Ecology*, 50, 247–256. <https://doi.org/10.1007/s10144-008-0090-8>
- Cubaynes, S., Pradel, R., Choquet, R., Duchamp, C., Gaillard, J. M., Lebreton, J. D., ... Gimenez, O. (2010). Importance of accounting for detection heterogeneity when estimating abundance: The case of French wolves. *Conservation Biology*, 24, 621–626. <https://doi.org/10.1111/j.1523-1739.2009.01431.x>
- Denoël, M., Dalleur, S., Langrand, E., Besnard, A., & Cayuela, H. (2018). Dispersal and alternative breeding site fidelity strategies in an amphibian. *Ecography*, 41, 1543–1555. <https://doi.org/10.1111/ecog.03296>
- de Valpine, P., & Hastings, A. (2002). Fitting population models incorporating process noise and observation error. *Ecological Monographs*, 72, 57–76. [https://doi.org/10.1890/0012-9615\(2002\)072\[0057:FP-MIPN\]2.0.CO;2](https://doi.org/10.1890/0012-9615(2002)072[0057:FP-MIPN]2.0.CO;2)
- Engen, S., & Sæther, B. E. (2005). Generalizations of the Moran effect explaining spatial synchrony in population fluctuations. *The American Naturalist*, 166, 603–612. <https://doi.org/10.1086/491690>
- Gascoigne, J., Berec, L., Gregory, S., & Courchamp, F. (2009). Dangerously few liaisons: A review of mate-finding Allee effects. *Population Ecology*, 51, 355–372. <https://doi.org/10.1007/s10144-009-0146-4>
- Gelman, A., & Hill, J. (2006). *Data analysis using regression and multilevel/hierarchical models*. Cambridge, UK: Cambridge University Press.
- Griffiths, R. A. (1996). *Newts and salamanders of Europe*. London, UK: Academic Press.
- Griffiths, R. A., & De Wijer, P. (1994). Differential effects of pH and temperature on embryonic development in the British newts (*Triturus*). *Journal of Zoology*, 234, 613–622.
- Griffiths, R. A., De Wijer, P., & May, R. T. (1994). Predation and competition within an assemblage of larval newts (*Triturus*). *Ecography*, 17, 176–181. <https://doi.org/10.1111/j.1600-0587.1994.tb00091.x>
- Griffiths, R. A., Sewell, D., & McCrea, R. S. (2010). Dynamics of a declining amphibian metapopulation: Survival, dispersal and the impact of climate. *Biological Conservation*, 143, 485–491. <https://doi.org/10.1016/j.biocon.2009.11.017>
- Grosbois, V., Gimenez, O., Gaillard, J. M., Pradel, R., Barbraud, C., Clobert, J., ... Weimerskirch, H. (2008). Assessing the impact of climate variation on survival in vertebrate populations. *Biological Reviews*, 83, 357–399. <https://doi.org/10.1111/j.1469-185X.2008.00047.x>
- Grosbois, V., Harris, M. P., Anker-Nilssen, T., McCleery, R. H., Shaw, D. N., Morgan, B. J., & Gimenez, O. (2009). Modeling survival at multi-population scales using mark–recapture data. *Ecology*, 90, 2922–2932. <https://doi.org/10.1890/08-1657.1>
- Hagström, T. (1980). Growth of newts (*Triturus cristatus* and *Triturus vulgaris*) at various ages (Amphibia: Caudata: Salamandridae). *Salamandra*, 16, 248–251.
- Harper, E. B., & Semlitsch, R. D. (2007). Density dependence in the terrestrial life history stage of two anurans. *Oecologia*, 153, 879–889. <https://doi.org/10.1007/s00442-007-0796-x>
- Haynes, K. J., Bjørnstad, O. N., Allstadt, A. J., & Liebhold, A. M. (2013). Geographical variation in the spatial synchrony of a forest-defoliating insect: Isolation of environmental and spatial drivers. *Proceedings of the Royal Society B: Biological Sciences*, 280, 20122373.
- Hedlund, L. (1990). Factors affecting differential mating success in male crested newts, *Triturus cristatus*. *Journal of Zoology*, 220, 33–40. <https://doi.org/10.1111/j.1469-7998.1990.tb04292.x>
- Hugueny, B. (2006). Spatial synchrony in population fluctuations: Extending the Moran theorem to cope with spatially heterogeneous dynamics. *Oikos*, 115, 3–14. <https://doi.org/10.1111/j.2006.0030-1299.14686.x>
- Ims, R. A., & Andreassen, H. P. (2000). Spatial synchronization of vole population dynamics by predatory birds. *Nature*, 408, 194–196. <https://doi.org/10.1038/35041562>
- IUCN. (2020). The IUCN red list of threatened species. Version 2020–1. Retrieved from <https://www.iucnredlist.org>
- Jehle, R., & Arntzen, J. W. (2000). Post-breeding migrations of newts (*Triturus cristatus* and *T. marmoratus*) with contrasting ecological requirements. *Journal of Zoology*, 251, 297–306. <https://doi.org/10.1111/j.1469-7998.2000.tb01080.x>
- Jehle, R., Thiesmeier, B., & Foster, J. (2011). *The crested newt. A dwindling pond-dweller*. Bielefeld, Germany: Laurenti-Verlag.
- Jeyam, A., McCrea, R. S., Bregnballe, T., Frederiksen, M., & Pradel, R. (2018). A test of positive association for detecting heterogeneity in capture for capture–recapture data. *Journal of Agricultural, Biological and Environmental Statistics*, 23, 1–19. <https://doi.org/10.1007/s13253-017-0315-4>
- Kellner, K. (2018). A Wrapper Around 'rjags' to Streamline 'JAGS' Analyses. Retrieved from <https://cran.r-project.org/web/packages/jagsUI>
- Kendall, B. E., Bjørnstad, O. N., Bascompte, J., Keitt, T. H., & Fagan, W. F. (2000). Dispersal, environmental correlation, and spatial synchrony in population dynamics. *The American Naturalist*, 155, 628–636. <https://doi.org/10.1086/303350>
- Kéry, M., & Schaub, M. (2011). *Bayesian population analysis using WinBUGS: A hierarchical perspective*. San Diego, CA: Academic Press.
- Koenig, W. D. (2001). Spatial autocorrelation and local disappearances in wintering North American birds. *Ecology*, 82, 2636–2644. [https://doi.org/10.1890/0012-9658\(2001\)082\[2636:SAALDI\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2001)082[2636:SAALDI]2.0.CO;2)
- Koenig, W. D. (2002). Global patterns of environmental synchrony and the Moran effect. *Ecography*, 25, 283–288. <https://doi.org/10.1034/j.1600-0587.2002.250304.x>
- Koenig, W. D., & Liebhold, A. M. (2016). Temporally increasing spatial synchrony of North American temperature and bird populations. *Nature Climate Change*, 6, 614–617. <https://doi.org/10.1038/nclimate2933>
- Korpimäki, E., Norrdahl, K., Huitu, O., & Klemola, T. (2005). Predator-induced synchrony in population oscillations of coexisting small mammal species. *Proceedings of the Royal Society B: Biological Sciences*, 272, 193–202. <https://doi.org/10.1098/rspb.2004.2860>
- Liebhold, A., Koenig, W. D., & Bjørnstad, O. N. (2004). Spatial synchrony in population dynamics. *Annual Review of Ecology, Evolution, and*

- Systematics*, 35, 467–490. <https://doi.org/10.1146/annurev.ecolsys.34.011802.132516>
- Miaud, C., Joly, P., & Castanet, J. (1993). Variation in age structures in a subdivided population of *Triturus cristatus*. *Canadian Journal of Zoology*, 71, 1874–1879. <https://doi.org/10.1139/z93-267>
- Moran, P. A. P. (1953). The statistical analysis of the Canadian lynx cycle. II. Synchronization and meteorology. *Australian Journal of Zoology*, 1, 291–298. <https://doi.org/10.1071/ZO9530291>
- Morin, P. J. (1987). Predation, breeding asynchrony, and the outcome of competition among treefrog tadpoles. *Ecology*, 68, 675–683. <https://doi.org/10.2307/1938473>
- Muths, E., Chambert, T., Schmidt, B. R., Miller, D. A. W., Hossack, B. R., Joly, P., ... Fisher, R. N. (2017). Heterogeneous responses of temperate-zone amphibian populations to climate change complicates conservation planning. *Scientific Reports*, 7, 17102. <https://doi.org/10.1038/s41598-017-17105-7>
- Nakagawa, S., Johnson, P. C., & Schielzeth, H. (2017). The coefficient of determination  $R^2$  and intra-class correlation coefficient from generalized linear mixed-effects models revisited and expanded. *Journal of the Royal Society Interface*, 14, 20170213.
- Ousterhout, B. H., & Semlitsch, R. D. (2016). Non-additive response of larval ringed salamanders to intraspecific density. *Oecologia*, 180, 1137–1145. <https://doi.org/10.1007/s00442-015-3516-y>
- Patrick, D. A., Harper, E. B., Hunter, M. L., & Calhoun, A. J. (2008). Terrestrial habitat selection and strong density-dependent mortality in recently metamorphosed amphibians. *Ecology*, 89, 2563–2574. <https://doi.org/10.1890/07-0906.1>
- Pearman, P. B. (1995). Effects of pond size and consequent predator density on two species of tadpoles. *Oecologia*, 102, 1–8. <https://doi.org/10.1007/BF00333303>
- Peltonen, M., Liebhold, A. M., Bjørnstad, O. N., & Williams, D. W. (2002). Spatial synchrony in forest insect outbreaks: Roles of regional stochasticity and dispersal. *Ecology*, 83, 3120–3129. [https://doi.org/10.1890/0012-9658\(2002\)083\[3120:SSIFIO\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2002)083[3120:SSIFIO]2.0.CO;2)
- Péron, G., Crochet, P. A., Choquet, R., Pradel, R., Lebreton, J. D., & Gimenez, O. (2010). Capture–recapture models with heterogeneity to study survival senescence in the wild. *Oikos*, 119, 524–532. <https://doi.org/10.1111/j.1600-1706.2009.17882.x>
- Petranks, J. W. (1989). Density-dependent growth and survival of larval *Ambystoma*: Evidence from whole-pond manipulations. *Ecology*, 70, 1752–1767.
- Pledger, S. (2000). Unified maximum likelihood estimates for closed capture–recapture models using mixtures. *Biometrics*, 56, 434–442. <https://doi.org/10.1111/j.0006-341X.2000.00434.x>
- Plummer, M. (2003). JAGS: A program for analysis of Bayesian graphical models using Gibbs sampling. In *Proceedings of the 3rd international workshop on distributed statistical computing* (Vol. 124, p. 125).
- Pradel, R. (1996). Utilization of capture-mark-recapture for the study of recruitment and population growth rate. *Biometrics*, 52, 703–709. <https://doi.org/10.2307/2532908>
- Ranta, E., Kaitala, V., Lindström, J., & Linden, H. (1995). Synchrony in population dynamics. *Proceedings of the Royal Society of London B: Biological Sciences*, 262, 113–118.
- Reading, C. J. (2007). Linking global warming to amphibian declines through its effects on female body condition and survivorship. *Oecologia*, 151, 125–131. <https://doi.org/10.1007/s00442-006-0558-1>
- Relyea, R. A. (2004). Growth and survival of five amphibian species exposed to combinations of pesticides. *Environmental Toxicology and Chemistry: An International Journal*, 23, 1737–1742. <https://doi.org/10.1897/03-493>
- Royama, T. (2005). Moran effect on nonlinear population processes. *Ecological Monographs*, 75, 277–293. <https://doi.org/10.1890/04-0770>
- Schaub, M., Hirschheydt, J., & Gruebler, M. U. (2015). Differential contribution of demographic rate synchrony to population synchrony in barn swallows. *Journal of Animal Ecology*, 84, 1530–1541. <https://doi.org/10.1111/1365-2656.12423>
- Schaub, M., & Kéry, M. (2012). Combining information in hierarchical models improves inferences in population ecology and demographic population analyses. *Animal Conservation*, 15, 125–126. <https://doi.org/10.1111/j.1469-1795.2012.00531.x>
- Selwood, K. E., McGeoch, M. A., & Mac Nally, R. (2015). The effects of climate change and land-use change on demographic rates and population viability. *Biological Reviews*, 90, 837–853. <https://doi.org/10.1111/brv.12136>
- Trenham, P. C., Koenig, W. D., Mossman, M. J., Stark, S. L., & Jagger, L. A. (2003). Regional dynamics of wetland-breeding frogs and toads: Turnover and synchrony. *Ecological Applications*, 13, 1522–1532. <https://doi.org/10.1890/02-5206>
- Unglaub, B., Steinfartz, S., Drechsler, A., & Schmidt, B. R. (2015). Linking habitat suitability to demography in a pond-breeding amphibian. *Frontiers in Zoology*, 12, 9. <https://doi.org/10.1186/s12983-015-0103-3>
- Unglaub, B., Steinfartz, S., Kühne, D., Haas, A., & Schmidt, B. R. (2018). The relationships between habitat suitability, population size and body condition in a pond-breeding amphibian. *Basic and Applied Ecology*, 27, 20–29. <https://doi.org/10.1016/j.baae.2018.01.002>
- Van Buskirk, J., & Smith, D. C. (1991). Density-dependent population regulation in a salamander. *Ecology*, 72, 1747–1756.
- Vonesh, J. R., & De la Cruz, O. (2002). Complex life cycles and density dependence: Assessing the contribution of egg mortality to amphibian declines. *Oecologia*, 133, 325–333. <https://doi.org/10.1007/s00442-002-1039-9>
- Wells, K. D. (2010). *The ecology and behavior of amphibians*. Chicago, IL: University of Chicago Press.
- Ylikarjula, J., Alaja, S., Laakso, J., & Tesar, D. (2000). Effects of patch number and dispersal patterns on population dynamics and synchrony. *Journal of Theoretical Biology*, 207, 377–387. <https://doi.org/10.1006/jtbi.2000.2181>
- Zakaria, N. B. (2018). *Long-term population ecology of the great crested newt in Kent* (PhD thesis). University of Kent.

## SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

**How to cite this article:** Cayuela H, Griffiths RA, Zakaria N, et al. Drivers of amphibian population dynamics and asynchrony at local and regional scales. *J Anim Ecol*. 2020;89:1350–1364. <https://doi.org/10.1111/1365-2656.13208>